



# Strangeness Nuclear Physics with PANDA

Michael Papenbrock

Uppsala University



Super-FRS Coll. Meeting  
Walldorf  
28. April 2023

# Outline

- Motivation
- The PANDA detector
- Antihyperons in nuclei
- Neutron skin measurements
- Hyperatom spectroscopy
- Double-strange hypernuclei
- Summary





# Hyperon Puzzle


- Neutron stars among the most enigmatic objects in the Universe
- Extreme conditions: Masses of to 2 solar masses, but radii as low as approx. 10 km
- Formation and fate determined by Equation of State (EoS)
- Gravity, strong force, and Pauli principle compete
  - Rapid increase of chemical potential at centre
  - Conversion of nucleons to hyperons energetically favourable
  - Relief of Fermi pressure softens EoS, reducing maximum mass to 1.4 solar masses
  - Contradiction → Hyperon Puzzle

# Experimental approaches to hyperon few-body interaction

- Hypernuclei
  - 2+3-body forces
  - High-precision  $\gamma$ -ray spectroscopy
  - Spin-dependent forces
  - Significant contribution to solving hyperon puzzle
- Hyperon femtoscopy
  - Low-energy scattering parameters
  - NY and YY pairs
  - E.g. ALICE and HADES




# Facility for Antiproton and Ion Research (FAIR)




# High Energy Storage Ring (HESR)

- Anti-protons with  $1.5 < p_{beam} < 15 \text{ GeV}/c$
- Internal targets
  - Cluster-jet and pellet ( $\bar{p}p$ )
  - Foils ( $\bar{p}A$ )
- Luminosity
  - Design  $\sim 2 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
  - Phase One  $\sim 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$
- Quasi-continuous beam





# PANDA – full setup









# Antihyperons in nuclei

- Measure asymmetry of  $\bar{Y}$  and  $Y$



$$\alpha_T = \frac{p_T(Y) - p_T(\bar{Y})}{p_T(Y) + p_T(\bar{Y})}, \alpha_L = \frac{p_L(Y) - p_L(\bar{Y})}{p_L(Y) + p_L(\bar{Y})}$$

- Momentum asymmetries relate to  $\bar{Y}$  interaction potential
- Possible within an hour of data taking
- PANDA unique



# Probing neutron skin




- Many different approaches
  - Hadronic probes
  - Electromagnetic probes
  - Weak interaction
  - Astrophysical observations
- The interesting case of  $^{48}\text{Ca}$ 
  - Measure skin difference to  $^{40}\text{Ca}$
  - Ab-initio calculations possible




# Probing neutron skin

- $\Lambda\bar{\Lambda}$  only in  $\bar{p}p$ ,  $\Sigma^-\bar{\Lambda}$  only in  $\bar{p}n$
- Double ratio of probabilities  

$$DR = \frac{p_{\Sigma^-\bar{\Lambda}}^{II}/p_{\Lambda\bar{\Lambda}}^{II}}{p_{\Sigma^-\bar{\Lambda}}^I/p_{\Lambda\bar{\Lambda}}^I} = \frac{1+p_{abs}}{1-p_{abs}}$$
- $p_{abs}$ : antiproton absorption probability, related to integrated skin density
- Study evolution of neutron skin thickness for isotope chains, e.g.  $^{129-136}\text{Xe}$
- Adds to systematic uncertainties of hyperatom observables
- Preprint: arxiv:2209.03875






# Hyperatoms and hypernuclei





# Hyperatoms and hypernuclei




- Target positioning resolution:  $5\mu m$
- Repeatability:  $\pm 18\mu m$  (14000 measurements, requirement:  $300\mu m$ )
- Preprint: arxiv:2303.13359






# Hyperatoms and hypernuclei





# X-ray spectroscopy of $\Xi^-$ hyperatoms

- Observe nuclear cascade
- Shift of low atomic level sensitive to  $\Xi A$  potential near the nuclear surface
- Different neutron/proton content for different nuclei  $\rightarrow$  isospin dependence of  $\Xi A$  force
- PANDA unique



# Sensitivity to nuclear structure


- Systematic uncertainties due to uncertainty of neutron skin thickness
- Best know nucleus:  $^{208}\text{Pb}$





# $\Lambda\Lambda$ hypernuclei

- Active secondary target
  - Boron- $\mu$ Strip sandwich
- Form excited  $\Lambda\Lambda$  hypernuclei by  $\Xi^-$  capture
- Measure  ${}_{\Lambda\Lambda}X$   $\gamma$ -transitions and momentum correlations
- Explore structure of light, double-strange hypernuclei
- Study few-body forces in baryonic matter



Nuclei	$E_x$ (MeV)	$J^P$	production probability				
			${}^9\text{Be}$	${}^{10}\text{B}$	${}^{11}\text{B}$	${}^{\text{nat}}\text{B}$	${}^{12}\text{C}$
${}_{\Lambda\Lambda}^4\text{H}$	0.0	$1^+$	0.00866	0.02410	0.00000	0.00472	0.0000
${}_{\Lambda\Lambda}^5\text{H}$	0.0	$1^+$	0.02120	0.02209	0.03199	0.03005	0.00633
${}_{\Lambda\Lambda}^5\text{He}$	0.0	$1^+$	0.00000	0.00330	0.00000	0.00065	0.00000
${}_{\Lambda\Lambda}^6\text{He}$	0.0	$0^+$	0.02350	0.03175	0.00977	0.01408	0.03304
${}_{\Lambda\Lambda}^7\text{He}$	0.0	$1^+$	0.10201	0.03038	0.04407	0.04139	0.00649
${}_{\Lambda\Lambda}^8\text{He}$	0.0	$0^+$	0.01880	0.00445	0.00490	0.00481	0.00000
${}_{\Lambda\Lambda}^8\text{He}$	1.80	$2^+$	0.08201	0.01846	0.02351	0.02252	0.00000
${}_{\Lambda\Lambda}^9\text{He}$	0.0	$1^+$	0.00426	0.00017	0.00292	0.00238	0.00000
${}_{\Lambda\Lambda}^9\text{He}$	2.92	$1^+$	0.01859	0.00021	0.00435	0.00354	0.00000
${}_{\Lambda\Lambda}^7\text{Li}$	0.0	$1^+$	0.00016	0.00635	0.00000	0.00124	0.00000
${}_{\Lambda\Lambda}^8\text{Li}$	0.0	$1^+$	0.01055	0.01991	0.00233	0.00578	0.00209
${}_{\Lambda\Lambda}^8\text{Li}$	1.36	$3^+$	0.01998	0.03976	0.00212	0.00950	0.00150
${}_{\Lambda\Lambda}^9\text{Li}$	5.63	$2^+$	0.00617	0.01747	0.00000	0.00342	0.00000
${}_{\Lambda\Lambda}^9\text{Li}$	0.0	$1^+$	0.02199	0.03041	0.03948	0.03770	0.02574
${}_{\Lambda\Lambda}^9\text{Li}$	0.73	$1^+$	0.01079	0.01452	0.01803	0.01734	0.01236
${}_{\Lambda\Lambda}^9\text{Li}$	4.55	$1^+$	0.03997	0.04398	0.04528	0.04502	0.04207
${}_{\Lambda\Lambda}^9\text{Li}$	5.96	$1^+$	0.02870	0.02975	0.02907	0.02920	0.02864
${}_{\Lambda\Lambda}^{10}\text{Li}$	0.0	$2^+$	0.00000	0.00702	0.03799	0.03192	0.00000
${}_{\Lambda\Lambda}^{10}\text{Li}$	0.98	$1^+$	0.00000	0.00404	0.02138	0.01798	0.00000
${}_{\Lambda\Lambda}^{10}\text{Li}$	2.255	$3^+$	0.00000	0.00929	0.04422	0.03737	0.00000
${}_{\Lambda\Lambda}^9\text{Be}$	0.0	$1^+$	0.00000	0.00497	0.00003	0.00100	0.00000
${}_{\Lambda\Lambda}^9\text{Be}$	0.71	$1^+$	0.00000	0.00227	0.00001	0.00045	0.00000

# Summary

- PANDA excellent tool for hypernuclear physics from the start
- Explore strong interaction in the nuclear periphery
- Observables relate to equation of state of neutron stars
- Key topics
  - Antihyperon potential in cold baryonic matter
  - Hyperatoms:  $\Xi^-$  potential in neutron-rich environments
  - Structure of  $\Lambda\Lambda$  hypernuclei



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 824093.




Thank you for your attention!






# PANDA – Phase One setup



# Prospects

- JPARC
  - Kaon beam
  - Extended target (cm)
  - $\frac{\rho_n}{\rho_p} \sim 1$
- PANDA
  - Stored antiproton beam
  - Thin secondary target (mm)
  - $\frac{\rho_n}{\rho_p} \sim 2$
- PANDA has unique explanatory potential





# PANDA is a strangeness factory

- New simulation studies of single- and double-strange hyperons\*
  - Exclusive measurements of
    - $\bar{p}p \rightarrow \bar{\Lambda}\Lambda, \Lambda \rightarrow p\pi^-, \bar{\Lambda} \rightarrow \bar{p}\pi^+$
    - $\bar{p}p \rightarrow \bar{\Sigma}^0\Lambda, \Lambda \rightarrow p\pi^-, \bar{\Sigma}^0 \rightarrow \bar{\Lambda}\gamma, \bar{\Lambda} \rightarrow \bar{p}\pi^+$
    - $\bar{p}p \rightarrow \bar{\Xi}^+\Xi^-, \Xi^- \rightarrow \Lambda\pi^-, \Lambda \rightarrow p\pi^-, \bar{\Xi}^+ \rightarrow \bar{\Lambda}\pi^+, \bar{\Lambda} \rightarrow \bar{p}\pi^+$
  - Ideal pattern recognition and PID
  - Background using Dual Parton Model

$p_{beam}$ (GeV/c)	Reaction	$\sigma(\mu\text{b})$	$\varepsilon(\%)$	Rate ( $\text{s}^{-1}$ ) @ $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$	S/B	Events / day
1.64	$\bar{p}p \rightarrow \bar{\Lambda}\Lambda$	64.0	16.0	44	114	$3.8 \cdot 10^6$
1.77	$\bar{p}p \rightarrow \bar{\Sigma}^0\Lambda$	10.9	5.3	2.4	>11**	207000
6.0	$\bar{p}p \rightarrow \bar{\Sigma}^0\Lambda$	20	6.1	5.0	21	432000
4.6	$\bar{p}p \rightarrow \bar{\Xi}^+\Xi^-$	$\sim 1$	8.2	0.3	274	26000
7.0	$\bar{p}p \rightarrow \bar{\Xi}^+\Xi^-$	$\sim 0.3$	7.9	0.1	65	86000

\*\* 90% C.L.

\*By W. Ikegami Andersson (PhD thesis, Uppsala 2020) and G. Perez Andrade (Master thesis, Uppsala 2019)


# Components in EoS

- Strong interaction in the confinement domain
- Three-body forces
  - Nucleon-nucleon-hyperon (NNY)
  - Nucleon-hyperon-hyperon (NYY)
    - Could be repulsive → counteract softening of EoS
- Two-body force
  - Nucleon-hyperon (NY)
  - Hyperon-hyperon (YY)
    - Repulsive core would stiffen EoS or make hyperon presence energetically unfavourable
- Data on hyperon interaction too scarce to constrain theoretical models





# HADES / PANDA Phase 0



- Next beam time: p+p@4.5 GeV
- Opportunity for YN and YY interaction studies
- High sensitivity in low energy region

