Feasibility Study of $Zc(3900)$ and $Zcs(3985)$ in $\bar{p}p$ with the PANDA Detector

Ali YILMAZ & Haluk DENİZLİ

PANDA Collaboration Meeting 23/1

March 6-10, 2023

Intro

- The charmonium like state $Zc^{\pm}(3900)$ was observed by the BESIII [1] and Belle [2] collaborations in the $\pi^{\pm}J/\psi$ invariant mass spectrum of $e^+e^- \rightarrow \pi^+\pi^-$ at $\sqrt{s}$=4.26 GeV in 2013 and then confirmed by CLEO-c collaboration in the same process at s $\sqrt{s}$=4.17 GeV [3], which makes Zc(3900) the first confirmed charged charmonium like state.

- The Zcs(3985)– was observed in π–J/ψ invariant mass distribution in the study of e+e– → π+π–J/ψ at BESIII and Belle experiments *[M. Ablikim et al., C. Z. Yuan et al.].*

- [1] M. Ablikim et al. (BESIII Collaboration), Phys. Rev. Lett. 126, 102001 (2021)

- [1] M. Ablikim et al. (BESIII Collaboration), Phys. Rev. Lett. 110, 252001 (2013).
- [2] Z. Q. Liu et al. (Belle Collaboration), Phys. Rev. Lett. 110, 252002 (2013); 111, 019901(E) (2013).
- [3] T. Xiao, S. Dobbs, A. Tomaradze, and K. K. Seth, Phys. Lett. B 727, 366 (2013).

Outline

- Zc(3900) & Zcs(3985)
  - Event Generation
  - Reconstruction & Analysis
  - Background
- Summary

Event Generation

- 2 million events at each
- $P_{\bar{p}} = 8.5454$ GeV/c (at resonance of $\psi_{(4260)}$)
- assuming the branching ratio of 100% for $Zc(3900) \rightarrow \pi + J/\psi$
- Mass of $Zc(3900)^{\pm}$: $m_{Zc(3900)}$ = 3887.2 ± 2.3 [Mev/c^2]
- Width: $\Gamma_{Zc(3900)}$ = 28.2 ± 2.6 [Mev/c^2]



- 1.99 million events at each
- $P_{\bar{p}} = 15$ GeV/c (at max of PANDA)
- assuming the branching ratio of 100% for $Zcs(3985) \rightarrow K + J/\psi$
- Mass of $Zcs(3985)^{\pm}$: $m_{Zcs(3985)}$ = 3982.5 ± 2.3 [Mev/c^2]
- Width: Γ = 12.8 ± 3.0 [Mev/c^2]



Reconstruction

Production and Reco:

- Simulation of transport through the detector
  - Production & Reco: Using FairSoft jun19p2 / FairRoot v18.2.1 / PandaRoot dev:
  - Analysis: Using FairSoft apr22 / FairRoot v18.6.8 / PandaRoot dev
- Transport and reconstruction of particles is done with the PandaRoot framework
- Follow the decay tree
- Best PID algorithm is used (MuonBestPlus for μ^+,…)

Analysis

Reconst. Final States efficiencies

- Used decay pattern recognition and “best” particle identification (PID)

Zc

- Reconstructed FS: $\mu^-, \mu^+, \pi^-, \pi^+$

Particle type	ε[%]
μ^+	95.47
μ^-	94.19
π^+	83.46
π^-	79.07

$\bar{p}p \to Zc(3900)^+\pi^-, (Zc(3900)^+ \to J/\psi\pi^+, (J/\psi \to \mu^+\mu^-))$

Particle type	ε[%]
e^+	89.48
e^-	85.41
π^+	83.34
π^-	78.89

$\bar{p}p \to Zc(3900)^+\pi^-, (Zc(3900)^+ \to J/\psi\pi^+, (J/\psi \to e^+e^-))$

Particle type	ε[%]
μ^+	96.61
μ^-	94.11
π^+	78.77
π^-	83.39

$\bar{p}p \to Zc(3900)^-\pi^+, (Zc(3900)^- \to J/\psi\pi^-, (J/\psi \to \mu^+\mu^-))$

Particle type	ε[%]
e^+	89.64
e^-	85.32
π^+	78.57
π^-	83.2

$\bar{p}p \to Zc(3900)^-\pi^+, (Zc(3900)^- \to J/\psi\pi^-, (J/\psi \to e^+e^-))$

Zcs

- Reconstructed FS: μ^+, μ^-, K^+, K^-

Particle type	ε[%]
μ^+	99.84
μ^-	94.12
K^+	77.2
K^-	79.39

$\bar{p}p \to K^- Z_{cs}(3985)^+, (Z_{cs}(3985)^+ \to K^+J/\psi), (J/\psi \to \mu^+\mu^-))$

Particle type	ε[%]
e^+	90.8
e^-	82.36
K^+	76.95
K^-	79.16

$\bar{p}p \to K^- Z_{cs}(3985)^+, (Z_{cs}(3985)^+ \to K^+J/\psi), (J/\psi \to e^+e^-))$

Particle type	ε[%]
μ^+	94.91
μ^-	94.15
K^+	80.86
K^-	75.61

$\bar{p}p \to K^+ Z_{cs}(3985)^-, (Z_{cs}(3985)^- \to K^-J/\psi), (J/\psi \to \mu^+\mu^-))$

Particle type	ε[%]
e^+	82.74
e^-	82.43
K^+	80.78
K^-	75.4

$\bar{p}p \to K^+ Z_{cs}(3985)^-, (Z_{cs}(3985)^- \to K^-J/\psi), (J/\psi \to e^+e^-))$

Analysis

Reconstruction of FS: u+

Zc

pt vs pz



P vs Θ



dP vs p



dP/p vs p













Analysis

Reconstruction of FS: u+

Zcs

pt vs pz



P vs Θ



d*P vs p*



dP/*p vs p*













Analysis

Momentum Resolutions

Zc

Reconstructed FS: $\mu^+, \mu^-, \pi^+, \pi^-$

Particle type	*dp/p* [%]
μ^+	1.158
μ^-	1.154
π^+	1.138
π^-	1.136

$\bar{p}p \rightarrow Zc(3900)^{\mp}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	*dp/p* [%]
e+	1.426
e-	1.403
π+	1.146
π-	1.134

$\bar{p}p \rightarrow Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow e^{+}e^{-}))$

Particle type	*dp/p* [%]
μ+	1.167
μ-	1.155
π+	1.123
π-	1.152

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	*dp/p* [%]
e+	1.447
e-	1.435
π+	1.121
π-	1.151

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow e^{+}e^{-}))$

Zcs

Reconstructed FS: μ^+, μ^-, K^+, K^-

Particle type	dp/p [%]
μ^+	0.8109
μ^-	0.8095
K^+	0.7916
K^-	0.8464

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
e^+	0.6003
e^-	0.482
K^+	0.4194
K^-	0.3626

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Particle type	dp/p [%]
μ^+	0.8257
μ^-	0.7994
K^+	0.8447
K^-	0.783

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
e^+	1.209
e^-	1.223
K^+	0.8424
K^-	0.7857

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Analysis

Reconstruction of Resonance State : J/ψ

- Invariant mass cut on $\mu^+\mu^-$ (e^+e^-) to select J/ψ cands $m_{J/\psi}$: (3.0969 ± 0.5) GeV/c^2
- Perform RhoDecayTreeFitter fit
- Select candidate with DecayTree fit prob > 0.01

Analysis

Resonance States: J/ψ

- Reconstructed: efficiency

Zc

Particle type	ε[%]
J/Ψ	67.29

$\bar{p}p \rightarrow Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	ε[%]
J/Ψ	30.44

$\bar{p}p \rightarrow Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow e^{+}e^{-}))$

Particle type	ε[%]
J/Ψ	67.26

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	ε[%]
J/Ψ	30.47

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow e^{+}e^{-}))$

Zcs

Particle type	ε[%]
J/Ψ	67.37

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	ε[%]
J/Ψ	27.97

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Particle type türü	ε[%]
J/Ψ	67.24

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	ε[%]
J/Ψ	27.94

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Analysis

Resonance States: J/ψ

- Reconstructed: m, chi2, prob

Zc







Zcs







Analysis

Resonance States: Momentum Distributions

Zc

- Reconstructed: J/ψ



















Analysis

Resonance States: Momentum Distributions

Zcs

- Reconstructed: J/ψ



















Analysis

Reconst. Resonance States: J/ψ

- Reconstructed: Momentum Resolution

Zc

Particle type	d*p/p[%]*
J/Ψ	1.032

$\bar{p}p \rightarrow Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	*dp/p[%]*
J/Ψ	1.336

$\bar{p}p \rightarrow Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \rightarrow J/\psi\pi^{+}, (J/\psi \rightarrow e^{+}e^{-}))$

Particle type	*dp/p[%]*
J/Ψ	1.028

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	*dp/p[%]*
J/Ψ	1.365

$\bar{p}p \rightarrow Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \rightarrow J/\psi\pi^{-}, (J/\psi \rightarrow e^{+}e^{-}))$

Zcs

Particle type	dp/p [%]
J/Ψ	0.6985

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
J/Ψ	1.145

$\bar{p}p \rightarrow K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \rightarrow K^{+}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Particle type	dp/p [%]
J/Ψ	0.6956

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
J/Ψ	1.126

$\bar{p}p \rightarrow K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \rightarrow K^{-}J/\psi), (J/\psi \rightarrow e^{+}e^{-}))$

Analysis

Reconst. Resonance States: J/ψ

- Reconstructed: m, mDiff

$M_{J/\psi^{evt.pdl}}$: 3.09690 GeV/c²

Zc



$M_{J/\psi^{reco}}$: 3.09746 GeV/c²



Zcs



$M_{J/\psi^{reco}}$: 3.0968 GeV/c²



VoigtFit:: quadratic Background + relativistic BW

Analysis

Reconst. Resonance States:

- Combine J/ψ and π

Zc

- Mass cut with window $m_{Zc(3900)}$: [3.8872 ± 0.5] GeV/c²

Zcs

- Mass cut with window $m_{Zcs(3985)}$: [3.9825 ± 0.5] GeV/c²

- Perform RhoDecayTreeFitter fit
- Select candidate with DecayTree fit prob > 0.01

Analysis

Reconst. Resonance States

- Reconstructed: efficiency

Zc

Particle type	ε [%]
Zc(3900)	45.87
$\bar{p}p \to Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \to J/\psi\pi^{+}, (J/\psi \to \mu^{+}\mu^{-}))$	

Particle type	ε [%]
Zc(3900)	19.98
$\bar{p}p \to Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \to J/\psi\pi^{+}, (J/\psi \to e^{+}e^{-}))$	

Particle type	ε [%]
Zc(3900)	45.94
$\bar{p}p \to Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \to J/\psi\pi^{-}, (J/\psi \to \mu^{+}\mu^{-}))$	

Particle type	ε [%]
Zc(3900)	20.02
$\bar{p}p \to Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \to J/\psi\pi^{-}, (J/\psi \to e^{+}e^{-}))$	

Zcs

Particle type	ε[%]
Zcs(3985)	45.44
$\bar{p}p \to K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \to K^{+}J/\psi), (J/\psi \to \mu^{+}\mu^{-}))$	

Particle type	ε[%]
Zcs(3985)	18.31
$\bar{p}p \to K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \to K^{+}J/\psi), (J/\psi \to e^{+}e^{-}))$	

Particle type	ε[%]
Zcs(3985)	44.31
$\bar{p}p \to K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \to K^{-}J/\psi), (J/\psi \to \mu^{+}\mu^{-}))$	

Particle type	ε[%]
Zcs(3985)	17.88
$\bar{p}p \to K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \to K^{-}J/\psi), (J/\psi \to e^{+}e^{-}))$	

Analysis

Reconst. Resonance States

- Reconstructed: momentum distributions

Zc















Analysis

Reconst. Resonance States

- Reconstructed: momentum distributions

Zcs















Analysis

Reconst. Resonance States:

- Reconstructed: momentum resolutions

Zc

Particle type	*dp/p* [%]
Zc(3900)	0.9954

$\bar{p}p \to Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \to J/\psi\pi^{+}, (J/\psi \to \mu^{+}\mu^{-}))$

Particle type	*dp/p*[%]
Zc(3900)	1.299

$\bar{p}p \to Zc(3900)^{+}\pi^{-}, (Zc(3900)^{+} \to J/\psi\pi^{+}, (J/\psi \to e^{+}e^{-}))$

Particle type	*dp/p*[%]
Zc(3900)	0.9976

$\bar{p}p \to Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \to J/\psi\pi^{-}, (J/\psi \to \mu^{+}\mu^{-}))$

Particle type	*dp/p* [%]
Zc(3900)	1.293

$\bar{p}p \to Zc(3900)^{-}\pi^{+}, (Zc(3900)^{-} \to J/\psi\pi^{-}, (J/\psi \to e^{+}e^{-}))$

Zcs

Particle type	dp/p [%]
Zcs(3985)	0.7922

$\bar{p}p \to K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \to K^{+}J/\psi), (J/\psi \to \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
Zcs(3985)	0.7892

$\bar{p}p \to K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \to K^{-}J/\psi), (J/\psi \to \mu^{+}\mu^{-}))$

Particle type	dp/p [%]
Zcs(3985)	1.174

$\bar{p}p \to K^{-}Z_{cs}(3985)^{+}, (Z_{cs}(3985)^{+} \to K^{+}J/\psi), (J/\psi \to e^{+}e^{-}))$

Particle type	dp/p [%]
Zcs(3985)	1.177

$\bar{p}p \to K^{+}Z_{cs}(3985)^{-}, (Z_{cs}(3985)^{-} \to K^{-}J/\psi), (J/\psi \to e^{+}e^{-}))$

Analysis

Reconst. Resonance States

- Reconstructed: m, mDiff

Zc





$M_{Zc(3900)^{reco}}$: [3.8869 ± 0.00003] GeV/c²

$\Gamma_{Zc(3900)^{reco}}$: [0.0308 ± 0.00007] GeV/c²

$M_{Zc(3900)^{evt.pdl}}$: [3.8872] GeV/c²

$\Gamma_{Zc(3900)^{evt.pdl}}$: [0.0282] GeV/c²

Zcs





$M_{Zcs(3985)^{reco}}$: [3.98279 ± 0.00002] GeV/c²

$\Gamma_{Zcs(3985)^{reco}}$: [0.0205 ± 0.00004] GeV/c²

$M_{Zcs(3985)^{evt.pdl}}$: [3.9825] GeV/c²

$\Gamma_{Zcs(3985)^{evt.pdl}}$: [0.0128] GeV/c²

VoigtFit:: quadratic Background + relativistic BW

Analysis

Background

- 30 million events were generated with Dual Parton Model (DPM)
- Same analysis strategy applied to background events
  - no event out of 30 million survived after the applied cuts.
  - The non-observation of any background events corresponds to a 90% confidence upper limit of 2.3 events. (means reco eff, $\epsilon_{bkg} = 2.3 \cdot 10^{-8}$)

Zc

- $\sigma_{sig} = [22.0 \pm 1.0]$ pb from Ref.[1]
- at a beam momentum of 8.54 GeV/c, the inelastic cross section is $\sigma_{bkg} = 45$ mb Ref.[2].



Zcs

- $\sigma_{sig} = [4.4 \pm 1.4]$ pb from Ref.[3] @ √s = 4.681 GeV
- at a beam momentum of 15 GeV/c, the inelastic cross section is $\sigma_{bkg} = 40$ mb Ref.[2].

- The branching ratio of J/psi is set to 100% during event generation. To correct this value for the following calculations, the branching ratio $Br_{sig} = Br_{J/\psi} = 5.961$ for the J/psi decay in the decay tree is taken into account.

Scaling factor is $F_{bkg} = \dfrac{N^{gen}_{sig} \cdot \sigma_{bkg}}{N^{gen}_{bkg} \cdot \sigma_{sig} \cdot Br_{sig}}$

Signal-to-Background ratio is defined as

$$\frac{S}{B} = \frac{\sigma_{sig} \cdot \epsilon_{sig} \cdot Br_{sig}}{\sigma_{bkg} \cdot \epsilon_{bkg}}$$

Signal significance is defined as

$$S_{sig} = \frac{N_{sig}}{\sqrt{N_{sig} \cdot + N_{bkg} \cdot F_{bkg}}}$$

[1] M. Ablikim et al. (BESIII Collaboration). Phys. Rev. Lett. 119, 072001 (2017)
[2] https://pdg.lbl.gov/2022/hadronic-xsections/
[3] M. Ablikim et al. (BESIII Collaboration), Phys. Rev. Lett. 126, 102001 (2021)

Summary

Particle type	ε^{reco} [%]	S / B x10^{-4}	S_{sig}
$Zc(3900)^+$ (from $\mu^+\mu^-$)	45.87	5.8	10.7391*
$Zc(3900)^+$ (from e^+e^-)	19.98	2.5	3.6877*
$Zc(3900)^-$ (from $\mu^+\mu^-$)	45.94	5.8	10.7035*
$Zc(3900)^-$ (from e^+e^-)	20.02	2.5	3.6905*

* assuming at least 1 background event

Particle type	ε^{reco} [%]	S / B	S_{sig}
$Zcs(3985)^+$ (from $\mu^+\mu^-$)	45.44	0.00013	5.4617*
$Zcs(3985)^+$ (from e^+e^-)	18.31	5.23x10^{-5}	1.6705*
$Zcs(3985)^-$ (from $\mu^+\mu^-$)	44.31	0.00013	5.4446*
$Zcs(3985)^-$ (from e^+e^-)	17.88	5.11x10^{-5}	1.6705*

* assuming at least 1 background event

Feature Works

- Increase DPM background statistics
- Generate events for Zc background study
  - $\bar{p}p \rightarrow \pi^+ + \pi^- + \mu^+ + \mu^-$,
  - $\bar{p}p \rightarrow \pi^+ + \pi^- + e^+ + e^-$

- Increase DPM background statistics
- Generate events for Zcs background study
  - $\bar{p}p \rightarrow K^+ + K^- + \mu^+ + \mu^-$,
  - $\bar{p}p \rightarrow K^+ + K^- + e^+ + e^-$

Analysis

Reconstruction of FS: transverse vs. longitudinal Momentum Distributions

μ^+ μ^- π^+ π^-



Analysis

Reconstruction of FS: total momentum vs. Θ angle Distributions

μ^+ μ^- π^+ π^-

















Analysis

Reconstruction of FS: Momentum Resolution

μ^+	μ^-	π^+	π^-
momentum diff vs. momentum for μ^+	momentum diff vs. momentum for μ^-	reco π^+ from Zc(3900)$^+$	momentum diff vs. momentum for π^- from $\bar{p}p$
reco momentum resolution for μ^+ from J/ψ	reco momentum resolution for μ^- from J/ψ	reco momentum resolution for π^+ from Zc(3900)$^+$	reco momentum resolution for π^+ from Zc(3900)$^+$
momentum resolution vs. momentum for μ^+	momentum resolution vs. momentum for μ^-	momentum resolution vs. momentum for π^+(Zc(3900)$^+$)	momentum resolution vs. momentum for π^- from $\bar{p}p$
momentum resolution vs. momentum for μ^+	momentum resolution vs. momentum for μ^-	momentum resolution vs. momentum for π^+(Zc(3900)$^+$)	momentum resolution vs. momentum for π^- from $\bar{p}p$

Backups

Electron channel : e^-













Backups

Electron channel : e^-













Backups

Electron channel : e^+













Backups

Electron channel : e^+













Backups

Electron channel : J/ψ













Backups

Electron channel : J/ψ