

# Subthreshold charm and strangeness production at GSI/FAIR energies

Marcus Bleicher  
(and Jan Steinheimer)

Institut für Theoretische Physik  
Goethe Universität - Frankfurt

# What is sub-threshold particle production?

...

And why is it interesting for us?

# Production of hadrons below threshold

- In elementary reactions, e.g. pp, it is not possible to produce a particle with mass  $m_{\text{new}}$ , if  $m_p + m_p + m_{\text{new}} > E_{\text{CM},pp}$  (energy conservation)
- However, in p+A and A+A reaction this is possible
- The question is, what mechanism allows for the production and are they realized


# Mechanisms

- Generally three different mechanisms are available:
  - 1) Fermi motion  
(more energy available than we thought)
  - 2) mass reduction/potentials  
(lowers the threshold for production)
  - 3) multi-step/multi-particle processes  
(collect energy to reach the threshold)

# This talk...

- Explores multi-strange particle production  
i.e.  $\phi$  and  $\Xi$  production  
→ solves a long standing puzzle at GSI energies
- Explores charm production  
i.e.  $J/\Psi$ ,  $L_c$  and D-mesons  
→ new road for a charm program at FAIR

# Motivation: $\phi$




## $\phi$ production

HADES and FOCI reported unexpected large  $\phi$  contribution to the  $K^-$  yield.

G. Agakishiev et al. [HADES Collaboration], Phys. Rev. C **80**, 025209 (2009)

# Motivation: $\Xi$



G. Agakishiev *et al.* [HADES Collaboration], Phys. Rev. C **80**, 025209 (2009)

## $\phi$ production

HADES and FOPI reported unexpected large  $\phi$  contribution to the  $K^-$  yield.

## $\Xi$ production

$\Xi^-$  yield, measured in Ar+KCl much larger than thermal model.

Confirmed in  $p+Nb \rightarrow$  No  $Y+Y$  exchange!!

Both particles are not well described in microscopic transport models and thermal fits are also not convincing.


Threshold for  $p+p \rightarrow p+p+\phi \approx 2.895$  GeV

Threshold for  $p+p \rightarrow N+\Xi+K+K \approx 3.24$  GeV

# Probabilities

## Sub-threshold production baseline

- Fermi momenta lift the collision energy above the threshold.
- Secondary interactions accumulate energy.




Why not introduce these decays for the less known resonances?



# New resonances


$N^*(1650)$	$\Delta(1232)$
$N^*(1710)$	$\Delta(1600)$
$N^*(1720)$	$\Delta(1620)$
$N^*(1875)$	$\Delta(1700)$
$N^*(1900)$	$\Delta(1900)$
$N^*(1990)$	$\Delta(1905)$
$N^*(2080)$	$\Delta(1910)$
$N^*(2190)$	$\Delta(1920)$
$N^*(2220)$	$\Delta(1930)$
$N^*(2250)$	$\Delta(1950)$
$N^*(2600)$	$\Delta(2440)$
$N^*(2700)$	$\Delta(2750)$
$N^*(3100)$	$\Delta(2950)$
$N^*(3500)$	$\Delta(3300)$
$N^*(3800)$	$\Delta(3500)$
$N^*(4200)$	$\Delta(4200)$



Important: New resonances replace the strings, no additional pp cross section is introduced

# Fixing the branching ratio

We use ANKE data on the  $\phi$  production cross section to fix the  $N^* \rightarrow N + \phi$  branching fraction.



Only 1 parameter

$$\Gamma_{N^* \rightarrow N\phi} / \Gamma_{\text{tot}} = 0.2\%$$

1 parameter fits all 3 points!


Y. Maeda *et al.* [ANKE Collaboration], Phys. Rev. C **77**, 015204 (2008) [arXiv:0710.1755 [nucl-ex]].

# Cross sections

Detailed balance  $\rightarrow$  absorption cross section

$$\frac{d\sigma_{b \rightarrow a}}{d\Omega} = \frac{\langle p_a^2 \rangle}{\langle p_b^2 \rangle} \frac{(2S_1 + 1)(2S_2 + 1)}{(2S_3 + 1)(2S_4 + 1)} \sum_{J=J_-}^{J_+} \frac{\langle j_1 m_1 j_2 m_2 || JM \rangle^2}{\langle j_3 m_3 j_4 m_4 || JM \rangle^2} \frac{d\sigma_{a \rightarrow b}}{d\Omega}$$

- $\phi + p$  cross section from detailed balance is very small.




# $\phi$ transparency ratios I

Model 1: The eikonal approximation of the Valencia group.

Model 2: Paryev developed the spectral function approach for  $\phi$  production in both the primary proton- nucleon and secondary pion- nucleon channels.

Model 3: BUU transport calculation of the Rossendorf group. Accounts for baryon- baryon and meson- baryon  $\phi$  production processes.

→ **20 mb absorption cross section for  $\phi+N$**




# Transparency ratios II

Detailed balance  $\rightarrow$  absorption cross section

$$\frac{d\sigma_{b \rightarrow a}}{d\Omega} = \frac{\langle p_a^2 \rangle}{\langle p_b^2 \rangle} \frac{(2S_1 + 1)(2S_2 + 1)}{(2S_3 + 1)(2S_4 + 1)} \sum_{J=J_-}^{J_+} \frac{\langle j_1 m_1 j_2 m_2 || JM \rangle^2}{\langle j_3 m_3 j_4 m_4 || JM \rangle^2} \frac{d\sigma_{a \rightarrow b}}{d\Omega}$$

- $\phi + p$  cross section from detailed balance is very small.
- Still the transparency ratio is well reproduced. Remember: this is what lead to the 20 mb cross section from ANKE.
- Cross section from transparency ratio is model dependent!





# New explanation

- $\phi + p$  cross section from detailed balance is very small.
- Still the transparency ratio is well reproduced. Remember: this is what lead to the 20 mb cross section from ANKE.
- Cross section from transparency ratio is model dependent!

- Not 'absorption' of the  $\phi$ , but of the mother resonance.
- Reactions of the type:  


$$N^* + N \rightarrow N'^* + N'^*$$

$$N^* + N \rightarrow N'^* + N'^*$$
 where the mass of  $N'^* < N^*$  so no  $\phi$  can be produced.



# Extrapolation to AA

When applied to nuclear collisions:




- Qualitative behavior nicely reproduced
- Predicted maximum at 1.25 A GeV
- High energies: too low due to string production
- HADES preliminary results for 1.23 A GeV, see HADES talks by R. Holzmann and T. Scheib.

Even centrality dependence is very well reproduced: Signal for multi step processes.

# Centrality

Even centrality dependence works well:





- Centrality dependence nicely reproduced.
- Good indicator for multi step production.

Data from: K. Piasecki et al., arXiv:1602.04378 [nucl-ex].



# Plain Kaon yields



Good description of the Kaon data

# Now for the $\Xi$

No elementary measurements near threshold.


We use  $p+Nb$  at  $E_{\text{lab}} = 3.5$  GeV data  $\rightarrow \Gamma_{N^* \rightarrow \Xi + K + K} / \Gamma_{\text{tot}} = 3.0\%$

HADES data	
$\langle \Xi^- \rangle$	$\Xi^- / \Lambda$
$(2.0 \pm 0.3 \pm 0.4) \times 10^{-4}$	$(1.2 \pm 0.3 \pm 0.4) \times 10^{-2}$
UrQMD	
$\langle \Xi^- \rangle$	$\Xi^- / \Lambda$
$(1.44 \pm 0.05) \times 10^{-4}$	$(0.71 \pm 0.03) \times 10^{-2}$

**Table:**  $\Xi^-$  production yield and  $\Xi^- / \Lambda$  ratio for minimum bias  $p + Nb$  collision at a beam energy of  $E_{\text{lab}} = 3.5$  GeV, compared with recent HADES results

G. Agakishiev *et al.*, Phys.Rev.Lett. 114 (2015) no.21, 212301.

# Comparison to data for $\Xi$



- $\Xi^-$  yield in Ar+KCl collisions is nicely reproduced
- Consistent with the p+Nb data.
- Indication for  $\Xi$  production from non-thermal 'tails' of particle production.
- All other strange particle ratios are also in line with experiment

# Can we also use this for charm?


• • •

Bold..., but possible...

J. Steinheimer, A. Botvina and M. Bleicher, arXiv:1605.03439 [nucl-th].

# Fixing the branching ratio

We use data from p+p at  $\sqrt{s} = 6.7$  GeV to fix the  $N^* \rightarrow N + J/\Psi$  branching fraction.



Only 1 parameter

$$\Gamma_{N^* \rightarrow N J \Psi} / \Gamma_{\text{tot}} = 2.5 \cdot 10^{-5}$$


## Assumptions

- We assume the associated production of  $N^* \rightarrow \Lambda_c + \bar{D}$  to be a factor 15 larger at that beam energy and to contribute about the half of the total charm production.
- We neglect  $D + \bar{D}$  pair production as it has a significantly higher threshold
- We neglect string production
- All the contributions should even increase the expected yield.

# $J/\Psi$ cross section

## Detailed balance $\rightarrow$ absorption cross section


- $J/\Psi + p$  cross section from detailed balance is very small.
- Not 'absorption' of the  $J/\Psi$ , but of the mother resonance.
- Reactions of the type:  
$$N^* + N \rightarrow N'^* + N'^*$$
$$N^* + N \rightarrow N'^* + N'^*$$
where the mass of  $N'^* < N^*$  so no  $J/\Psi$  can be produced.



Comparable to: D. Kharzeev and H. Satz, Phys. Lett. B **334**, 155 (1994).

# Predictions for SIS-100

When applied to central nuclear collisions (min. bias: divide by 5):





$$E_{\text{lab}} = 11 \text{ A GeV}$$

- $1.5 \cdot 10^{-6} J/\Psi$  per event
- $2 \cdot 10^{-5} \Lambda_c$  per event
- $\approx 3 - 4 \cdot 10^{-5} \bar{D}$  per event

# Comparison to others I

Parametrized cross section for  $J/\Psi$

$$\sigma_i^{NN}(s) = f_i a \left(1 - \frac{m_i}{\sqrt{s}}\right)^\alpha \left(\frac{\sqrt{s}}{m_i}\right)^\beta \theta(\sqrt{s} - \sqrt{s_{0i}})$$



HSD results taken from:


O. Linnyk, E. L. Bratkovskaya and W. Cassing, Int. J. Mod. Phys. E **17**, 1367 (2008)



# Comparison to others II

Cross section for  $p + p \rightarrow p + \bar{D}^0 + \Lambda_c$

Hadronic Lagrangian



Taken from:

W. Liu, C. M. Ko and S. H. Lee, Nucl. Phys. A **728**, 457 (2003)

# Summary

- A new mechanism for the production of  $\Xi$  and  $\phi$  is introduced and validated in elementary collisions
- This new branching ratio of high mass resonances is fitted to available data and extrapolated to AA
- It allows for the first time to describe the sub-threshold multi-strange particle production
- If this mechanism is also be applicable to charm production it may open a new road for charm studies at FAIR-SIS 100