



Contribution ID: 68

Type: Poster

## Predicting QCD phase diagram with chiral relativistic mean field model fitted to $\mu_B = 0$ lattice data

*Monday, 21 May 2018 18:30 (0 minutes)*

The SU(3) flavor parity-doublet quark-hadron model is used to investigate the phase diagram of QCD matter. The quark sector of the model is tuned to the  $\mu_B = 0$  lattice QCD data on trace anomaly. The structure of the baryon number susceptibilities in the temperature/chemical potential plane is studied in some detail. The model predicts three phase transitions - nuclear first-order liquid-gas phase transition, chiral symmetry restoration, and deconfinement transition.

At  $\mu_B = 0$ , a good agreement with the corresponding lattice data is obtained.

The deviations from the free hadron gas baseline in the crossover temperature region at  $\mu_B = 0$  are mainly attributed to the leftover of the liquid-gas transition in nuclear matter, the chiral phase transition determines the baryon fluctuations at much higher  $\mu_B$ , and at even higher baryon densities the behavior of fluctuations is controlled by deconfinement transition.

**Primary author:** Mr MOTORNENKO, Anton (Goethe Universität Frankfurt)

**Co-authors:** Prof. STOECKER, Horst (GSI Helmholtzzentrum fuer Schwerionenforschung); Dr STEINHEIMER, Jan (FIAS); Prof. SCHRAMM, Stefan (FIAS); Mr VOVCHENKO, Volodymyr (Frankfurt Institute for Advanced Studies(FIAS))

**Presenter:** Mr MOTORNENKO, Anton (Goethe Universität Frankfurt)

**Session Classification:** Poster Session