

# Production of $\Sigma(1385)$ at $p(4.5)p$

07.11.2018 | Waleed Esmail

Institut für Kernphysik (IKP), Forschungszentrum Jülich

# Outlines:


- Introduction.
- HADES spectrometer.
- Analysis strategy.
- Summary and outlook.

# Introduction:

- **Ultimate Goal:** Measurements of the *electromagnetic decays* of excited hyperons provide an important insight into their structure.
- *Investigation of the  $\Sigma(1385)^+$  resonance via its hadronic decay.*
  - its investigation can supply some important information concerning the  $\Sigma(1385)^0$ .
  - also the line shapes of the two resonances should at least be similar, what can serve as a cross check for the  $\Sigma(1385)^0$  reconstruction.
- In addition the *hyperon reconstruction* will benefit from the proposed *Forward Detector* that extends HADES acceptance to forward direction and essential for the reconstruction of the proton from hyperon decays.


# HADES Spectrometer and future forward detector:

HADES




Forward Detector

- Polar angle  $0.5^\circ - 6.5^\circ$





- Eight double layers 0, 90, 0, 90, and 0, 90, +45, -45




# Analysis Strategy:

- Production of  $\Sigma(1385)^+$  (Pluto++) 1 million events:



# Analysis Strategy:



- Utilize energy loss and velocity information to train an artificial neural network (ANN) to identify signal particles.



# Analysis Strategy:

- Lambda Reconstruction:

- Primary vertex: intersection between  $\pi^+$  and  $K^+$  tracks.
- Decay vertex: intersection between  $\pi^-$  and  $P$  tracks.



Distance of secondary vertex to primary vertex

$d_\Lambda > 15$  mm

Minimum distance between proton and  $\pi^-$  track



$d_{p\pi^-} < 20$  mm

Distance of  $\pi^-$  track to primary vertex

$d_{\pi^-} > d_p$

# Analysis Strategy:


- Lambda Reconstruction:




# Analysis Strategy:

- Lambda Reconstruction with Forward Detector:

P<sup>+</sup> HADES





P<sup>+</sup> HADES+FwDet





➤ combine all tracks from the forward detector and  $\pi^-$  from HADES.

Distance between tracks (hades, FwDet)





# Analysis Strategy:

- Lambda Reconstruction (HADES+FwDet):



# Analysis Strategy:

- Sigma Reconstruction:



# Summary and Outlook:

- Starting the analysis of  $\Sigma(1385)^+$  at beam energy 4.5 GeV.
- Study of the proposed forward detector
- Simulation of  $\Sigma(1385)^+$  to real/virtual photon decays.
- Optimizing the analysis strategy to real/virtual photon decays.
- Kinematic refit.
- Including background processes.

# References:

- J. Siebenson, Technische Universität München (2013)

Strange baryonic resonances below the KN threshold - Results from p+p reactions at the HADES experiment

- REPORT on  $\Xi$  and  $\Lambda$  production with the use of Pluto framework  
UPDATE

<https://indico.gsi.de/event/6041/>