Application of the 3-Fluid Hydrodynamic Event Generator THESEUS to CBM

Elena Volkova(Eberhard Karls Universität Tübingen(UT-PIT)) for the CBM collaboration

THESEUS

- Three-fluid hydrodynamics (3FH) was developed to simulate heavyion collisions at moderately relativistic energies.
- 3 ideals fluids: projectile, target and fireball nucleons
- Particlization is the process of changing from a fluid to a particle description. It allows to apply experimental cuts to particle distributions
- Final state hadronic rescattering processes are simulated using the UrQMD code.



Exploring QCD phase diagram



Where is critical point?

Is there phase transition between Hadron gas and QGP?

If yes, which type?

THESEUS EoS and rapidity distributions

3 different equations of state: purely hadronic, 1st order phase transition (2-phase) **EoS** with deconfinement and crossover EoS with deconfinement Will we see the difference after CBM acceptance cuts?



THESEUS for CBM, Elena Volkova

CBM setup

CBM is fixed target experiment with a polar angle acceptance between $2.5^{\circ} - 25^{\circ}$

Energy range: 2-10 AgeV at SIS100 for AuAu collisions



Effect of CBM acceptance cuts



Rapidity distribution for hadron gas before and after CBM



Rapidity distribution for crossover before and after CBM



Rapidity distribution for 1st order pht before and after CBM







Rapidity distribution for different EoS at 4 AGeV after CBM



Quantitative analysis of rapidity distributions

Rapidity distribution can be analyzed quantitatively by looking at the curvature at mid-rapidity for different energies.

$$C_{y} \equiv \left(y_{cm}^{3} \frac{d^{3}N}{dy^{3}}\right)_{y=y_{cm}} / \left(y_{cm} \frac{dN}{dy}\right)_{y=y_{cm}}$$





Conclusions

- Features of different EoS are (partly) presented in this talk;
- Rapidity distributions allows in principle to distinguish 1st order phase transition EoS and crossover with gas EoS;
- This features are preserved to some extend also with CBM acceptance cuts.
- The midrapidity shape can be measured by its «curvature»;
- Energy scan for the curvature C_y of the net proton rapidity distribution is observable for differentiation EoS. («Event simulation based on three-fluid hydrodynamics for collisions at energies available at the NICA and at the FAIR» P.Batyuk, D.Blaschke, Yu. B.Ivanov, Iu.Karpenko)



Backup

$$Y = \frac{1}{2} \ln(E + pz/E - pz)$$

$$C_{y} \equiv \left(y_{cm}^{3} \frac{d^{3}N}{dy^{3}}\right)_{y=y_{cm}} / \left(y_{cm} \frac{dN}{dy}\right)_{y=y_{cm}}$$



Rapidity evolution with energy

