

Contribution ID: 3

Type: Talk

Fission hindrance of high-K isomers in transfermium nuclei

Wednesday, 4 May 2022 11:45 (15 minutes)

To date, the fission branch of only a handful of identified high-K isomers has been measured or a lower limit inferred and, except in the cases of 250 No [1-3] and 254 Rf [4], all partial fission half-lives or their lower limits are reported to be shorter than the partial fission half-life of the corresponding ground state [5]. This is at odds with estimates of expected fission hindrances (defined as the ratio of the isomer and ground state partial fission half lives) due to the specialisation energy and reduced pairing fields associated with high-K states [6] and clearly calls for a revision of the available data. We report here on new measurements of the fission hindrances of high-K isomers in 250,252,254 No using the GABRIELA detector array [7,8] at the focal plane of the recoil separator SHELS [9].

[1] A. Belozerov et al., Eur. Phys. J. A 16 (2003) 447

- [2] D. Peterson et al., Phys. Rev. C 74 (2006) 014316
- [3] J. Kallunkathariyil et al. Phys. Rev. C 101 (2020) 011301(R)
- [4] H. David et al., Phys. Rev. Lett. 115 (2015) 132502}
- [5] F. Kondev, G. Dracoulis and T. Kibedi, At. Dat. And Nucl. Dat. Tables 103-104 (2015) 50
- [6] R.M. clark, EPJ Web of Conferences 131 (2016) 02002

[7] K. Hauschild et al., Nucl. Instr. Meth. A 560 (2006) 388-394

- [8] R. Chakma et al., Eur. Phys. J. A 56 (2020) 245
- [9] A. Popeko et al., Nucl. Instr. Meth. B 376 (2016) 140}

Primary authors: LOPEZ-MARTENS, Araceli (IJCLab); GALL, Benoit (IPHC CNRS)

Presenter: GALL, Benoit (IPHC CNRS)

Session Classification: Isomers in Heavy and Superheavy Nuclei