

Contribution ID: 34

Type: Talk

Auger and X-ray K-shell fluorescence measurements for Sc-44 isomeric decays

Wednesday, 4 May 2022 09:15 (15 minutes)

Despite being a well-studied nucleus close to stability, the K-shell X-ray and Auger fluorescence yield for scandium-44 are not very well defined. However, the low-lying nuclear structure of $^{44}_{21}$ Sc and its population in 44 Ti electron capture decay lends itself to extracting these quantities.

The first two excited states in ⁴⁴Sc are isomeric and lie at 68 keV and 146 keV with half-lives of 154.8(8) ns and 51.0(3) μ s respectively. The 146 keV level is populated by the electron capture decay of ⁴⁴Ti >99% of the time. By carefully measuring coincident *K* X-rays (at 4-5 keV) and γ decays over several months using an optimised ⁴⁴Ti source, the half-lives of the isomeric states can be fitted. This allows extraction of the fractional X-ray intensities for the initial electron capture decay as well as the subsequent internal electron conversion that competes with γ emission to de-excite the lowest two ⁴⁴Sc excited states. Thus, the relative X-ray-to-Auger *K*-shell fluorescence can be obtained for the three decay processes.

These fluorescence values are being compared to BrIccEmiss [1,2] predictions for which Monte-Carlo simulations and fits to the Evaluated Atomic Data Library (EADL) are combined. The results of this study will be reported.

- [1] B.Q. Lee, T. Kibedi, et al., Computational and Mathematical Methods in Medicine (2012).
- [2] B.Q. Lee, PhD thesis, Department of Nuclear Physics, The Australian National University (2017).

Primary authors: WHELDON, Carl (University of Birmingham); Prof. KOKALOVA, Tzany (University of Birmingham); Mr PIRRIE, Stuart (University of Birmingham); Dr TUNGATE, Garry (University of Birmingham); Prof. KIBEDI, Tibor (ANU)

Presenter: WHELDON, Carl (University of Birmingham)

Session Classification: Isomers in Nuclear Structure and Astrophysics ONLINE