



Contribution ID: 25

Type: Oral presentation

Can 3N force affect spectroscopic factors extracted from transfer and knockout reactions?

Wednesday, 26 June 2024 11:20 (20 minutes)

The direct reaction theory widely used to study single-particle spectroscopic strength in nucleon transfer experiments is based on a Hamiltonian with two-nucleon interactions only. We point out that in reactions where three-body effects are important, for example, such as (d, p) and $(p, 2p)$, an additional three-body force arises due to three-nucleon ($3N$) interaction between nucleons belonging to different fragments. We develop calculations of this $3N$ -induced force for one-nucleon removal reactions thus making an essential step towards bringing together nuclear structure theory, where $3N$ force is routinely used, and nuclear direct reaction theory, based on two-nucleon interactions only.

We study the effects of the $3N$ force on nucleon transfer in (d, p) and (d, n) reactions on ^{56}Ni , ^{48}Ca , ^{26m}Al and ^{24}O targets at deuteron incident energies between 4 and 40 MeV/nucleon. Deuteron breakup is treated exactly within a continuum discretized coupled-channel approach. We found that an additional three-body force can noticeably alter the angular distributions at forward angles, with consequences for spectroscopic factors' studies. We also present the study of transfer to $2p$ continuum in the $^{25}\text{F}(p, 2p)^{24}\text{O}$ reaction, involving the same overlap function as in the $^{24}\text{O}(d, n)^{25}\text{F}$ case, quantifying the differences in the spectroscopic factors due to additional $3N$ -induced force.

Collaboration

Primary author: TIMOFEYUK, Natalia (University of Surrey)

Co-authors: Dr MOSCHINI, Laura (University of Surrey); Dr GÓMEZ-RAMOS, Mario (University of Seville)

Presenter: TIMOFEYUK, Natalia (University of Surrey)

Session Classification: Wednesday morning 2