



Contribution ID: 96

Type: Oral presentation

Intermediate-energy Coulomb excitation of $N = 52$ isotones towards ^{100}Sn

Thursday, 27 June 2024 09:20 (20 minutes)

The Sn isotopes, containing the longest chain of isotopes between two doubly-magic nuclei, offer a fundamental testing ground for nuclear theories. Between the $N = 50$ and $N = 82$ shell closures, the 2_1^+ energies of all Sn isotopes are well established and show an almost constant value, as expected in the generalized seniority scheme. Within the same framework, the $B(E2)$ values should resemble an inverted parabola peaking at mid-shell. However, measurements in the most proton-rich Sn isotopes have shown a clear deviation from the expected behavior, with an enhancement of the transition probabilities towards ^{100}Sn . Although different calculations tend to agree on the neutron-rich side of the chain, significant differences are observed in the proton-rich side. This is particularly true for ^{102}Sn , where the difference between the predictions amounts to almost a factor of 3, making this isotope a good candidate for the investigation of the effects driving the nuclear structure in the vicinity of ^{100}Sn .

An experiment to measure for the first time the $B(E2)$ values in the $N = 52$ isotones towards ^{100}Sn , including ^{102}Sn , was performed at the Radioactive Isotope Beam Factory in Japan. A 345 MeV/nucleon beam of ^{124}Xe was fragmented on a 5-mm-thick Be target at the entrance of the BigRIPS separator. The $N = 52$ isotones of interest were identified on an event-by-event basis using the $B\rho - \Delta E - B\rho$ technique. A 0.5-mm Au target placed at the F8 focal plane was used to induce Coulomb excitation. Outgoing fragments were identified using the ZeroDegree spectrometer. The Au target was surrounded by the high-efficiency DALI2⁺ γ -detector array, composed of 226 NaI(Tl) detectors. Preliminary results on the Coulomb excitation cross sections and transition probabilities for ^{98}Pd , ^{100}Cd and ^{102}Sn will be presented, and their comparison with shell model and ab-initio calculations will be discussed.

Collaboration

Primary authors: CORTES, Martha Liliana (RIKEN Nishina Center); ALGORA, Alejandro (IFIC (CSIC-Univ. Valencia)); BROWNE, Frank (RIKEN Nishina Center); BRUGNARA, Daniele (Istituto Nazionale di Fisica Nucleare (INFN)(INFN-Legnaro)); CEDERKALL, Joakim (Lund University); CHEN, Sidong (The University of Hong Kong); CORAGGIO, Luigi (INFN); CORSI, Anna (CEA Saclay); DOORNENBAL, Pieter (RIKEN); GARGANO, Angela (INFN Milano); GERL, Jürgen (GSI Helmholtzzentrum für Schwerionenforschung GmbH(GSI)); GORSKA-OTT, Magdalena (GSI Helmholtzzentrum für Schwerionenforschung GmbH(GSI)); DE GREGORIO, Giovanni (INFN Napoli); HAEFNER, G. (Université Paris-Saclay); ITACO, Nunzio (INFN Napoli); KOIWAI, Takuma (The university of Tokyo); KOKUBUN, K. (The University of Tokyo); KOSEOGLOU, Pavlos (IKP TU-Darmstadt / GSI); KUBONO, S. (RIKEN Nishina Center); LEE, Jenny (The University of Hong Kong); LI, Pengjie (Institute of Modern Physics, Chinese Academy of Sciences); LIANG, P. (The University of Hong Kong); LIU, J. (The University of Hong Kong); LIU, zhong (Institute of Modern Physics, Chinese Academy of Sciences); LOKOTKO, Taras (The University of Hong Kong); MIYAGI, Takayuki (TU Darmstadt); PARK, Joochun (Lund University); Prof. SAKURAI, Hiroshi (RIKEN, Univ. of Tokyo); SARMIENTO PICO, Luis Gerardo (Lund University(LuU)); SCHWENK, Achim (Technische Universität Darmstadt); TANIUCHI, Ryo (University of York); WIMMER, Kathrin (GSI Helmholtzzentrum für Schwerionenforschung GmbH(GSI)); XIAN, W. (The University of Hong Kong); ZANON, Irene (INFN-LNL)

Presenter: CORTES, Martha Liliana (RIKEN Nishina Center)

Session Classification: Thursday morning 1