





# XYZs with Effective Field Theories

Nora Brambilla




Quark Confinement and  
the Hadron Spectrum since 1994



We are currently through a new revolution in particle physics

The present revolution: new particles discoveries beyond the Quark Model in the QQbar sector at/above the strong decay thres



$D^0 \bar{D}^0$  strong threshold



Plots for each experiment available at our




<https://qwg.ph.nat.tum.de/exoticshub/>

upcoming experiments



Samuel Ting: It was like to stumble in a village where people were living 70000 years

- Discovery of the first quark of heavy type  $Q$  ( $m_c > \Lambda_{\text{QCD}}$ )
- Confirmation of the quark model and QCD




Aubert et al. BNL 74

Samuel Ting: It was like to stumble in a village where people were living 70000 years

- Discovery of the first quark of heavy type  $Q$  ( $m_c > \Lambda_{\text{QCD}}$ )
- Confirmation of the quark model and QCD


—> **narrow width and asymptotic freedom:** annihilation at large scale controlled by a small coupling constant  $\alpha_s(2m_c) \ll 1$

—> **energy levels and confinement:** structure of levels controlled by a Cornell potential in a Schroedinger eq.




Aubert et al. BNL 74

### Cornell potential



Samuel Ting: It was like to stumble in a village where people were living 70000 years




- Discovery of the first quark of heavy type  $Q$  ( $m_c > \Lambda_{\text{QCD}}$ )
- Confirmation of the quark model and QCD

—> **narrow width and asymptotic freedom:** annihilation at large scale controlled by a small coupling constant  $\alpha_s(2m_c) \ll 1$

—> **energy levels and confinement:** structure of levels controlled by a Cornell potential in a Schroedinger eq.

### Cornell potential




Aubert et al. BNL 74


Heavy quarkonia are nonrelativistic systems: multiscale systems

Many scales: a challenge and an opportunity

# Quarkonium scales



Levels normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$



$$M(Y(1S)) = 9460 \text{ GeV}$$

$$M(J/\Psi) = 3097 \text{ GeV}$$


THE MASS SCALE IS PERTURBATIVE

$$m_Q \gg \Lambda_{\text{QCD}}$$


$$m_b \simeq 5 \text{ GeV}; m_c \simeq 1.5 \text{ GeV}$$

Normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$

# Quarkonium scales



Levels normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$



$$M(Y(1S)) = 9460 \text{ GeV}$$

$$M(J/\Psi) = 3097 \text{ GeV}$$

NR BOUND STATES HAVE AT LEAST 3 SCALES

$$m \gg mv \gg mv^2 \quad v \ll 1$$

$$mv \sim r^{-1}$$

THE SYSTEM IS NONRELATIVISTIC(NR)

$$\Delta E \sim mv^2, \Delta_{fs} E \sim mv^4$$

$$v_b^2 \sim 0.1, v_c^2 \sim 0.3$$

THE MASS SCALE IS PERTURBATIVE


$$m_Q \gg \Lambda_{\text{QCD}}$$

$$m_b \simeq 5 \text{ GeV}; m_c \simeq 1.5 \text{ GeV}$$


Normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$



# Quarkonium scales



Levels normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$



$$M(Y(1S)) = 9460 \text{ GeV}$$

$$M(J/\Psi) = 3097 \text{ GeV}$$

NR BOUND STATES HAVE AT LEAST 3 SCALES

$$m \gg mv \gg mv^2 \quad v \ll 1$$

$$mv \sim r^{-1}$$

and  $\Lambda_{\text{QCD}}$

THE SYSTEM IS NONRELATIVISTIC(NR)

$$\Delta E \sim mv^2, \Delta_{fs} E \sim mv^4$$

$$v_b^2 \sim 0.1, v_c^2 \sim 0.3$$


THE MASS SCALE IS PERTURBATIVE

$$m_Q \gg \Lambda_{\text{QCD}}$$

$$m_b \simeq 5 \text{ GeV}; m_c \simeq 1.5 \text{ GeV}$$

Normalized with respect to  $\chi_b(1P)$  and  $\chi_c(1P)$


- The different quarkonium radii provide different measures of the transition from a Coulombic to a confined bound state.



○ Godfrey Isgur PRD 32(85)189

quarkonia probe the perturbative (high energy) and non perturbative region (low energy) as well as the transition region in dependence of their radius  $r$

- The different quarkonium radii provide different measures of the transition from a Coulombic to a confined bound state.




quarkonia probe the perturbative (high energy) and non perturbative region (low energy) as well as the transition region in dependence of their radius  $r$

The present revolutions: nuclear matter phase diagram

Quarkonia are probe of QGP formation

Matsui Satz 1986  
idea of color screening  
in medium




Debye charge screening

$$V(r) \sim -\alpha_s \frac{e^{-m_D r}}{r}$$

$$m_D \sim gT$$

$$r \sim \frac{1}{m_D} \xrightarrow{\text{Bound state dissolve}}$$




Sequential  
Melting at  
different  
Temperature

The present revolutions: nuclear matter phase diagram

Quarkonia are probe of QGP formation

Matsui Satz 1986  
idea of color screening  
in medium




Debye charge screening

$$V(r) \sim -\alpha_s \frac{e^{-m_D r}}{r}$$

$$m_D \sim gT$$


$$r \sim \frac{1}{m_D} \xrightarrow{\text{Bound state dissolve}}$$



Sequential  
Melting at  
different  
Temperature

Experimental measurements:

$R_{AA}$  is the nuclear modification factor = yield of quarkonium in PbPb / yield in pp.




- CMS PLB 790 (2019) 270
- ALICE PLB 822 (2021) 136579
- ATLAS PRC 107 (2023) 054912

The present revolutions: nuclear matter phase diagram

Quarkonia are probe of QGP formation

Matsui Satz 1986  
idea of **color screening**  
in medium




Debye charge screening

$$V(r) \sim -\alpha_s \frac{e^{-m_D r}}{r}$$

$$m_D \sim gT$$


$$r \sim \frac{1}{m_D} \xrightarrow{\text{Bound state dissolve}}$$



Sequential  
Melting at  
different  
Temperature

Experimental measurements:

$R_{AA}$  is the nuclear modification factor = yield of quarkonium in PbPb / yield in pp.




- CMS PLB 790 (2019) 270
- ALICE PLB 822 (2021) 136579
- ATLAS PRC 107 (2023) 054912

Today a new paradigm emerged **beyond screening** relating the  $R_{AA}$  to the **nonequilibrium evolution of the heavy pair in medium**: medium induced dissociation and color singlet/octet recombination. **Quantum phenomenon to be addresses with quantum master equations**

# The present revolution: XYZ striking characteristics


Belle 2003: discovery of X(3872)



surprisingly narrow above strong decay thresholds, with surprising decay patterns

Sizeable hadronic prompt production cross sections (Tevatron, CMS, ATLAS, LHCb)


production in heavy ions where the strongly coupled QCD medium is formed



opens completely new perspectives!

# The present revolution: XYZ striking characteristics


Belle 2003: discovery of X(3872)



surprisingly narrow above strong decay thresholds, with surprising decay patterns

Sizeable hadronic prompt production cross sections (Tevatron, CMS, ATLAS, LHCb)

production in heavy ions where the strongly coupled QCD medium is formed



opens completely new perspectives!

XYZs not merely composite particles, have unique properties

Novel strongly correlated exotics systems

It is of fundamental interest to provide first principle predictions for spectra, transitions, production and medium evolution from QCD





The present revolution: XYZ a formidable theoretical challenge

Close/above threshold new degrees of freedom like **glue and light quarks** are nonperturbative part in the binding.

# The present revolution: XYZ a formidable theoretical challenge

Close/above threshold new degrees of freedom like **glue and light quarks** are nonperturbative part in the binding.

- Models assume some special degrees of freedom and a model interaction





- Lattice calculation of exotics masses are limited by the large number of open decay modes and they are not immediately suited for production and in medium studies

# The present revolution: XYZ a formidable theoretical challenge

Close/above threshold new degrees of freedom like **glue and light quarks** are nonperturbative part in the binding.

- Models assume some special degrees of freedom and a model interaction




- Lattice calculation of exotics masses are limited by the large number of open decay modes and they are not immediately suited for production and in medium studies

We need a flexible approach rooted in QCD that can address all properties of XYZ spectra, production and propagation in medium : Born Oppenheimer Effective Field Theory (BOEFT)

For quarkonium to become a probe of strong interactions, it should be treated in QCD :a very hard problem

Close to the bound state  $\alpha_s \sim v$




$$\sim \frac{1}{E - \left( \frac{p^2}{m} + V \right)}$$

- From  $\left( \frac{p^2}{m} + V \right) \phi = E \phi \rightarrow p \sim mv$  and  $E = \frac{p^2}{m} + V \sim mv^2$ .

For quarkonium to become a probe of strong interactions, it should be treated in QCD :a very hard problem

Close to the bound state  $\alpha_s \sim v$




$$\sim \frac{1}{E - \left(\frac{p^2}{m} + V\right)}$$

- From  $\left(\frac{p^2}{m} + V\right)\phi = E\phi \rightarrow p \sim mv$  and  $E = \frac{p^2}{m} + V \sim mv^2$ .

multiscale diagrams have a complicate power counting and contribute to all orders in the coupling


Difficult also for the lattice!



$$L^{-1} \ll \lambda \ll \Lambda \ll a^{-1}$$

For quarkonium to become a probe of strong interactions, it should be treated in QCD :a very hard problem

Close to the bound state  $\alpha_s \sim v$




From  $(\frac{p^2}{m} + V)\psi = E\psi \rightarrow p \sim mv$  and  $E = \frac{p^2}{m} + V \sim mv^2$ .

The problem is greatly simplified and predictivity is achieved by using Nonrelativistic Effective Field Theories

multiscale diagrams have a complicate power counting and contribute to all orders in the coupling


Difficult also for the lattice!

$$L^{-1} \ll \lambda \ll \Lambda \ll a^{-1}$$



# QQbar systems with NR EFT

Color degrees of freedom  
 $3 \times 3 = 1 + 8$   
 singlet and octet QQbar



Hard

Soft  
 (relative  
 momentum)


Ultrasoft  
 (binding energy)

$$\mathcal{L}_{\text{EFT}} = \sum_n c_n(E_\Lambda/\mu) \frac{O_n(\mu, \lambda)}{E_\Lambda}$$

$$\langle O_n \rangle \sim E_\lambda^n$$

# QQbar systems with NR EFT

Color degrees of freedom  
 $3 \times 3 = 1 + 8$   
 singlet and octet QQbar



$$\frac{E_\lambda}{E_\Lambda} = \frac{mv}{m}$$

Hard

Soft  
 (relative momentum)

Ultrasoft  
 (binding energy)


$$\mathcal{L}_{\text{EFT}} = \sum_n c_n(E_\Lambda/\mu) \frac{O_n(\mu, \lambda)}{E_\Lambda}$$

$$\langle O_n \rangle \sim E_\lambda^n$$



# QQbar systems with NR EFT

Color degrees of freedom  
 $3 \times 3 = 1 + 8$   
 singlet and octet QQbar



Hard

Soft  
 (relative momentum)

$$\frac{E_\lambda}{E_\Lambda} = \frac{mv}{m}$$

$$\frac{E_\lambda}{E_\Lambda} = \frac{mv^2}{mv}$$



Ultrasoft  
 (binding energy)

$$\mathcal{L}_{\text{EFT}} = \sum_n c_n(E_\Lambda/\mu) \frac{O_n(\mu, \lambda)}{E_\Lambda}$$

$$\langle O_n \rangle \sim E_\lambda^n$$



# QQbar systems with NR EFT: Non Relativistic QCD (NRQCD)

Caswell, Lepage 86, Lepage Thacker 88,  
Bodwin, Braaten, Lepage 95





# QQbar systems with NR EFT: Non Relativistic QCD (NRQCD)

Caswell, Lepage 86, Lepage Thacker 88,  
Bodwin, Braaten, Lepage 95



# QQbar systems with NR EFT: Non Relativistic QCD (NRQCD)



Caswell, Lepage 86, Lepage Thacker 88,  
Bodwin, Braaten, Lepage 95



$$\mathcal{L}_{\text{NRQCD}} = \sum_n c(\alpha_s(m/\mu)) \times \frac{O_n(\mu, \lambda)}{m^n}$$

# QQbar systems with NR EFT: Non Relativistic QCD (NRQCD)



Caswell, Lepage 86, Lepage Thacker 88, Bodwin, Braaten, Lepage 95





Only at the level of pNRQCD we obtain the potentials from QCD and the zero order problem is the Schroedinger equations

$$\mathcal{L}_{\text{NRQCD}} = \sum_n c(\alpha_s(m/\mu)) \times \frac{O_n(\mu, \lambda)}{m^n}$$



# Quarkonium with NR EFT: potential Non Relativistic QCD (pNRQCD)



# Quarkonium with NR EFT: potential Non Relativistic QCD (pNRQCD)





# Quarkonium with NR EFT: potential Non Relativistic QCD (pNRQCD)



$$\mathcal{L}_{\text{pNRQCD}} = \sum_k \sum_n \frac{1}{m^k} c_k(\alpha_s(m/\mu)) \times V(r\mu', r\mu) \times O_n(\mu', \lambda) r^n$$



# Quarkonium with NR EFT: potential Non Relativistic QCD (pNRQCD)



$$\mathcal{L}_{pNRQCD} = \sum_k \sum_n \frac{1}{m^k} c_k(\alpha_s(m/\mu)) \times V(r\mu', r\mu) \times O_n(\mu', \lambda) r^n$$

## Weakly coupled pNRQCD at the perturbative soft scale

○ Pineda Soto NP PS 64 (1998) 428

Brambilla Pineda Soto Vairo NPB 566 (2000) 275

- If  $mv \gg \Lambda_{\text{QCD}}$ , the matching is perturbative

Non-analytic behaviour in  $r \rightarrow$  matching coefficients  $V$

The gauge fields are multipole expanded:

$$A(R, r, t) = A(R, t) + \mathbf{r} \cdot \nabla A(R, t) + \dots$$

$\mathbf{R}$  = center of mass

$\mathbf{r}$  =  $Q\bar{Q}$  distance

$$\begin{aligned} \mathcal{L}^{\text{pNRQCD}} = & \int d^3r \text{Tr} \left\{ S^\dagger \left( i\partial_0 - \frac{\mathbf{p}^2}{m} - V_S + \dots \right) S + O^\dagger \left( iD_0 - \frac{\mathbf{p}^2}{m} - V_O + \dots \right) O \right. \\ & \left. + V_A (S^\dagger \mathbf{r} \cdot g\mathbf{E} O + O^\dagger \mathbf{r} \cdot g\mathbf{E} S) + \frac{V_B}{2} (O^\dagger \mathbf{r} \cdot g\mathbf{E} O + O^\dagger O \mathbf{r} \cdot g\mathbf{E}) \right\} + \dots \end{aligned}$$

$$-\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} + \sum_{i=1}^{n_f} \bar{q}_i i\not{D} q_i$$

LO in  $r$

NLO in  $r$

# Weakly coupled pNRQCD at the perturbative soft scale

o Pineda Soto NP PS 64 (1998) 428

Brambilla Pineda Soto Vairo NPB 566 (2000) 275

- If  $mv \gg \Lambda_{\text{QCD}}$ , the matching is perturbative

Non-analytic behaviour in  $r \rightarrow$  matching coefficients  $V$

The gauge fields are multipole expanded:

$$A(R, r, t) = A(R, t) + \mathbf{r} \cdot \nabla A(R, t) + \dots$$

$\mathbf{R}$  = center of mass

$\mathbf{r}$  =  $Q\bar{Q}$  distance

$$\mathcal{L}^{\text{pNRQCD}} = \int d^3r \text{Tr} \left\{ S^\dagger \left( i\partial_0 - \frac{\mathbf{p}^2}{m} - V_S + \dots \right) S + O^\dagger \left( iD_0 - \frac{\mathbf{p}^2}{m} - V_O + \dots \right) O + \right. \\ \left. + V_A (S^\dagger \mathbf{r} \cdot g\mathbf{E} O + O^\dagger \mathbf{r} \cdot g\mathbf{E} S) + \frac{V_B}{2} (O^\dagger \mathbf{r} \cdot g\mathbf{E} O + O^\dagger O \mathbf{r} \cdot g\mathbf{E}) \right\} + \dots$$

LO in  $r$

NLO in  $r$

$$-\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} + \sum_{i=1}^{n_f} \bar{q}_i i\not{D} q_i$$

The matching coefficients are the Coulomb potential

$$V_S(r) = -C_F \frac{\alpha_s}{r} + \dots,$$


$$V_O(r) = \frac{1}{2N} \frac{\alpha_s}{r} + \dots,$$

$$| V_A = 1 + \mathcal{O}(\alpha_s^2), V_B = 1 + \mathcal{O}(\alpha_s^2).$$


## Feynman rules

$$\text{---} = \theta(t) e^{-it(\mathbf{p}^2/m + V)}$$

$$\text{====} = \theta(t) e^{-it(\mathbf{p}^2/m + V_O)} \left( e^{-i \int dt A^{\text{adj}}} \right)$$




$$= O^\dagger \mathbf{r} \cdot g\mathbf{E} S$$



$$= O^\dagger \{ \mathbf{r} \cdot g\mathbf{E}, O \}$$

## Energies at order $m\alpha^5$ (NNNLO)

$$E_n = 2m + \langle n | \frac{p^2}{m} + V_s | n \rangle + \langle n | \text{---} \text{---} \text{---} | n \rangle$$


$$E_n = \langle n | H_s(\mu) | n \rangle - i \frac{g^2}{3N_c} \int_0^\infty dt \langle n | \mathbf{r} e^{it(E_n^{(0)} - H_o)} \mathbf{r} | n \rangle \langle \mathbf{E}(t) \mathbf{E}(0) \rangle(\mu)$$

$$E_n^{(0)} - H_o \gg \Lambda_{\text{QCD}} \Rightarrow \langle \mathbf{E}(t) \mathbf{E}(0) \rangle(\mu) \rightarrow \langle \mathbf{E}^2(0) \rangle$$

local condensates as predicted in a paper by Misha Voloshin in 1979


→ used to extract precise (NNNLO) determination of  $m_c$  and  $m_b$

$m\alpha_s^5 \ln \alpha_s$  Brambilla Pineda Soto Vairo 99, Kniehl Penin 99  
 $m\alpha_s^5$  Kniehl Penin Smirnov Steinhauser 02 NNLL Pineda 02

NNLL Peset Pineda et al 2018,2019, Kiyo Sumino 2014, Beneke, Kiyo Schuler 05,08

$\sim e^{i\Lambda_{\text{QCD}}t}$

$E_n^{(0)} - H_o \sim \Lambda_{\text{QCD}} \Rightarrow$  no expansion possible, non-local condensates, analogous to the Lamb shift in QED




Applications of weakly coupled pNRQCD include: precise alphas extraction from the static energy,  $t\bar{t}$  production, quarkonia spectra, decays, E1 and M1 transitions, QQq and QQQ energies, thermal masses and potentials

At the nonperturbative soft scale


Hitting the scale  $\Lambda_{\text{QCD}}$   $r \sim \Lambda_{\text{QCD}}^{-1}$

The degrees of freedoms now are

$(Q\bar{Q})_1$



$(Q\bar{Q})_1 + \text{Glueball}$



$(Q\bar{Q})_8 G$

hybrid /tetraquarks

with gluons/light quarks becoming part of the binding over the strong decay threshold


—> nonperturbative problem, use lattice

At the nonperturbative soft scale


Hitting the scale  $\Lambda_{QCD}$   $r \sim \Lambda_{QCD}^{-1}$

The degrees of freedoms now are

$(Q\bar{Q})_1$



$(Q\bar{Q})_1 + \text{Glueball}$



$(Q\bar{Q})_8 G$

hybrid /tetraquarks

with gluons/light quarks becoming part of the binding over the strong decay threshold

—> nonperturbative problem, use lattice


Use symmetry and scale separation:

$m > \Lambda_{QCD}$  NRQCD holds

$\Lambda_{QCD} > mv^2$  fast (gluons, light quarks) and slow (heavy quarks)

like in molecular physics (fast-electrons, slow nuclei)

The spectrum of static energies can be calculated in NRQCD



Symmetry of a system with a static  
 $Q$  in  $x_1$  and a  $\bar{Q}$  in  $x_2$


Irreducible representations of  $D_{\infty h}$

- $\mathbf{K}$ : angular momentum of light d.o.f.  
 $\lambda = \hat{\mathbf{r}} \cdot \mathbf{K} = 0, \pm 1, \pm 2, \pm 3, \dots$   
 $\Lambda = |\lambda| = 0, 1, 2, 3, \dots$  ( $\Sigma, \Pi, \Delta, \Phi, \dots$ )
- Eigenvalue of  $CP$ :  $\eta = +1$  ( $g$ ),  $-1$  ( $u$ )
- $\sigma$ : eigenvalue of reflection about a plane containing  $\hat{\mathbf{r}}$  (only for  $\Sigma$  states)

$$\Lambda_{\eta}^{\sigma}$$

The spectrum of static energies can be calculated in NRQCD

Symmetry of a system with a static Q in  $x_1$  and a  $\bar{Q}$  in  $x_2$



NRQCD static energies

Irreducible representations of  $D_{\infty h}$

- $\mathbf{K}$ : angular momentum of light d.o.f.  
 $\lambda = \hat{\mathbf{r}} \cdot \mathbf{K} = 0, \pm 1, \pm 2, \pm 3, \dots$   
 $\Lambda = |\lambda| = 0, 1, 2, 3, \dots$  ( $\Sigma, \Pi, \Delta, \Phi, \dots$ )
- Eigenvalue of  $CP$ :  $\eta = +1$  ( $g$ ),  $-1$  ( $u$ )
- $\sigma$ : eigenvalue of reflection about a plane containing  $\hat{\mathbf{r}}$  (only for  $\Sigma$  states)

$$\Lambda_{\eta}^{\sigma}$$

$$\mathcal{H}^{(0)} = \int d^3\mathbf{x} \frac{1}{2} (\boldsymbol{\Pi}^a \boldsymbol{\Pi}^a + \mathbf{B}^a \mathbf{B}^a) - \sum_{n_f} \bar{q} i \mathbf{D} \cdot \boldsymbol{\gamma} q$$


$$\mathcal{H}^{(0)} |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = E_n^{(0)}(\mathbf{x}_1, \mathbf{x}_2) |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$

$$|\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = \psi^\dagger(\mathbf{x}_1) \chi(\mathbf{x}_2) |n; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$

$$E_n^{(0)}(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle X_n, T/2 | X_n, -T/2 \rangle$$

$$|X_n\rangle = \chi(\mathbf{x}_2) \phi(\mathbf{x}_2, \mathbf{R}) T^a H^a(\mathbf{R}) \phi(\mathbf{R}, \mathbf{x}_1) \psi^\dagger(\mathbf{x}_1) |\text{vac}\rangle$$

Phi = Wilson lines and H = gluonic and light quarks







The spectrum of static energies can be calculated in NRQCD

Symmetry of a system with a static Q in  $x_1$  and a  $\bar{Q}$  in  $x_2$

Lattice Spectrum of NRQCD  
hybrid static energies  $E_n^{(0)}$



NRQCD static energies

- Irreducible representations of  $D_{\infty h}$
- $\mathbf{K}$ : angular momentum of light d.o.f.  
 $\lambda = \hat{r} \cdot \mathbf{K} = 0, \pm 1, \pm 2, \pm 3, \dots$   
 $\Lambda = |\lambda| = 0, 1, 2, 3, \dots$  ( $\Sigma, \Pi, \Delta, \Phi, \dots$ )
  - Eigenvalue of  $CP$ :  $\eta = +1$  ( $g$ ),  $-1$  ( $u$ )
  - $\sigma$ : eigenvalue of reflection about a plane containing  $\hat{r}$  (only for  $\Sigma$  states)
- $\Lambda_{\eta}^{\sigma}$

$$\mathcal{H}^{(0)} = \int d^3\mathbf{x} \frac{1}{2} (\mathbf{\Pi}^a \mathbf{\Pi}^a + \mathbf{B}^a \mathbf{B}^a) - \sum_{n_f} \bar{q} i \mathbf{D} \cdot \boldsymbol{\gamma} q$$


$$\mathcal{H}^{(0)} |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = E_n^{(0)}(\mathbf{x}_1, \mathbf{x}_2) |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$

$$|\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = \psi^\dagger(\mathbf{x}_1) \chi(\mathbf{x}_2) |n; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$

$$E_n^{(0)}(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle X_n, T/2 | X_n, -T/2 \rangle$$

$$|X_n\rangle = \chi(\mathbf{x}_2) \phi(\mathbf{x}_2, \mathbf{R}) T^a H^a(\mathbf{R}) \phi(\mathbf{R}, \mathbf{x}_1) \psi^\dagger(\mathbf{x}_1) |vac\rangle$$

Phi = Wilson lines and H = gluonic and light quarks



Juge Kuti Mornigstar 98-06


Schlosser, Wagner 2111.00741, Bali Pineda 2004

The spectrum of static energies can be calculated in NRQCD

Symmetry of a system with a static Q in  $x_1$  and a Qbar in  $x_2$


## Tetraquark static energies

Sadl, Prevlosek,  
211014568



$Z_b$  channel  $\vec{S}_h = \vec{S}_b + \vec{S}_{\bar{b}}$

Eigen-energies  $E_n(r)$  : channel  $S_h=1, CP=-1, \epsilon=-1$



Irreducible representations of  $D_{\infty h}$

- $\mathbf{K}$ : angular momentum of light d.o.f.  
 $\lambda = \hat{r} \cdot \mathbf{K} = 0, \pm 1, \pm 2, \pm 3, \dots$   
 $\Lambda = |\lambda| = 0, 1, 2, 3, \dots$  ( $\Sigma, \Pi, \Delta, \Phi, \dots$ )
- Eigenvalue of  $CP$ :  $\eta = +1$  ( $g$ ),  $-1$  ( $u$ )
- $\sigma$ : eigenvalue of reflection about a plane containing  $\hat{r}$  (only for  $\Sigma$  states)

$$\Lambda_{\eta}^{\sigma}$$

$$\mathcal{H}^{(0)} = \int d^3\mathbf{x} \frac{1}{2} (\mathbf{\Pi}^a \mathbf{\Pi}^a + \mathbf{B}^a \mathbf{B}^a) - \sum_{n_f} \bar{q} i \mathbf{D} \cdot \boldsymbol{\gamma} q$$

NRQCD static energies


$$\mathcal{H}^{(0)} |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = E_n^{(0)}(\mathbf{x}_1, \mathbf{x}_2) |\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$

$$|\underline{n}; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)} = \psi^\dagger(\mathbf{x}_1) \chi(\mathbf{x}_2) |n; \mathbf{x}_1, \mathbf{x}_2\rangle^{(0)}$$


$$E_n^{(0)}(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle X_n, T/2 | X_n, -T/2 \rangle$$

$$|X_n\rangle = \chi(\mathbf{x}_2) \phi(\mathbf{x}_2, \mathbf{R}) T^a H^a(\mathbf{R}) \phi(\mathbf{R}, \mathbf{x}_1) \psi^\dagger(\mathbf{x}_1) |\text{vac}\rangle$$

Phi = Wilson lines and H = gluonic and light quarks



Notice: in presence of light quark in the binding one adds isospin quantum numbers and measure tetraquark static energies



QED —


$$\Lambda_{QCD} \gg mv^2$$

Braaten PRL 111 (2013) 162003

Braaten Langmack Smith PRD 90 (2014) 014044


Born Oppenheimer  
Description

Higher excitations  
develop a gap of order  $\Lambda_{QCD}$



Introducing a finite mass m:

- The spectrum of the  $mv^2$  fluctuations around the lowest static energy is the **quarkonium spectrum**
- The spectrum of the  $mv^2$  fluctuations around the higher excitations is the **exotic spectrum (hybrids and tetraquarks)**



QED


$$\Lambda_{\text{QCD}} \gg mv^2$$

Braaten PRL 111 (2013) 162003

Braaten Langmack Smith PRD 90 (2014) 014044

Born Oppenheimer Description

Higher excitations develop a gap of order  $\Lambda_{\text{QCD}}$



Introducing a finite mass m:

- The spectrum of the  $mv^2$  fluctuations around the lowest static energy is the **quarkonium spectrum**
- The spectrum of the  $mv^2$  fluctuations around the higher excitations is the **exotic spectrum (hybrids and tetraquarks)**

Nonperturbative matching to the pNREFT


$$|0; \mathbf{x}_1 \mathbf{x}_2\rangle \rightarrow |(Q\bar{Q})_1\rangle \rightarrow \text{Quarkonium Singlet}$$

$$E_0(r) \rightarrow V_0(r) \quad (\text{Strongly coupled}) \text{pNRQCD}$$

$$|n > 0; \mathbf{x}_1 \mathbf{x}_2\rangle \rightarrow |(Q\bar{Q})_g^{(n)}\rangle \rightarrow \text{Higher Gluonic Excitations}$$

$$|Q\bar{Q}q\bar{q}\rangle \quad \text{Tetraquarks}$$

$$E_n^{(0)}(r) \rightarrow V_n^{(0)}(r) \quad \text{BOEFT}$$




QED

$$\Lambda_{\text{QCD}} \gg mv^2$$

Braaten PRL 111 (2013) 162003  
 Braaten Langmack Smith PRD 90 (2014) 014044

Born Oppenheimer Description

Higher excitations develop a gap of order  $\Lambda_{\text{QCD}}$



Introducing a finite mass m:

- The spectrum of the  $mv^2$  fluctuations around the lowest static energy is the **quarkonium spectrum**
- The spectrum of the  $mv^2$  fluctuations around the higher excitations is the **exotic spectrum (hybrids and tetraquarks)**

Nonperturbative matching to the pNREFT

systematically

$$\langle H | \mathcal{H} | H \rangle = \langle nljs | \frac{\mathbf{p}^2}{m} + \sum_n \frac{V_s^{(n)}}{m^n} | nljs \rangle$$

expand quantummechanically NRQCD states and energies in  $1/m$  around the zero order and identify the QCD potentials


$$| \underline{0}; \mathbf{x}_1 \mathbf{x}_2 \rangle \rightarrow | (Q\bar{Q})_1 \rangle \rightarrow \text{Quarkonium Singlet}$$

$$E_0(r) \rightarrow V_0(r) \quad (\text{Strongly coupled}) \text{pNRQCD}$$

$$| \underline{n} > 0; \mathbf{x}_1 \mathbf{x}_2 \rangle \rightarrow | (Q\bar{Q})_g^{(n)} \rangle \rightarrow \text{Higher Gluonic Excitations}$$

$$| Q\bar{Q}q\bar{q} \rangle \quad \text{Tetraquarks}$$

$$E_n^{(0)}(r) \rightarrow V_n^{(0)}(r) \quad \text{BOEFT}$$




pNRQCD and the potentials come from integrating  $mv^2$  out all scales up to

- gluonic excitations develop a gap  $\Lambda_{QCD}$  and are integrated out

Brambilla Pineda Soto Vairo 00

Bali et al. 98



pNRQCD and the potentials come from integrating out all scales up to  $mv^2$

- gluonic excitations develop a gap  $\Lambda_{QCD}$  and are integrated out
- ⇒ The singlet quarkonium field  $S$  of energy  $mv^2$  is the only the degree of freedom of pNRQCD (up to ultrasoft light quarks, e.g. pions).

Brambilla Pineda Soto Vairo 00

$$\mathcal{L} = \text{Tr} \left\{ S^\dagger \left( i\partial_0 - \frac{\mathbf{p}^2}{m} - V_s \right) S \right\} + \Delta\mathcal{L}(\text{US light quarks})$$

Bali et al. 98

- A pure potential description emerges from the EFT however this is not the constituent quark model, alphas and masses are the QCD fundamental parameters
- The potentials  $V = \text{Re}V + ImV$  from QCD in the matching: get spectra and decays
- We obtain the form of the nonperturbative potentials  $V$  in terms of generalized Wilson loops (stat that are low energy pure gluonic correlators: all the flavour dependence is pulled out)

Applications regard: Spectrum, decays, production at LHC, studies of confinement

The singlet potential has the general structure


the fact that spin dependent corrections appear at order  $1/m^2$  is called Heavy Quark Spin Symmetry

$$V = V_0 + \frac{1}{m} V_1 + \frac{1}{m^2} (V_{SD} + V_{VD})$$

static

spin dependent

velocity dependent

$$V^{(1)} = -\frac{1}{2} \int_0^\infty dt t \langle \text{Wilson Loop} \rangle$$


**gauge invariant wilson loops can be calculated also in QCD vacuum model and large N**

$$V_{SD}^{(2)} = -\frac{r^k}{r^2} c_F \epsilon^{kij} i \int_0^\infty dt t \langle \text{Wilson Loop} \rangle \mathbf{L}_1 \cdot \mathbf{S}_2 + (1 \leftrightarrow 2) |V_{LS}^{(2)}$$

$$-\frac{r^k}{r^2} \left( c_F \epsilon^{kij} i \int_0^\infty dt t \langle \text{Wilson Loop} \rangle - \frac{2c_F - 1}{2} \nabla^k V^{(0)} \right) \mathbf{L}_1 \cdot \mathbf{S}_1 + (1 \leftrightarrow 2) |V_{LS}^{(1)}$$

$$-c_F^2 \hat{r}_i \hat{r}_j i \int_0^\infty dt \left( \langle \text{Wilson Loop} \rangle - \frac{\delta_{ij}}{3} \langle \text{Wilson Loop} \rangle \right) \left( \mathbf{S}_1 \cdot \mathbf{S}_2 - 3(\mathbf{S}_1 \cdot \hat{r})(\mathbf{S}_2 \cdot \hat{r}) \right) |V_T$$

$$+ \left( \frac{2}{3} c_F^2 i \int_0^\infty dt \langle \text{Wilson Loop} \rangle - 4 \left( d_{sv} + \frac{4}{3} d_{vv} \right) \delta^{(3)}(\mathbf{r}) \right) \mathbf{S}_1 \cdot \mathbf{S}_2 |V_S$$



› Pineda Vairo PRD 63 (2001) 054007  
› Brambilla Pineda Soto Vairo PRD 63 (2001) 014023




The singlet potential has the general structure

the fact that spin dependent corrections appear at order  $1/m^2$  is called Heavy Quark Spin Symmetry

$$V = \underbrace{V_0}_{\text{static}} + \frac{1}{m} V_1 + \frac{1}{m^2} (V_{SD} + V_{VD})$$

↑ spin dependent
↑ velocity dependent

$$V^{(1)} = -\frac{1}{2} \int_0^\infty dt t \langle \text{Wilson Loop} \rangle$$


$$V_{SD}^{(2)} = -\frac{r^k}{r^2} c_F \epsilon^{kij} i \int_0^\infty dt t \langle \text{Wilson Loop} \rangle \mathbf{L}_1 \cdot \mathbf{S}_2 + (1 \leftrightarrow 2) |V_{LS}^{(2)}$$

$$-\frac{r^k}{r^2} \left( c_F \epsilon^{kij} i \int_0^\infty dt t \langle \text{Wilson Loop} \rangle - \frac{2c_F - 1}{2} \nabla^k V^{(0)} \right) \mathbf{L}_1 \cdot \mathbf{S}_1 + (1 \leftrightarrow 2) |V_{LS}^{(1)}$$

$$-c_F^2 \hat{r}_i \hat{r}_j i \int_0^\infty dt \left( \langle \text{Wilson Loop} \rangle - \frac{\delta_{ij}}{3} \langle \text{Wilson Loop} \rangle \right) \left( \mathbf{S}_1 \cdot \mathbf{S}_2 - 3(\mathbf{S}_1 \cdot \hat{r})(\mathbf{S}_2 \cdot \hat{r}) \right) |V_T$$





$$+ \left( \frac{2}{3} c_F^2 i \int_0^\infty dt \langle \text{Wilson Loop} \rangle - 4 \left( d_{sv} + \frac{4}{3} d_{vv} \right) \delta^{(3)}(\mathbf{r}) \right) \mathbf{S}_1 \cdot \mathbf{S}_2 |V_S$$

**gauge invariant wilson loops can be calculated also in QCD vacuum model and large N**

› Pineda Vairo PRD 63 (2001) 054007  
 › Brambilla Pineda Soto Vairo PRD 63 (2001) 014023

- the potentials contain the contribution of the scale  $m$  inherited from NRQCD matching coefficients—> they cancel any QM divergences, good UV behaviour
- the flavour dependent part is extracted in the NRQCD matching coefficients
- the nonperturbative part is universal:factorized and depends only on the glue —> only one lattice calculation to get the dynamics and the observables instead of an ab initio calculation of multiple Green functions

# Lattice evaluation of the spin dependent potentials



Koma Koma Wittig 05, Koma Koma 06

Terrific advance in the data precision with Lüscher multivel algorithm!

Such data can distinguish different models for the dynamics of low energy QCD e.g. effective string model

BOEFT for HYBRIDS

# Hybrids static energies at short distances We can calculate the perturbative behaviour of the potential for short distance

In the short-range hybrids become **gluelumps**, i.e., quark-antiquark octets,  $O^a$ , in the presence of a gluonic field,  $H^a: H(R, r, t) = H^a(R, t)O^a(R, r, t)$ .

the hybrid static energy can be written as a (multipole) expansion in  $r$ :


octet potential

$$E_g = \frac{\alpha_s}{6r} + \Lambda_g + a_g r^2 + \dots$$


non perturbative coefficient

$\Lambda_g$  is the **gluelump mass**:  $\Lambda_g = \lim_{T \rightarrow \infty} \frac{i}{T} \ln \langle H^a(T/2) \phi_{ab}^{\text{adj}}(T/2, -T/2) H^b(-T/2) \rangle$   
 calculated on the lattice

Foster Michael PRD 59 (1999) 094509  
 Bali Pineda PRD 69 (2004) 094001  
 Lewis Marsh PRD 89 (2014) 014502



$a_g$  can be expressed as field correlators (single line = singlet, double line = octet), e.g.,




In the limit  $r \rightarrow 0$  more symmetry:  $D_{\infty h} \rightarrow O(3) \times C$

- ▶ Several  $\Lambda_{\eta}^{\sigma}$  representations contained in one  $J^{PC}$  representation:
- ▶ Static energies in these multiplets have same  $r \rightarrow 0$  limit.

The gluelump multiplets  $\Sigma_u^-, \Pi_u; \Sigma_g^{+'}, \Pi_g; \Sigma_g^-, \Pi_g', \Delta_g; \Sigma_u^+, \Pi_u', \Delta_u$  are degenerate.

Gluonic excitation operators up to dim 3		
$\Lambda_{\eta}^{\sigma}$	$K^{PC}$	$H^a$
$\Sigma_u^-$	$1^{+-}$	$\mathbf{r} \cdot \mathbf{B}, \mathbf{r} \cdot (\mathbf{D} \times \mathbf{E})$
$\Pi_u$	$1^{+-}$	$\mathbf{r} \times \mathbf{B}, \mathbf{r} \times (\mathbf{D} \times \mathbf{E})$
$\Sigma_g^{+'}$	$1^{--}$	$\mathbf{r} \cdot \mathbf{E}, \mathbf{r} \cdot (\mathbf{D} \times \mathbf{B})$
$\Pi_g$	$1^{--}$	$\mathbf{r} \times \mathbf{E}, \mathbf{r} \times (\mathbf{D} \times \mathbf{B})$
$\Sigma_g^-$	$2^{--}$	$(\mathbf{r} \cdot \mathbf{D})(\mathbf{r} \cdot \mathbf{B})$
$\Pi_g'$	$2^{--}$	$\mathbf{r} \times ((\mathbf{r} \cdot \mathbf{D})\mathbf{B} + \mathbf{D}(\mathbf{r} \cdot \mathbf{B}))$
$\Delta_g$	$2^{--}$	$(\mathbf{r} \times \mathbf{D})^i (\mathbf{r} \times \mathbf{B})^j + (\mathbf{r} \times \mathbf{D})^j (\mathbf{r} \times \mathbf{B})^i$
$\Sigma_u^+$	$2^{+-}$	$(\mathbf{r} \cdot \mathbf{D})(\mathbf{r} \cdot \mathbf{E})$
$\Pi_u'$	$2^{+-}$	$\mathbf{r} \times ((\mathbf{r} \cdot \mathbf{D})\mathbf{E} + \mathbf{D}(\mathbf{r} \cdot \mathbf{E}))$
$\Delta_u$	$2^{+-}$	$(\mathbf{r} \times \mathbf{D})^i (\mathbf{r} \times \mathbf{E})^j + (\mathbf{r} \times \mathbf{D})^j (\mathbf{r} \times \mathbf{E})^i$

BOEFT for HYBRIDS



## BOEFT for $E_{\Pi_u}$ and $E_{\Sigma_u^-}$ hybrids

$$P_{\kappa\lambda}^{i\dagger} O^a(\mathbf{r}, \mathbf{R}, t) H_{\kappa}^{ia}(\mathbf{R}, t) = Z_{\kappa} \Psi_{\kappa\lambda}(\mathbf{r}, \mathbf{R}, t)$$

$$\mathcal{L}_{\text{BOEFT for } 1^{+-}} = \int d^3r \sum_{\lambda\lambda'} \text{Tr} \left\{ \Psi_{1^{+-}\lambda}^{\dagger} \left( i\partial_0 - V_{1^{+-}\lambda\lambda'}(r) + \hat{r}_{\lambda}^{i\dagger} \frac{\nabla_r^2}{m} \hat{r}_{\lambda'}^i \right) \Psi_{1^{+-}\lambda'} \right\}$$

- $\lambda = \pm 1, 0$ ;  $\hat{r}_0^i = \hat{r}^i$  and  $\hat{r}_{\pm 1}^i = \mp (\hat{\theta}^i \pm i\hat{\phi}^i) / \sqrt{2}$ .

- $V_{1^{+-}\lambda\lambda'} = V_{1^{+-}\lambda\lambda'}^{(0)} + \frac{V_{1^{+-}\lambda\lambda'}^{(1)}}{m} + \frac{V_{1^{+-}\lambda\lambda'}^{(2)}}{m^2} + \dots$

- For the static potential:  $V_{1^{+-}\lambda\lambda'}^{(0)} = \delta_{\lambda\lambda'} V_{1^{+-}\lambda}^{(0)}$ , with  $V_{1^{+-}0}^{(0)} = E_{\Sigma_u^-}$ ,  $V_{1^{+-}\pm 1}^{(0)} = E_{\Pi_u}$

fitted from the lattice hybrids static energies

## BOEFT for $E_{\Pi_u}$ and $E_{\Sigma_u^-}$ hybrids

$$P_{\kappa\lambda}^{i\dagger} O^a(\mathbf{r}, \mathbf{R}, t) H_{\kappa}^{ia}(\mathbf{R}, t) = Z_{\kappa} \Psi_{\kappa\lambda}(\mathbf{r}, \mathbf{R}, t)$$

$$\mathcal{L}_{\text{BOEFT for } 1^{+-}} = \int d^3r \sum_{\lambda\lambda'} \text{Tr} \left\{ \Psi_{1^{+-}\lambda}^{\dagger} \left( i\partial_0 - V_{1^{+-}\lambda\lambda'}(r) + \hat{r}_{\lambda}^{i\dagger} \frac{\nabla_r^2}{m} \hat{r}_{\lambda'}^i \right) \Psi_{1^{+-}\lambda'} \right\}$$

- $\lambda = \pm 1, 0$ ;  $\hat{r}_0^i = \hat{r}^i$  and  $\hat{r}_{\pm 1}^i = \mp (\hat{\theta}^i \pm i\hat{\phi}^i) / \sqrt{2}$ .

- $V_{1^{+-}\lambda\lambda'} = V_{1^{+-}\lambda\lambda'}^{(0)} + \frac{V_{1^{+-}\lambda\lambda'}^{(1)}}{m} + \frac{V_{1^{+-}\lambda\lambda'}^{(2)}}{m^2} + \dots$

- For the static potential:  $V_{1^{+-}\lambda\lambda'}^{(0)} = \delta_{\lambda\lambda'} V_{1^{+-}\lambda}^{(0)}$ , with  $V_{1^{+-}0}^{(0)} = E_{\Sigma_u^-}$ ,  $V_{1^{+-}\pm 1}^{(0)} = E_{\Pi_u}$

fitted from the lattice hybrids static energies

The LO e.o.m. for the fields  $\Psi_{1^{+-}\lambda}^{\dagger}$  are a set of coupled Schrödinger equations:

$$i\partial_0 \Psi_{1^{+-}\lambda} = \left[ \left( -\frac{\nabla_r^2}{m} + V_{1^{+-}\lambda}^{(0)} \right) \delta_{\lambda\lambda'} - \sum_{\lambda'} C_{1^{+-}\lambda\lambda'}^{\text{nad}} \right] \Psi_{\kappa\lambda'}$$

The eigenvalues  $\mathcal{E}_N$  give the masses  $M_N$  of the states as  $M_N = 2m + \mathcal{E}_N$ .

$$\hat{r}_{\lambda}^{i\dagger} \left( \frac{\nabla_r^2}{m} \right) \hat{r}_{\lambda'}^i = \delta_{\lambda\lambda'} \frac{\nabla_r^2}{m} + C_{1^{+-}\lambda\lambda'}^{\text{nad}}$$

with  $C_{1^{+-}\lambda\lambda'}^{\text{nad}} = \hat{r}_{\lambda}^{i\dagger} \left[ \frac{\nabla_r^2}{m}, \hat{r}_{\lambda'}^i \right]$  called the **nonadiabatic coupling**.

# BOEFT for $E_{\Pi_u}$ and $E_{\Sigma_u^-}$ hybrids

○ Berwein Brambilla Tarrus Vairo PRD 92 (2015) 114019  
 Oncala Soto PRD 96 (2017) 014004  
 Brambilla Krein Tarrus Vairo PRD 97 (2018) 016016

$$\mathcal{L}_{\text{BOEFT for } 1^{+-}} = \int d^3r \sum_{\lambda\lambda'} \text{Tr} \left\{ \Psi_{1^{+-}\lambda}^\dagger \left( i\partial_0 - V_{1^{+-}\lambda\lambda'}(r) + \hat{r}_\lambda^{i\dagger} \frac{\nabla_r^2}{m} \hat{r}_{\lambda'}^i \right) \Psi_{1^{+-}\lambda'} \right\}$$

- $\lambda = \pm 1, 0$ ;  $\hat{r}_0^i = \hat{r}^i$  and  $\hat{r}_{\pm 1}^i = \mp (\hat{\theta}^i \pm i\hat{\phi}^i) / \sqrt{2}$ .
- $V_{1^{+-}\lambda\lambda'} = V_{1^{+-}\lambda\lambda'}^{(0)} + \frac{V_{1^{+-}\lambda\lambda'}^{(1)}}{m} + \frac{V_{1^{+-}\lambda\lambda'}^{(2)}}{m^2} + \dots$
- For the static potential:  $V_{1^{+-}\lambda\lambda'}^{(0)} = \delta_{\lambda\lambda'} V_{1^{+-}\lambda}^{(0)}$ , with  $V_{1^{+-}0}^{(0)} = E_{\Sigma_u^-}$ ,  $V_{1^{+-}\pm 1}^{(0)} = E_{\Pi_u}$ .

fitted from the lattice hybrids static energies

$$\left[ -\frac{1}{mr^2} \partial_r r^2 \partial_r + \frac{1}{mr^2} \begin{pmatrix} l(l+1) + 2 & 2\sqrt{l(l+1)} \\ 2\sqrt{l(l+1)} & l(l+1) \end{pmatrix} + \begin{pmatrix} E_\Sigma^{(0)} & 0 \\ 0 & E_\Pi^{(0)} \end{pmatrix} \right] \begin{pmatrix} \psi_\Sigma^{(N)} \\ \psi_{-\Pi}^{(N)} \end{pmatrix} = \mathcal{E}_N \begin{pmatrix} \psi_\Sigma^{(N)} \\ \psi_{-\Pi}^{(N)} \end{pmatrix}$$



$$\left[ -\frac{1}{mr^2} \partial_r r^2 \partial_r + \frac{l(l+1)}{mr^2} + E_\Pi^{(0)} \right] \psi_{+\Pi}^{(N)} = \mathcal{E}_N \psi_{+\Pi}^{(N)}$$

Mixing remove the degeneration among opposite parity states:  
 -> Lambda doubling

- $l(l+1)$  is the eigenvalue of angular momentum  $\mathbf{L}^2 = (\mathbf{L}_{Q\bar{Q}} + \mathbf{L}_g)^2$  existing also in molecular physics
- the two solutions correspond to **opposite parity** states:  $(-1)^l$  and  $(-1)^{l+1}$
- corresponding eigenvalues under charge conjugation:  $(-1)^{l+s}$  and  $(-1)^{l+s+1}$



Hybrid multiplets as predicted by BOEFT (coloured rectangles) compared to the neutral isoscalar states observed in charmonium/bottomonium sector (crosses)



	$l$	$J^{PC} \{s=0, s=1\}$	$E_n^{(0)}$
$H_1$	1	$\{1^{--}, (0, 1, 2)^{-+}\}$	$\Sigma_u^-, \Pi_u$
$H_2$	1	$\{1^{++}, (0, 1, 2)^{+-}\}$	$\Pi_u$
$H_3$	0	$\{0^{++}, 1^{+-}\}$	$\Sigma_u^-$
$H_4$	2	$\{2^{++}, (1, 2, 3)^{+-}\}$	$\Sigma_u^-, \Pi_u$
$H_5$	2	$\{2^{--}, (1, 2, 3)^{-+}\}$	$\Pi_u$

**Note:** Band in the mass value for each multiplet is due to the error (150 MeV) on the gluelump mass measured on the lattice

Spin dependent interactions

# The **BOEFT** gives a prescription to calculate the **hybrids spin dependent potentials at order 1/m and 1/m<sup>2</sup>**

1/m

$$V_{1^{+-}\lambda\lambda'}^{(1)}_{SD}(\mathbf{r}) = V_{SK}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S}$$

$$+ V_{SKb}(r) \left[ \left( \mathbf{r} \cdot \hat{r}_\lambda^\dagger \right) \left( r^i \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} + \left( r^i \mathbf{K}^{ij} \hat{r}_\lambda^{j\dagger} \right) \cdot \mathbf{S} \left( \mathbf{r} \cdot \hat{r}_{\lambda'} \right) \right]$$

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2$$

$$S_{12} = 12(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - 4(\mathbf{S}_1 \cdot \mathbf{S}_2)$$

1/m<sup>2</sup>

$$V_{1^{+-}\lambda\lambda'}^{(2)}_{SD}(\mathbf{r}) = V_{LSa}^{(2)}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{L} \hat{r}_{\lambda'}^i \right) \cdot \mathbf{S} + V_{LSb}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \left( L^i S^j + S^i L^j \right) \hat{r}_{\lambda'}^j$$

$$+ V_{S^2}^{(2)}(r) \mathbf{S}^2 \delta_{\lambda\lambda'} + V_{S_{12}a}^{(2)}(r) S_{12} \delta_{\lambda\lambda'} + V_{S_{12}b}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \hat{r}_{\lambda'}^j \left( S_1^i S_2^j + S_2^i S_1^j \right)$$

$(K^{ij})^k = i\epsilon^{ikj}$  is the angular momentum of the spin one gluons

$\mathbf{L}$  is the orbital angular momentum of the heavy-quark-antiquark pair.

# The BOEFT gives a prescription to calculate the hybrids spin dependent potentials at order $1/m$ and $1/m^2$

$1/m$

$$V_{1^{+-}\lambda\lambda'}^{(1)}_{SD}(\mathbf{r}) = V_{SK}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} \\ + V_{SKb}(r) \left[ \left( \mathbf{r} \cdot \hat{r}_\lambda^\dagger \right) \left( r^i \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} + \left( r^i \mathbf{K}^{ij} \hat{r}_\lambda^{j\dagger} \right) \cdot \mathbf{S} \left( \mathbf{r} \cdot \hat{r}_{\lambda'} \right) \right] \quad \mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2$$

$$S_{12} = 12(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - 4(\mathbf{S}_1 \cdot \mathbf{S}_2)$$

$1/m^2$

$$V_{1^{+-}\lambda\lambda'}^{(2)}_{SD}(\mathbf{r}) = V_{LSa}^{(2)}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{L} \hat{r}_{\lambda'}^i \right) \cdot \mathbf{S} + V_{LSb}^{(2)}(r) \hat{r}_\lambda^{i\dagger} (L^i S^j + S^i L^j) \hat{r}_{\lambda'}^j \\ + V_{S^2}^{(2)}(r) \mathbf{S}^2 \delta_{\lambda\lambda'} + V_{S_{12}a}^{(2)}(r) S_{12} \delta_{\lambda\lambda'} + V_{S_{12}b}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \hat{r}_{\lambda'}^j \left( S_1^i S_2^j + S_2^i S_1^j \right)$$

$(K^{ij})^k = i\epsilon^{ikj}$  is the angular momentum of the spin one gluons  $\mathbf{L}$  is the orbital angular momentum of the heavy-quark-antiquark pair.

## Features:

- New spin structures with respect to the quarkonium case: all terms at order  $1/m$  and two terms at order  $1/m^2$

Differently from the quarkonium case, the hybrid potential gets a first contribution already at order  $\Lambda_{\text{QCD}}^2/m_h$ . The corresponding operator does not contribute at LO to matrix elements of quarkonium states as its projection on quark-antiquark color singlet states vanishes. Hence, **spin splittings are remarkably less suppressed in heavy quarkonium hybrids than in heavy quarkonia.**

# Hybrid spin dependent potentials at order 1/m and 1/m^2

1/m

$$V_{1^{+-}\lambda\lambda'}^{(1)}_{SD}(\mathbf{r}) = V_{SK}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} \\ + V_{SKb}(r) \left[ \left( \mathbf{r} \cdot \hat{r}_\lambda^\dagger \right) \left( r^i \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} + \left( r^i \mathbf{K}^{ij} \hat{r}_\lambda^{j\dagger} \right) \cdot \mathbf{S} \left( \mathbf{r} \cdot \hat{r}_{\lambda'} \right) \right]$$

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2$$

$$S_{12} = 12(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - 4(\mathbf{S}_1 \cdot \mathbf{S}_2)$$

1/m^2

$$V_{1^{+-}\lambda\lambda'}^{(2)}_{SD}(\mathbf{r}) = V_{LSa}^{(2)}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{L} \hat{r}_{\lambda'}^i \right) \cdot \mathbf{S} + V_{LSb}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \left( L^i S^j + S^i L^j \right) \hat{r}_{\lambda'}^j \\ + V_{S^2}^{(2)}(r) \mathbf{S}^2 \delta_{\lambda\lambda'} + V_{S_{12}a}^{(2)}(r) S_{12} \delta_{\lambda\lambda'} + V_{S_{12}b}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \hat{r}_{\lambda'}^j \left( S_1^i S_2^j + S_2^i S_1^j \right)$$

$(K^{ij})^k = i\epsilon^{ikj}$  is the angular momentum of the spin one gluons       $\mathbf{L}$  is the orbital angular momentum of the heavy-quark-antiquark pair.

Features:

- The nonperturbative part in  $V_i(r)$  depend on nonperturbative gluonic correlators non local in time not yet calculated on the lattice: six unknowns, the octet perturbative part can be calculated in perturbation theory
- The only flavor dependence is carried by the perturbative NRQCD matching coefficients

# Hybrid spin dependent potentials at order 1/m and 1/m^2

1/m

$$V_{1^{+-}\lambda\lambda' \text{ SD}}^{(1)}(\mathbf{r}) = V_{SK}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} \\ + V_{SKb}(r) \left[ \left( \mathbf{r} \cdot \hat{r}_\lambda^\dagger \right) \left( r^i \mathbf{K}^{ij} \hat{r}_{\lambda'}^j \right) \cdot \mathbf{S} + \left( r^i \mathbf{K}^{ij} \hat{r}_\lambda^{j\dagger} \right) \cdot \mathbf{S} \left( \mathbf{r} \cdot \hat{r}_{\lambda'} \right) \right]$$

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2$$

$$S_{12} = 12(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - 4(\mathbf{S}_1 \cdot \mathbf{S}_2)$$

1/m^2

$$V_{1^{+-}\lambda\lambda' \text{ SD}}^{(2)}(\mathbf{r}) = V_{LSa}^{(2)}(r) \left( \hat{r}_\lambda^{i\dagger} \mathbf{L} \hat{r}_{\lambda'}^i \right) \cdot \mathbf{S} + V_{LSb}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \left( L^i S^j + S^i L^j \right) \hat{r}_{\lambda'}^j \\ + V_{S^2}^{(2)}(r) \mathbf{S}^2 \delta_{\lambda\lambda'} + V_{S_{12}a}^{(2)}(r) S_{12} \delta_{\lambda\lambda'} + V_{S_{12}b}^{(2)}(r) \hat{r}_\lambda^{i\dagger} \hat{r}_{\lambda'}^j \left( S_1^i S_2^j + S_2^i S_1^j \right)$$

$(K^{ij})^k = i\epsilon^{ijk}$  is the angular momentum of the spin one gluons       $\mathbf{L}$  is the orbital angular momentum of the heavy-quark-antiquark pair.


Features:

- The nonperturbative part in  $V_i(r)$  depend on nonperturbative gluonic correlators non local in time not yet calculated on the lattice: six unknowns, the octet perturbative part can be calculated in perturbation theory
- The only flavor dependence is carried by the perturbative NRQCD matching coefficients

USE LATTICE CALCULATION OF THE CHARMONIUM SPIN MULTIPLETS TO EXTRACT the 6 UNKNOWNs and PREDICT THE BOTTOMONIUM SPIN MULTIPLETS, learn also about the **DYNAMICS**

# Charmonium Hybrids Multiplets $H_1$

lattice data from (violet) from  
G. K. C. Cheung, C. O'Hara, G. Moir, M. Peardon, S. M.  
Ryan, C. E. Thomas, and D. Tims (Hadron Spectrum),  
JHEP **12**, 089 (2016), arXiv:1610.01073 [hep-lat].  
with a pion of about 240 MeV




# Charmonium Hybrids Multiplets $H_1$

lattice data from (violet) from

G. K. C. Cheung, C. O'Hara, G. Moir, M. Peardon, S. M. Ryan, C. E. Thomas, and D. Tims (Hadron Spectrum), JHEP 12, 089 (2016), arXiv:1610.01073 [hep-lat].

with a pion of about 240 MeV

height of the boxes is an estimate of the uncertainty:  
 estimated by the parametric size of higher order corrections,  $m \alpha_s^5$  for the perturbative part, powers of  $\Lambda_{\text{qcd}}/m$  for the nonperturbative part, plus the statistical error on the fit





# Charmonium Hybrids Multiplets $H_1$

lattice data from (violet) from


G. K. C. Cheung, C. O'Hara, G. Moir, M. Peardon, S. M. Ryan, C. E. Thomas, and D. Tims (Hadron Spectrum), JHEP 12, 089 (2016), arXiv:1610.01073 [hep-lat].


with a pion of about 240 MeV

height of the boxes is an estimate of the uncertainty:  
 estimated by the parametric size of higher order corrections,  $m \alpha_s^5$  for the perturbative part, powers of  $\Lambda_{\text{qcd}}/m$  for the nonperturbative part, plus the statistical error on the fit


the perturbative part produces a pattern opposite to the lattice and to ordinary quarkonia  $\rightarrow$  discrepancy can be reconciled thanks to the nonperturbative parts, especially the one at order  $1/m$  which goes like  $\Lambda^2/m$  and is parametrically larger than the perturbative contribution at order  $m v^4$

which is interesting as some models are taking the spin interaction from perturbation theory with a constituent gluon







### Summary of $\bar{c}c$ $1^{--}$ masses




G. Ray, C. McNeile, 2110.14101

# Charmonium Hybrids Multiplets H<sub>1</sub> and H<sub>2</sub>



H<sub>1</sub> and H<sub>2</sub> corresponds to  $l=1$  and are negative and positive parity resp. The mass splitting between H<sub>1</sub> and H<sub>2</sub> is a result of lambda-doubling


H<sub>3</sub> and H<sub>4</sub> are also calculated



here you find predictions for all H multiplets

# Bottomonium hybrid spin splittings


thanks to the BOEFT factorization we can fix the nonperturbative unknowns from a charmonium hybrid calculation the nonperturbative low energy unknowns do not depend on the flavor: we can predict the bottomonium hybrids spin splittings



and also the other H multiplets

# Bottomonium hybrid spin splittings


thanks to the BOEFT factorization we can fix the nonperturbative unknowns from a charmonium hybrid calculation the nonperturbative low energy unknowns do not depend on the flavor: we can predict the bottomonium hybrids splin splittings



and also the other H multiplets

Comparison of our prediction to the existing lattice data on H<sub>1</sub>

# Bottomonium H<sub>1</sub> hybrid spin splittings



blue BOEFT predictions (more precise),  
violet actual lattice calculation

○ Ryan et al arXiv:2008.02656 [2+1 flavors,  $m_\pi = 400$  MeV]  
unpublished plot by J. Segovia and J. Tarrus

## Spin effects

->difficult to insert in models

->this spin structure has huge impact in phenomenology : larger spin multiplets separation than in quarkonium

->less spin symmetry in decays due to quarkonium-hybrids mixing via a spin operator at  $1/m$

**↓**  
**Oncala & Soto, Phys. Rev. D. 96, (2017)**

# Spin effects

- >difficult to insert in models
- >this spin structure has huge impact in phenomenology : larger spin multiplets separation than in quarkonium
- >less spin symmetry in decays due to quarkonium-hybrids mixing via a spin operator at  $1/m$

↓  
**Oncala & Soto, Phys. Rev. D. 96, (2017)**

- Hybrid states in the same energy range and same quantum #'s as quarkonium can mix.

Ex.  $H_1 [1^{--}] (4155) \leftrightarrow c\bar{c} [1^{--}] (3S)$

Effect on decay:  $H_m \leftrightarrow Q'_m \rightarrow (\eta_c, J/\psi, \dots) + (\gamma, \dots)$

- Mixing potential  $V_{\kappa\lambda}^{\text{mix}}$  : determined from matching NRQCD and BOEFT at  $O(1/m)$


Expression after matching:

$$V_{|\lambda|}^{\text{mix}} = -\frac{g_{CF}}{2m_Q} \langle 1 | B^j(\mathbf{r}/2, 0) | 0 \rangle^{(0)} P_{\lambda}^j$$

Above expression can be computed on lattice if we identify:

$$|0\rangle^{(0)} = |\Sigma_g^+\rangle$$

$$|1\rangle_{\lambda=0}^{(0)} = |\Sigma_u^-\rangle, |1\rangle_{|\lambda|=1}^{(0)} = |\Pi_u\rangle$$



$$\Gamma_{H \rightarrow S} = -2 \langle H | \text{Im} \Delta V | H \rangle.$$

**we calculated spin conserving and spin flipping decays  
they are same size**

**Decay to open threshold states not accounted**




$$\Gamma_{H \rightarrow S} = -2 \langle H | \text{Im} \Delta V | H \rangle.$$

we calculated spin conserving and spin flipping decays  
they are same size


Decay to open threshold states not accounted

- Comparison: bottom exotic states with corresponding bottomonium hybrid state:



BOEFT calculation of semi inclusive hybrids decays to quarkonium

Comparison: charm exotic states with corresponding charmonium hybrid state:



**$H_m \rightarrow Q_n + X$**

spin-conserving + spin-flipping decays  
 =  
lower bound on the total decay widths of hybrids  
 which is compared with inclusive rate of physical states in PDG.

# Hybrid: Summary

Brambilla, Lai, AM, Vairo arXiv:2212.09187

Hybrids ( $Q\bar{Q}g$ ): Color singlet state of color octet  $Q\bar{Q}$  + gluon. ( $Q = c, b$ )

✓ Isoscalar neutral mesons (Isospin=0)

✓ Candidates for hybrids based on **mass, quantum numbers**, and **decays** to quarkonium:

## Charm sector:

➤  $X(4160)$  : could be charm hybrid  $H_1[2^{-+}](4155)$ .

➤  $\psi(4710)$  : could be charm hybrid  $H_1[(1^{- -})](4812)$ .

➤  $X(4630)$  : could be charm hybrid  $H_1[(1/2^{- +})](4507)$ .

➤  $X(4630)$  : could be charm hybrid  $H_1[(1/2^{- +})](4507)$ .

➤  $\psi(4390)$  : could be charm hybrid  $H_1[1^{--}](4507)$ .

➤  $\chi_{c1}(4685)$  : could be charm hybrid  $H_2[(1^{+ +})](4667)$ .

## Bottom sector:

➤  $Y(10753)$  : could be bottom hybrid  $H_1[(1^{- -})](10786)$ .

### DISCLAIMER!!!

All the above interpretation can differ accounting for decays to meson-pair threshold states and hybrid-quarkonium mixing.

# Hybrid Decays Hybrid decays to meson-pair threshold states:

Conventional Wisdom: Hybrid decays to two S-wave mesons forbidden!  $H_m \not\rightarrow D^{(*)} \bar{D}^{(*)}$

Kou & Pene, Phys Lett B 631 (2005)    Page, Phys Lett B 407 (1997)    Farina, Tecocoatzi, Giachino, Santopinto & Swanson, Phys Rev D 102 (2020)

Born Oppenheimer quantum numbers for hybrids and ground state meson pair  
**does allow for decay to two s-wave mesons.**      **Bruschini 2306.17120**

	$l$	$J^{PC} \{s = 0, s = 1\}$	$E_n^{(0)}$
$H_1$	1	$\{1^{--}, (0, 1, 2)^{-+}\}$	$\Sigma_u^-, \Pi_u$
$H_2$	1	$\{1^{++}, (0, 1, 2)^{+-}\}$	$\Pi_u$
$H_3$	0	$\{0^{++}, 1^{+-}\}$	$\Sigma_u^-$
$H_4$	2	$\{2^{++}, (1, 2, 3)^{+-}\}$	$\Sigma_u^-, \Pi_u$
$H_5$	2	$\{2^{--}, (1, 2, 3)^{-+}\}$	$\Pi_u$

Most quarkonium hybrids can decay into pair of s-wave mesons !

forbidden for decay into pair of s-wave mesons

Recent lattice computation for  $c\bar{c}$  hybrid  $1^{-+}$  **decay** to

$D_1 \bar{D} : 258(133) \text{ MeV}$

$D^* \bar{D} : 88(18) \text{ MeV}$

$D^* \bar{D}^* : 150(118) \text{ MeV}$

Shi et al 2306.12884

BOEFT for tetraquarks and pentaquarks

**BOEFT may be used to describe any system made by two heavy quarks bound adiabatically with some light quarks degrees of freedom (tetraquarks QQlight quarks, QQbar light quarks, pentaquarks)**  
**In case of light quarks isospin quantum numbers should be added**

Steps go as before:

—identify the symmetries, identify the interpolating operators  $\mathcal{O}_n$

$$\mathcal{O}_n(t, \mathbf{r}, \mathbf{R}) = \chi(t, \mathbf{R} - \mathbf{r}/2) \phi(t, \mathbf{R} - \mathbf{r}/2, \mathbf{R}) H_n(t, \mathbf{R}) \phi(t, \mathbf{R}, \mathbf{R} + \mathbf{r}/2) \psi^\dagger(t, \mathbf{R} + \mathbf{r}/2)$$

—define the static energies

$$E_n^{(0)}(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle \mathcal{O}_n(T, \mathbf{r}, \mathbf{R}) | \mathcal{O}_n(0, \mathbf{r}, \mathbf{R}) \rangle .$$

-obtain the coupled Schroedinger equations in BOEFT

**BOEFT may be used to describe any system made by two heavy quarks bound adiabatically with some light quarks degrees of freedom (tetraquarks QQlight quarks, QQbar light quarks, pentaquarks)**  
**In case of light quarks isospin quantum numbers should be added**

Steps go as before:

—identify the symmetries, identify the interpolating operators  $\mathcal{O}_n$

$$\mathcal{O}_n(t, \mathbf{r}, \mathbf{R}) = \chi(t, \mathbf{R} - \mathbf{r}/2)\phi(t, \mathbf{R} - \mathbf{r}/2, \mathbf{R})H_n(t, \mathbf{R})\phi(t, \mathbf{R}, \mathbf{R} + \mathbf{r}/2)\psi^\dagger(t, \mathbf{R} + \mathbf{r}/2)$$

—define the static energies

$$E_n^{(0)}(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle \mathcal{O}_n(T, \mathbf{r}, \mathbf{R}) | \mathcal{O}_n(0, \mathbf{r}, \mathbf{R}) \rangle .$$

-obtain the coupled Schroedinger equations in BOEFT

**Notice:**

-the perturbative part of the potentials can be calculated

-the structure of the spin corrections will be similar to the hybrids case (with a 1/m spin correction)  
 calculation of decays will use the same technology

$$O_{\kappa,\lambda}^{Q\bar{Q}}(t, \mathbf{r}, \mathbf{R}) = \chi^\dagger(t, \mathbf{x}_2)\phi(t, \mathbf{x}_2, \mathbf{R})H_{\kappa,\lambda}^{Q\bar{Q}}(t, \mathbf{R})\phi(t, \mathbf{R}, \mathbf{x}_1)\psi(t, \mathbf{x}_1)$$

**I=0**

$$H_{0^{-+},0}(t, \mathbf{x}) = [\bar{q}(t, \mathbf{x})\gamma^5 T^a q(t, \mathbf{x})] T^a,$$

$$H_{1^{--},0}(t, \mathbf{x}) = [\bar{q}(t, \mathbf{x}) (\hat{\mathbf{r}} \cdot \boldsymbol{\gamma}) T^a q(t, \mathbf{x})] T^a,$$

$$H_{1^{--},\pm 1}(t, \mathbf{x}) = [\bar{q}(t, \mathbf{x}) (\hat{\mathbf{r}} \times \boldsymbol{\gamma}) T^a q(t, \mathbf{x})] T^a$$

$$E_{\kappa,\Lambda}(r) = V_o(r) + \Lambda_{H_\kappa} + \mathcal{O}(r^2)$$

Adjoint meson

$Q\bar{Q}$ color state	Light spin $\mathbf{K}^{PC}$	Static energies	$l$	$J^{PC}$ $\{S_Q = 0, S_Q = 1\}$	Multiplets
Octet	$0^{-+}$	$\{\Sigma_u^-\}$	0	$\{0^{++}, 1^{+-}\}$	$T_1^0$
			1	$\{1^{--}, (0, 1, 2)^{-+}\}$	$T_2^0$
			2	$\{2^{++}, (1, 2, 3)^{+-}\}$	$T_3^0$
	$1^{--}$	$\{\Sigma_g^{+'}, \Pi_g\}$	1	$\{1^{+-}, (0, 1, 2)^{++}\}$	$T_1^1$
			0	$\{0^{-+}, 1^{--}\}$	$T_2^1$
			1	$\{1^{-+}, (0, 1, 2)^{--}\}$	$T_3^1$
			2	$\{2^{-+}, (1, 2, 3)^{--}\}$	$T_4^1$

Coupled schroedinger eqs set up in BOEFT, need lattice input on the static tetra energies



## Tetraquark static energies

$I=1$  S. Prelovsek, H. Bahtiyar, J. Petrovich eprint: 1912.02656

$I=0$  Bicudo Cichy Peters Wagner PRD 93 (2016) 034501

Tetraquark static energies

$I=1$  S. Prelovsek, H. Bahtiyar, J. Petrovich eprint: 1912.02656

$I=0$  Bicudo Cichy Peters Wagner PRD 93 (2016) 034501

Mixing between quarkonium, hybrids; hybrids, tetraquarks

**Preliminary studies N. B. , Schlosser, Wagner, Vairo**

Tetraquark static energies

$I=1$  S. Prelovsek, H. Bahtiyar, J. Petrovich eprint: 1912.02656

$I=0$  Bicudo Cichy Peters Wagner PRD 93 (2016) 034501

Mixing between quarkonium, hybrids; hybrids, tetraquarks


**Preliminary studies N. B. , Schlosser, Wagner, Vairo**

Cross talk with the heavy light static energies

# Cross talk with the heavy light static energies

Adiabatic energy levels of the static energy of quarkonium  
and heavy-light, heavy-light strange-  
avoided level crossing

Bulava et al 1902.04006




# Cross talk with the heavy light static energies

Adiabatic energy levels of the static energy of quarkonium and heavy-light, heavy-light strange-  
avoided level crossing

In the diabatic picture gives the coupling between quarkonium and heavy light states allowing to solve the coupled schoedinger eqs and determine the amount of quarkonium and molecular states

$$\begin{pmatrix} \hat{V}(r) & g_1 & g_2 \\ g_1 & \hat{E}_1 & 0 \\ g_2 & 0 & \hat{E}_2 \end{pmatrix}.$$


Bulava et al 1902.04006



# Cross talk with the heavy light static energies

Adiabatic energy levels of the static energy of quarkonium and heavy-light, heavy-light strange-  
avoided level crossing

Bulava et al 1902.04006



In the diabatic picture gives the coupling between quarkonium and heavy light states allowing to solve the coupled schoedinger eqs and determine the amount of quarkonium and molecular states

$$\begin{pmatrix} \hat{V}(r) & g_1 & g_2 \\ g_1 & \hat{E}_1 & 0 \\ g_2 & 0 & \hat{E}_2 \end{pmatrix}.$$

The same cross talk with the heavy light static energies should be studied for hybrids and tetraquarks


# Cross talk with the heavy light static energies

Adiabatic energy levels of the static energy of quarkonium and heavy-light, heavy-light strange-  
avoided level crossing

In the diabatic picture gives the coupling between quarkonium and heavy light states allowing to solve the coupled schoedinger eqs and determine the amount of quarkonium and molecular states

$$\begin{pmatrix} \hat{V}(r) & g_1 & g_2 \\ g_1 & \hat{E}_1 & 0 \\ g_2 & 0 & \hat{E}_2 \end{pmatrix}.$$

Bulava et al 1902.04006




The same cross talk with the heavy light static energies should be studied for hybrids and tetraquarks

In this way special states with strong molecular components and characteristics like the X(3872) can be originated In BOEFT

The BOEFT contains all models: what dominates and where depends on the QCD dynamics

Static energies for  $I \neq 0$  (schematic):



The static energies are defined in BOEFT that gives the appropriate set of operators to be used and could describe the short distance limit.


Being nonperturbative objects  $E(r)$  should be calculated on the lattice (or in QCD vacuum models)

Figure from J. Tarrus



The BOEFT contains all models: what dominates and where depends on the QCD dynamics

Static energies for  $I \neq 0$  (schematic):



The static energies are defined in BOEFT that gives the appropriate set of operators to be used and could describe the short distance limit.

Being nonperturbative objects  $E(r)$  should be calculated on the lattice (or in QCD vacuum models)


Figure from J. Tarrus

# XYZ production and evolution in medium: may be studied with the EFT tools developed for quarkonium

## Bottomonium Nuclear Modification factor

can be obtained using pNRQCD at finite temperature, density matrix, and open quantum systems

$$R_{AA}(nS) = \frac{\langle n, \mathbf{q} | \rho_s(t_F; t_F) | n, \mathbf{q} \rangle}{\langle n, \mathbf{q} | \rho_s(0; 0) | n, \mathbf{q} \rangle}$$



Bands are from the dependence in kappa and gamma parameters

N.B., M. Escobedo, M. Strickland, A. Vairo P. VanderGriend et al




arXiv:2302.11826  
arXiv:2107.06222

arXiv:1711.04515

arXiv:2205.10289

## Quarkonium production can factorized and calculated in pNRQCD

N.B., Chung, Vairo, Wang 2210.17345



## Outlook

NREFTs and lattice allows us to describe the physics of quarkonium away from the strong decay threshold in quantum field theory: higher order perturbative calculation can be performed and quarkonium can be used for precision physics/ factorisation allows to systematically study confinement

BOEFT allows to describe hybrids: new unexpected features are found (Lambda doubling, Spin structure, decays, mixing) that have important impact on the phenomenology

BOEFT allows to describe hybrids and calculate multiplets, mixing and decays: on going work

The same picture can be extended to tetraquarks and pentaquarks, once some lattice input on relevant correlators will be available.

NOTICE that the needed lattice calculations are simpler than the direct calculations of the X Y Z properties on the lattice, the knowledge of few correlators together with the BOEFT will allow to obtain many phenomenological information

NREFTs and lattice and open quantum system allows us to describe the nonequilibrium evolution quarkonium in the quark gluon plasma and production processes: same theory could be then used for XYZ production and evolution in medium in heavy ion collisions

**This picture has the possibility to give a unified description to exotics and to leave the dynamics decide which configuration will dominate in a given range**

# Material for discussion/references

N. Brambilla, A. Pineda, J. Soto, A. Vairo

Rev. Mod. Phys. 77 (2005) 1423 • hep-ph/0410047

pNRQCD

N. Brambilla, A. Pineda, J. Soto, A. Vairo

Nucl. Phys. B566 (2000) 275 • hep-ph/9907240

A. Pineda, J. Soto, hep-ph/9707481

N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas,  
A. Vairo and C. Z. Yuan

*The XYZ states: experimental and theoretical status and perspectives*

Phys.Rept. 873 (2020) 1-154 • e-Print:

[1907.07583](#) [hep-ex]

BOEFT

QCD spin effects in the heavy hybrid potentials and spectra  
Nora Brambilla, Wai Kin Lai, J. Segovia, J. Tarrus

Phys.Rev.D 101 (2020) 5, 054040 • e-Print:

[1908.11699](#)

**Quarkonium Hybrids with Nonrelativistic Effective Field Theories**

Matthias Berwein, Nora Brambilla, Jaume Tarrús Castellà, Antonio Vairo

Phys.Rev. D92 (2015) no.11, 114019

e-Print: [arXiv:1510.04299](#)

**Born-Oppenheimer approximation in an effective field theory language**

Nora Brambilla, Gastão Krein, Jaume Tarrús Castellà, Antonio Vairo

Phys.Rev. D97 (2018) no.1, 016016

e-Print: [arXiv:1707.09647](#)

**Spin structure of heavy-quark hybrids**

N. Brambilla, Wai Kin Lai, J. Segovia, J. Tarrus  
A. Vairo

Phys.Rev.D 99 (2019) 1, 014017, e-Print:

[1805.07713](#) [hep-ph]

BOEFT

**Long range properties of 1S bottomonium states**

N. Brambilla, G. Krein, J. Tarrus, A. Vairo

Phys.Rev.D 93 (2016) 5, 054002 • e-Print: [1510.05895](#)

**Heavy hybrids decays to quarkonia**

N. Brambilla, A. Mohapatra, A. Vairo

Phys.Rev.D 107 (2023) 5, 054034, e-Print:

[2212.09187](#)[hep-ph]

**Oncala and Soto**

Phys.Rev.D 96 (2017) 1, 014004 •

Heavy hybrids: spectrum, decay and mixing

**Nonrelativistic effective field theory for heavy exotic hadrons**

J. Soto, J. Tarrus

Published in: *Phys.Rev.D* 102 (2020) 1, 014012 •

# Material for discussion/references

## Inclusive production of $J/\psi$ $\psi(2s)$ $Y$ states in pNRQCD

N. Brambilla, H. S. Chung, A. Vairo, X.P. Wang

*JHEP* 03 (2023) 242 2210.17345

## Quarkonium Production

## Gradient Flow

## Perturbative calculations

### QCD static force in gradient flow

N. Brambilla et al

*JHEP* 01 (2022) 184

e-Print: 2111.07811[hep-ph]

## Determination of the QCD coupling from the static energy and the free energy

A. Bazavov N. Brambilla, et al TUMQCD

*Phys.Rev.D* 100 (2019) 11, 114511 e-Print:

1907.11747[hep-ph]

## $\alpha_s$ Extraction

## Gradient Flow Lattice calculations

### QCD Force

J. Mayer-Steudte N. Brambilla et al TUMQCD

e-Print: 2212.12400[hep-ph]

## Static energy in (2+1+1) flavor lattice QCD

N. Brambilla, et al TUMQCD

*Phys.Rev.D* 107 (2023) 7, 074503, e-Print:

2206.03156[hep-ph]

## LATTICE

### EFTs and Lattice for XYZ

N. Brambilla

e-Print: 2111.10788[hep-ph]

### Lattice computation of the static force

N. Brambilla, V. Leino et al TUMQCD

*Phys.Rev.D* 105 (2022) 5, 054514 e-Print:

2106.01794 hep-ph]

# Material for discussion/references

N. Brambilla, M. A. Escobedo, J. Soto and A. Vairo, Phys. Rev. D **96** (2017) no.3, 034021 [arXiv:1612.07248 [hep-ph]].

N. Brambilla, M. A. Escobedo, J. Soto and A. Vairo, Phys. Rev. D **97** (2018) no.7, 074009 [arXiv:1711.04515 [hep-ph]].

N. Brambilla, M. A. Escobedo, A. Vairo and P. Vander Griend, Phys. Rev. D **100** (2019) no.5, 054025 [arXiv:1903.08063 [hep-ph]].

N. Brambilla, M. A. Escobedo, M. Strickland, A. Vairo, P. Vander Griend and J. H. Weber, [arXiv:2012.01240 [hep-ph]]. and arXiv: 2107 .06222

M. Escobedo *Phys.Rev.D* 103 (2021) 3, 034010 • e-Print: 2010.10424 [  
N. Brambilla, M. Escobedo, A. Islam, M. Strickland . A. Tiwari, A. Vairo arXiv 2205.10289

**Regeneration of bottomonia in an open quantum systems approach**

N. Brambilla, M. Escobedo, M Strickland et al 2302.11826

Non-equilibrium  
evolution in QGP

N. Brambilla, J. Ghiglieri, A. Vairo, Peter Petreczky  
*Phys.Rev.D* 78 (2008) 014017 • e-Print: 0804.0993 [hep-ph]

N. Brambilla, M. Escobedo, J. Ghiglieri, A. Vairo  
*JHEP* 05 (2013) 130 • e-Print: 1303.6097 [hep-ph]

N. Brambilla, M. Escobedo, J. Ghiglieri, A. Vairo  
*JHEP* 12 (2011) 116 • e-Print: 1109.5826 [hep-ph]

N. Brambilla, M. Escobedo, J. Ghiglieri, J. Soto, A. Vairo  
*JHEP* 09 (2010) 038 • e-Print: 1007.4156 [hep-ph]

Potential and  
energies  
in medium

N. Brambilla, V. Leino, Peter Petreczky,  
A. Vairo e-Print 2007.10078 [hep-ph],  
N. Brambilla, V. Leino, J. Meyer  
Steudte, P. Petreczky 2206.02861

Lattice calculation of the  
heavy quark transport coefficient

Tetraquarks	Order	$M_{Q\bar{Q}Q\bar{Q}}$ [GeV]	$B_{Q\bar{Q}Q\bar{Q}}$ [MeV]
$T_{cc\bar{c}\bar{c}}$	LO	6.1276(3)	16.6(4)
	NLO	6.078(2)	67.9(1)
	NNLO'	6.018(3)	144(2)
$T_{cc\bar{c}\bar{b}}/T_{bc\bar{c}\bar{c}}$	LO	9.294(3)	23.0(4)
	NLO	9.312(4)	72(2)
	NNLO'	9.259(5)	139(2)
$T_{bb\bar{c}\bar{c}}/T_{cc\bar{b}\bar{b}}$	LO	12.503(1)	23.7(4)
	NLO	12.457(4)	79(2)
	NNLO'	12.386(3)	157(3)
$T_{bc\bar{c}\bar{c}}$	LO	12.471(5)	19.5(8)
	NLO	12.417(5)	69(2)
	NNLO'	12.354(6)	139(2)
$T_{bb\bar{b}\bar{c}}/T_{bc\bar{b}\bar{b}}$	LO	15.652(6)	27.9(7)
	NLO	15.50(2)	87(2)
	NNLO'	15.37(7)	169(4)
$T_{bb\bar{b}\bar{b}}$	LO	18.8693(5)	31.2(6)
	NLO	18.8207(6)	83.6(1)
	NNLO'	18.7598(6)	151(1)

Variational and Green function Monte Carlo method based on Weakly coupled pNRQCD potential calculated at LO NLO and NNLO' (prime means only two body forces are considered)




Decays may be calculated in the same framework

TABLE II. Predictions for tetraquark masses and binding energies for all combinations of tetraquarks involving only  $b$  and  $c$  quarks at each order of pNRQCD indicated. Pairs of tetraquarks in the same row have identical binding energies in our calculations due to charge conjugation.


# Low energy physics factorized in Wilson loops: can be used to probe the confinement mechanism

any QCD vacuum model is an assumption on the behaviour of the Wilson loop

Bali et al



Boryakov et al. 04






## Spectrum: general consideration

Multiplet	$T$	$J^{PC}(S=0)$	$J^{PC}(S=1)$	$E_{\Gamma}$
$H_1$	1	$1^{--}$	$(0, 1, 2)^{-+}$	$E_{\Sigma_u^-}, E_{\Pi_u}$
$H_2$	1	$1^{++}$	$(0, 1, 2)^{+-}$	$E_{\Pi_u}$
$H_3$	0	$0^{++}$	$1^{+-}$	$E_{\Sigma_u^-}$
$H_4$	2	$2^{++}$	$(1, 2, 3)^{+-}$	$E_{\Sigma_u^-}, E_{\Pi_u}$

**Spin degenerated**

## Spectrum: with mixing and $\Lambda$ -doubling



- The Schrödinger equation mixes states with the same parity. A consequence is  $\Lambda$ -doubling, i.e., the lifting of degeneracy between states with opposite parity. This happens also in molecular physics, however, there  $\Lambda$ -doubling is a subleading effect, while it is a LO effect in the quarkonium hybrid spectrum.
- The eigenstates are organized in the multiplets  $H_1, H_2, \dots$ . Neglecting off-diagonal terms, the multiplets  $H_1$  and  $H_2$  would be degenerate.
- We compute the spectrum using quark masses in the renormalon subtraction (RS) scheme:  $m_{c\text{RS}} = 1.477(40)$  GeV and  $m_{b\text{RS}} = 4.863(55)$  GeV.

The glueball masses, which enter in the normalization of the hybrid potentials, have been computed in the same scheme and assigned an uncertainty of  $\pm 0.15$  GeV, which is the largest source of uncertainty in the hybrid masses.

charmonium hybrids

○ Berwein Brambilla Tarrus Vairo PRD 92 (2015) 114019  
data without mixing (dashed) from Braaten et al PRD 90 (2014)

Without Lambda-doubling masses of opposite parity states are degenerate