

CP violation tests in hyperon decays at BESIII

Varvara Batozskaya
on behalf of the BESIII collaboration

Institute of High Energy Physics, Beijing, China
National Centre for Nuclear Research, Warsaw, Poland

FAIR next generation scientists
8th Edition Workshop
Donji Seget, Croatia
23-27 September 2024




Introduction to hadron physics

- Two main questions to particle physics:
 - What are the most elementary building blocks of matter?
 - What are the forces between them?
 - SM is well-established theoretical description of fundamental particles and their interactions
 - Building blocks:
 - Quarks and leptons
 - Forces mediated by exchange of gauge bosons associated with weak, electromagnetic and strong interactions
- | three generations of matter (fermions) | | | interactions / force carriers (bosons) |
|---|---|---|--|
| I | II | III | |
| $\approx 2.16 \text{ GeV}/c^2$
$\frac{1}{2}$
u
up | $\approx 1.273 \text{ GeV}/c^2$
$\frac{1}{2}$
c
charm | $\approx 172.57 \text{ GeV}/c^2$
$\frac{1}{2}$
t
top | 0
0
g
gluon |
| $\approx 4.7 \text{ MeV}/c^2$
$\frac{1}{2}$
d
down | $\approx 93.5 \text{ MeV}/c^2$
$\frac{1}{2}$
s
strange | $\approx 4.183 \text{ GeV}/c^2$
$\frac{1}{2}$
b
bottom | 0
0
1
0
γ
photon |
| QUARKS | | | |
| $\approx 0.511 \text{ MeV}/c^2$
-1
e
electron | $\approx 105.66 \text{ MeV}/c^2$
-1
μ
muon | $\approx 1.776933 \text{ GeV}/c^2$
-1
τ
tau | 0
0
1
0
Z
Z boson |
| $\approx 0.8 \text{ eV}/c^2$
0
ν_e
electron neutrino | $\approx 0.17 \text{ MeV}/c^2$
0
ν_μ
muon neutrino | $\approx 18.2 \text{ GeV}/c^2$
0
ν_τ
tau neutrino | $\approx 91.1881 \text{ GeV}/c^2$
± 1
W
W boson |
| LEPTONS | | | GAUGE BOSONS
VECTOR BOSONS |
- SM is successfully tested by numerous experiments discovered a number of:
 - Non-standard hadronic states with properties indicating their complex substructure [PhysRept873(2020)1]
 - Conventional hadrons: $q\bar{q}$ mesons and qqq baryons [PhysLett8(1964)214]


Hadrons: conventional & exotic

[NatureRevPhys1(2019)480]



Hadrons: conventional & exotic





[NatureRevPhys1(2019)480]







None of exotic hadrons is settled!

Main contributors to exotic hadrons

- e^+e^- collider



- Hadron collider




* Not all experiments are covered

Brief discussion of exotic particles: XYZ states



Brief discussion of exotic particles: XYZ states

- **X**: neutral non-vector states
 $B^\pm \rightarrow K^\pm \mathbf{X} \rightarrow K(\pi^+\pi^- J/\psi)$
- **X(3872)**:
 - First discovered by Belle, $N_{\text{evt}} \sim 36$
 - Later confirmed by other experiments



Brief discussion of exotic particles: XYZ states




- **X:** neutral non-vector states
 $B^\pm \rightarrow K^\pm \text{X} \rightarrow K(\pi^+\pi^- J/\psi)$
- **X(3872):**
 - First discovered by Belle, $N_{\text{evt}} \sim 36$
 - Later confirmed by other experiments
- **Y:** neutral vector states
 $e^+e^- \rightarrow \gamma_{\text{ISR}} \text{Y} \rightarrow \gamma_{\text{ISR}}(\pi^+\pi^- J/\psi)$
- **Y(4260):**
 - First discovered by Babar, $N_{\text{evt}} \sim 125$
 - Later confirmed by other experiments




Brief discussion of exotic particles: XYZ states




- **X:** neutral non-vector states
 $B^\pm \rightarrow K^\pm \text{X} \rightarrow K(\pi^+\pi^- J/\psi)$
- **X(3872):**
 - First discovered by Belle, $N_{\text{evt}} \sim 36$
 - Later confirmed by other experiments
- **Y:** neutral vector states
 $e^+e^- \rightarrow \gamma_{\text{ISR}} \text{Y} \rightarrow \gamma_{\text{ISR}}(\pi^+\pi^- J/\psi)$
- **Y(4260):**
 - First discovered by Babar, $N_{\text{evt}} \sim 125$
 - Later confirmed by other experiments
- **Z:** charged, clearly multi-quark states
 $B \rightarrow K \text{Z}^\pm \rightarrow K(\pi^\pm \psi')$
- **Z $^\pm(4430)$:**
 - First discovered by Belle, $N_{\text{evt}} \sim 121$
 - Not confirmed by Babar
 - Confirmed by LHCb



BESIII hadron discoveries




BESIII hadron discoveries



- States produced directly in e^+e^- collision with no clear interpretation
- States with exotic flavour combinations decaying into heavy mesons
- New light states decaying into mesons
- State consistent with conventional $c\bar{c}$ meson
- New light baryon states
- State with exotic J^{PC}

Experimental facility:
BEPCII and BESIII

- Beijing Electron-Positron Collider (BEPCII)
 - e^+e^- collider with $2.0 \text{ GeV} < E_{\text{CMS}} < 4.95 \text{ GeV}$
 - $\mathcal{L}_{\text{peak}} = 10^{33} \text{ cm}^{-2} \text{s}^{-1}$
 - Data taking since 2009



- Beijing Spectrometer (BESIII)
 - Optimized for flavour physics
 - Covers 93% of 4π solid angle
 - 1.0 T super-conducting solenoid
- Momentum resolution:
 $\sigma(p)/p = 0.5\%$ at 1 GeV/c
- Time resolution:
68 (65) ps in the barrel (end cap)

BESIII data sample

- World's largest charmonia sample:
 - $N_{J/\psi} \sim 10^{10}$
 - $N_{\psi(2S)} \sim 3 \cdot 10^9$
- Full baryon-antibaryon octet kinematically accessible

Resonance	Pair	$\epsilon(\%)$	$\mathcal{B}(\cdot 10^{-4})$	Reference
J/ψ	$\Lambda\Lambda$	42.37 ± 0.14	19.43 ± 0.03	[PRD95(2017)052003]
	$\Sigma^0\bar{\Sigma}^0$	17.83 ± 0.06	11.64 ± 0.04	[JHEP11(2021)226]
	$\Sigma^+\bar{\Sigma}^-$	24.1 ± 0.7	10.61 ± 0.04	[PRD93(2016)072003]
	${}^*\Xi^-\bar{\Xi}^+$	18.40 ± 0.04	10.40 ± 0.06	[PLB770(2017)217]
	$\Xi^0\bar{\Xi}^0$	14.05 ± 0.04	11.65 ± 0.04	
$\psi(2S)$	$\Lambda\Lambda$	42.83 ± 0.34	3.97 ± 0.02	[PRD95(2017)052003]
	$\Sigma^0\bar{\Sigma}^0$	14.79 ± 0.12	2.44 ± 0.03	[JHEP11(2021)226]
	$\Sigma^+\bar{\Sigma}^-$	18.6 ± 0.5	2.52 ± 0.04	[PRD93(2016)072003]
	${}^*\Xi^-\bar{\Xi}^+$	18.04 ± 0.04	2.78 ± 0.05	[PLB770(2017)217]
	$\Xi^0\bar{\Xi}^0$	14.10 ± 0.04	2.73 ± 0.03	
	$\Omega^-\bar{\Omega}^+$	$17.1/18.9$	0.59 ± 0.03	[PRL126(2021)092002]

numbers for $1.31 \cdot 10^9 J/\psi$ and $0.45 \cdot 10^9 \psi(2S)$

* numbers for $0.22 \cdot 10^9 J/\psi$ and $0.11 \cdot 10^9 \psi(2S)$

Introduction to CPV

- More than 50 years of the knowledge about CP violation (CPV)
 - Confirmed only in meson decays
- SM CPV is not sufficient to explain observed matter-antimatter asymmetry
- Baryogenesis requires C and CP violation in the processes

[[PismaZh.Eksp.Teor.Fiz.5\(1967\)32](#)]


created by Annika Rockström



- Systematical mapping with different hadronic systems and complementary methods are needed for understanding CPV in flavour sector

Ground-state strange baryons

- Spin- $\frac{1}{2}$ baryon octet
- Weak $\Delta S = 1$ transitions




Hyperon	Mass [GeV/c ²]	Decay (\mathcal{B})
$\Lambda(uds)$	1.116	$p\pi^-$ (64.1%) $n\pi^0$ (35.9%)
$\Sigma^-(dds)$	1.197	$n\pi^-$ (99.8%)
$\Sigma^+(uus)$	1.189	$p\pi^0$ (51.6%) $n\pi^+$ (48.3%)
$\Xi^0(uss)$	1.315	$\Lambda\pi^0$ (99.5%)
$\Xi^-(dss)$	1.322	$\Lambda\pi^-$ (99.9%)
$\Omega^-(sss)$	1.672	ΛK^- (67.8%) $\Xi^0\pi^-$ (23.6%) $\Xi^-\pi^0$ (8.6%)

Modular method to study full process:
 $e^+e^- \rightarrow (c\bar{c}) \rightarrow Y\bar{Y} \rightarrow (BM)(\bar{B}\bar{M})$


Production process

[PRD99(2019)056008]



- Spin $\frac{1}{2} + \frac{1}{2}$ baryon-antibaryon density matrix:

$$\rho_{1/2, \overline{1/2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$



- Spin $\frac{1}{2} + \frac{1}{2}$ baryon-antibaryon density matrix:

$$\rho_{1/2, \overline{1/2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$


$$C_{\mu\bar{\nu}} = \begin{pmatrix} 1 + \alpha_\psi \cos^2 \theta & 0 & \beta_\psi \sin \theta \cos \theta & 0 \\ 0 & \sin^2 \theta & 0 & \gamma_\psi \sin \theta \cos \theta \\ -\beta_\psi \sin \theta \cos \theta & 0 & \alpha_\psi \sin^2 \theta & 0 \\ 0 & -\gamma_\psi \sin \theta \cos \theta & 0 & -\alpha_\psi - \cos^2 \theta \end{pmatrix}$$

Y_1 transverse polarization
 $\beta_\psi \sin \theta \cos \theta$
 Y_1 transverse polarization
spin-correlation terms

$$\beta_\psi = \sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi), \quad \gamma_\psi = \sqrt{1 - \alpha_\psi^2} \cos(\Delta\Phi)$$

Production process

[PRD99(2019)056008]



- Spin $\frac{1}{2} + \frac{1}{2}$ baryon-antibaryon density matrix:

$$\rho_{1/2, \overline{1/2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$

$$C_{\mu\bar{\nu}} = \begin{pmatrix} 1 + \alpha_\psi \cos^2 \theta & 0 & \beta_\psi \sin \theta \cos \theta & 0 \\ 0 & \sin^2 \theta & 0 & \gamma_\psi \sin \theta \cos \theta \\ -\beta_\psi \sin \theta \cos \theta & 0 & \alpha_\psi \sin^2 \theta & 0 \\ 0 & -\gamma_\psi \sin \theta \cos \theta & 0 & -\alpha_\psi - \cos^2 \theta \end{pmatrix}$$

Y_1 transverse polarization spin-correlation terms

$$\beta_\psi = \sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi), \quad \gamma_\psi = \sqrt{1 - \alpha_\psi^2} \cos(\Delta\Phi)$$

- Unpolarised e^+e^- beams \Rightarrow transverse polarisation (if $\Delta\Phi \neq 0$):

$$P_y(\cos \theta) = \frac{\sqrt{1 - \alpha_\psi^2} \cos \theta \sin \theta}{1 + \alpha_\psi \cos^2 \theta} \sin(\Delta\Phi)$$

Decay amplitudes in hyperon decays

- S - and P -wave amplitudes:

$Y \rightarrow BM$ like $\Lambda \rightarrow p\pi^-$

$Y_1 \rightarrow Y_2 (\rightarrow BM) M$ like $\Xi^- \rightarrow \Lambda (\rightarrow p\pi^-)\pi^-$

$$\mathcal{A} = S + P \vec{\sigma} \cdot \hat{\mathbf{n}}$$

- $|\Delta I| = 1/2$
- Contribution of $|\Delta I| = 3/2$ is $\sim 10\%$

weak CP-odd phases

$$S = |S| \exp(\xi_S) \exp(i\delta_S)$$

$$P = |P| \exp(\xi_P) \exp(i\delta_P)$$

strong phases

Decay amplitudes in hyperon decays

- S - and P -wave amplitudes:

$$Y \rightarrow BM \text{ like } \Lambda \rightarrow p\pi^-$$

$$Y_1 \rightarrow Y_2 (\rightarrow BM) M \text{ like } \Xi^- \rightarrow \Lambda (\rightarrow p\pi^-)\pi^-$$

$$\mathcal{A} = S + P \vec{\sigma} \cdot \hat{\mathbf{n}}$$

- $|\Delta I| = 1/2$
- Contribution of $|\Delta I| = 3/2$ is $\sim 10\%$

weak CP-odd phases

$$S = |S| \exp(\xi_S) \exp(i\delta_S)$$

$$P = |P| \exp(\xi_P) \exp(i\delta_P)$$


strong phases

- Two measurable parameters

$$\alpha = \frac{2\text{Re}(S*P)}{|S|^2 + |P|^2}, \quad \beta = \frac{2\text{Im}(S*P)}{|S|^2 + |P|^2} = \sqrt{1 - \alpha^2} \sin \phi$$

Measured hyperon decay parameters


- Polarisation of hyperons is experimentally accessible in weak parity-violating decays



- Example:
angular distribution of $Y \rightarrow BM$

$$I(\cos \theta_B) \propto 1 + \alpha_B P_y \cos \theta_B$$

- Angle ϕ is accessible when polarisation of daughter baryon is measured
 - Example: $Y_1 \rightarrow Y_2 (\rightarrow BM) M$ like $\Xi^- \rightarrow \Lambda (\rightarrow p\pi^-)\pi^-$



CP tests in hyperon decays

- If CP conserved: $\bar{\alpha} = -\alpha$, $\bar{\beta} = -\beta$, $\bar{\phi} = -\phi$
- Possible CP tests:
- SM predictions [PRD105(2022)116022]

weak P-S phase difference

$$A_{CP} = \frac{\alpha + \bar{\alpha}}{\alpha - \bar{\alpha}} = -\sin \phi \tan(\xi_P - \xi_S) \frac{\sqrt{1-\alpha^2}}{\alpha}$$

$$\Phi_{CP} = \frac{\phi + \bar{\phi}}{2} = \cos \phi \tan(\xi_P - \xi_S) \frac{\alpha}{\sqrt{1-\alpha^2}}$$

$$-3 \cdot 10^{-5} \leq A_\Lambda \leq 3 \cdot 10^{-5}$$

$$0.5 \cdot 10^{-5} \leq A_\Xi \leq 6 \cdot 10^{-5}$$

Decay mode	$\xi_P - \xi_S$ [10^{-4} rad]
$\Lambda \rightarrow p\pi^-$	-0.2 ± 2.2
$\Xi^- \rightarrow \Lambda\pi^-$	-2.1 ± 1.7

- HyperCP: $A_{CP}^\Lambda + A_{CP}^\Xi = (0.0 \pm 5.1_{\text{stat}} \pm 4.4_{\text{syst}}) \cdot 10^{-4}$ [PRL93(2004)262001]
- HyperCP^[preliminary]: $A_{CP}^\Lambda + A_{CP}^\Xi = (-6.0 \pm 2.1_{\text{stat}} \pm 2.0_{\text{syst}}) \cdot 10^{-4}$ [BEACH2008]
signal statistics increased by factor 6-7

Joint angular amplitude

[PRD99(2019)056008] [PRD100(2019)114005]

- Production matrix:

$$\rho_{1/2, \overline{1}/\overline{2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$

- Decay matrix:

$$\sigma_\mu^{Y_1} \rightarrow \sum_{\mu'=0}^3 a_{\mu\mu'}^{Y_1}(\alpha_{Y_1}, \phi_{Y_1}; \theta_{Y_2}, \varphi_{Y_2}) \sigma_{\mu'}^{Y_2}$$

Joint angular amplitude

- Production matrix:

$$\rho_{1/2, \bar{1}/\bar{2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$

- Decay matrix:

$$\sigma_\mu^{Y_1} \rightarrow \sum_{\mu'=0}^3 a_{\mu\mu'}^{Y_1}(\alpha_{Y_1}, \phi_{Y_1}; \theta_{Y_2}, \varphi_{Y_2}) \sigma_{\mu'}^{Y_2}$$

- Joint angular amplitude of full decay chain takes into account **polarisation**, **entanglement** and **sequential decays**

$$\mathcal{W}(\xi, \omega) = \sum_{\mu, \bar{\nu}=0}^3 C_{\mu\bar{\nu}} \sum_{\mu', \bar{\nu}'=0}^3 a_{\mu\mu'}^{Y_1} a_{\bar{\nu}\bar{\nu}'}^{\bar{Y}_1} a_{\mu'0}^{Y_2} a_{\bar{\nu}'0}^{\bar{Y}_2}$$

$\xi = (\theta_{Y_1}, \theta_{Y_2}, \varphi_{Y_2}, \bar{\theta}_{Y_2}, \bar{\varphi}_{Y_2}, \theta_B, \varphi_B, \bar{\theta}_B, \bar{\varphi}_B)$ - set of helicity angles

$\omega = (\alpha_\psi, \Delta\Phi, \alpha_{Y_1}, \phi_{Y_1}, \bar{\alpha}_{Y_1}, \bar{\phi}_{Y_1}, \alpha_{Y_2}, \bar{\alpha}_{Y_2})$ - set of measured parameters

Joint angular amplitude

- Production matrix:

$$\rho_{1/2, \bar{1}/\bar{2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$

- Decay matrix:

$$\sigma_\mu^{Y_1} \rightarrow \sum_{\mu'=0}^3 a_{\mu\mu'}^{Y_1} (\alpha_{Y_1}, \phi_{Y_1}; \theta_{Y_2}, \varphi_{Y_2}) \sigma_{\mu'}^{Y_2}$$

- Joint angular amplitude of full decay chain takes into account **polarisation**, **entanglement** and **sequential decays**

$$\mathcal{W}(\xi, \omega) = \sum_{\mu, \bar{\nu}=0}^3 C_{\mu\bar{\nu}} \sum_{\mu', \bar{\nu}'=0}^3 a_{\mu\mu'}^{Y_1} a_{\bar{\nu}\bar{\nu}'}^{\bar{Y}_1} a_{\mu'0}^{Y_2} a_{\bar{\nu}'0}^{\bar{Y}_2}$$

$\xi = (\theta_{Y_1}, \theta_{Y_2}, \varphi_{Y_2}, \bar{\theta}_{Y_2}, \bar{\varphi}_{Y_2}, \theta_B, \varphi_B, \bar{\theta}_B, \bar{\varphi}_B)$ - set of helicity angles

$\omega = (\alpha_\psi, \Delta\Phi, \alpha_{Y_1}, \phi_{Y_1}, \bar{\alpha}_{Y_1}, \bar{\phi}_{Y_1}, \alpha_{Y_2}, \bar{\alpha}_{Y_2})$ - set of measured parameters

- $e^+e^- \rightarrow (c\bar{c}) \rightarrow Y\bar{Y} \rightarrow (BM)(\bar{B}\bar{M})$

- $\xi = 5$ angles, $\omega = 4$ parameters

Joint angular amplitude

- Production matrix:

$$\rho_{1/2, \bar{1}/\bar{2}} = \frac{1}{4} \sum_{\mu\bar{\nu}} C_{\mu\bar{\nu}} \sigma_\mu^{Y_1} \otimes \sigma_{\bar{\nu}}^{\bar{Y}_1}$$

- Decay matrix:

$$\sigma_\mu^{Y_1} \rightarrow \sum_{\mu'=0}^3 a_{\mu\mu'}^{Y_1} (\alpha_{Y_1}, \phi_{Y_1}; \theta_{Y_2}, \varphi_{Y_2}) \sigma_{\mu'}^{Y_2}$$

- Joint angular amplitude of full decay chain takes into account **polarisation**, **entanglement** and **sequential decays**

$$\mathcal{W}(\xi, \omega) = \sum_{\mu, \bar{\nu}=0}^3 C_{\mu\bar{\nu}} \sum_{\mu', \bar{\nu}'=0}^3 a_{\mu\mu'}^{Y_1} a_{\bar{\nu}\bar{\nu}'}^{\bar{Y}_1}$$

$$\xi = (\theta_{Y_1}, \theta_{Y_2}, \varphi_{Y_2}, \bar{\theta}_{Y_2}, \bar{\varphi}_{Y_2}, \theta_B, \varphi_B, \bar{\theta}_B, \bar{\varphi}_B) - \text{set of helicity angles}$$

$$\omega = (\alpha_\psi, \Delta\Phi, \alpha_{Y_1}, \phi_{Y_1}, \bar{\alpha}_{Y_1}, \bar{\phi}_{Y_1}, \alpha_{Y_2}, \bar{\alpha}_{Y_2}) - \text{set of measured parameters}$$

- $e^+e^- \rightarrow (c\bar{c}) \rightarrow Y\bar{Y} \rightarrow (BM)(\bar{B}\bar{M})$
 - $\xi = 5$ angles, $\omega = 4$ parameters
- $e^+e^- \rightarrow (c\bar{c}) \rightarrow Y_1\bar{Y}_1 \rightarrow (Y_2 M_1)(\bar{Y}_2 \bar{M}_1) \rightarrow (BM_2 M_1)(\bar{B}\bar{M}_2 \bar{M}_1)$
 - $\xi = 9$ angles, $\omega = 8$ parameters

Experimental results:

$$e^+ e^- \rightarrow Y\bar{Y} \rightarrow (BM)(\bar{B}\bar{M})$$

$$e^+ e^- \rightarrow J/\psi \rightarrow \Lambda \bar{\Lambda}, \Lambda \rightarrow p \pi^- + \text{c.c.} \quad (1)$$


- Increasing data statistics have allowed for the significant result improvement ($\sim 3\sigma$):

¹[Nature Phys.15(2019)631] \Rightarrow ²[PRL129(2022)131801]

	This work ²	Previous work ¹
$N_{J/\psi}$	10^{10}	$1.31 \cdot 10^9$
N_{sig}	$3.2 \cdot 10^6$	$421 \cdot 10^3$
N_{bkg}	3801 ± 63	399 ± 20

- Angular dependence of the moment for the acceptance-corrected data:

$$\mu(\cos \theta_\Lambda) = \frac{\alpha_\Lambda - \bar{\alpha}_\Lambda}{2} \frac{1 + \alpha_\psi \cos^2 \theta_\Lambda}{3 + \alpha_\psi} P_y(\cos \theta_\Lambda)$$



Parameters	This work ²	Previous results ¹
α_ψ	$0.4748 \pm 0.0022 \pm 0.0024$	$0.461 \pm 0.006 \pm 0.007$
$\Delta\Phi$ [rad]	$0.7521 \pm 0.0042 \pm 0.0080$	$0.740 \pm 0.010 \pm 0.009$
α_Λ	$0.7519 \pm 0.0036 \pm 0.0019$	$0.750 \pm 0.009 \pm 0.004$
$\bar{\alpha}_\Lambda$	$-0.7559 \pm 0.0036 \pm 0.0029$	$-0.758 \pm 0.010 \pm 0.007$


$$e^+e^- \rightarrow J/\psi \rightarrow \Lambda\bar{\Lambda}, \Lambda \rightarrow p\pi^- + \text{c.c.} \quad (2)$$

$$A_{\text{CP}}^{\Lambda} = \frac{\alpha_{\Lambda} + \bar{\alpha}_{\Lambda}}{\alpha_{\Lambda} - \bar{\alpha}_{\Lambda}} = -0.0025 \pm 0.0046_{\text{stat}} \pm 0.0011_{\text{syst}}$$



- BESIII: $A_{\text{CP}}^{\Lambda} = -0.006 \pm 0.012_{\text{stat}} \pm 0.007_{\text{syst}}$ [Nature Phys.15(2019)631]
- PS185: $A_{\text{CP}}^{\Lambda} = 0.013 \pm 0.021_{\text{tot}}$ [PRC54(1996)1877]

$$\langle \alpha_{\Lambda} \rangle = \frac{\alpha_{\Lambda} - \bar{\alpha}_{\Lambda}}{2} = 0.7542 \pm 0.0010_{\text{stat}} \pm 0.0020_{\text{syst}}$$

- BESIII: $\langle \alpha_{\Lambda} \rangle = 0.754 \pm 0.003_{\text{stat}} \pm 0.002_{\text{syst}}$ [Nature Phys.15(2019)631]
- CLAS: $\alpha_{\Lambda} = 0.721 \pm 0.006_{\text{stat}} \pm 0.005_{\text{syst}}$ [PRL123(2019)182301]



$$e^+ e^- \rightarrow J/\psi \rightarrow \Sigma^+ \bar{\Sigma}^-$$



First result

Parameters	$(p\pi^0)(\bar{p}\pi^0)$ ¹	$(p\pi^0)(\bar{n}\pi^-) + \text{c.c.}$ ²
$N_{J/\psi}$	$1.31 \cdot 10^9$	10^{10}
N_{sig}	$87 \cdot 10^3$ with 5% bkg	$(3.1 + 7.5) \cdot 10^5$ with 2% bkg
$\alpha_{J/\psi}$	$-0.508 \pm 0.006 \pm 0.004$	$-0.5156 \pm 0.0030 \pm 0.0061$
$\Delta\Phi_{J/\psi}$ [rad]	$-0.270 \pm 0.012 \pm 0.009$	$-0.2772 \pm 0.0044 \pm 0.0041$
$\langle \alpha_0 \rangle$	$-0.994 \pm 0.004 \pm 0.002$	
$\langle \alpha_+ \rangle$		$0.0506 \pm 0.0026 \pm 0.0019$

³[PRD67(2003)056001]



First result

A_{CP}^0	$-0.004 \pm 0.037 \pm 0.010$	$3.6 \cdot 10^{-6}$ (SM ³)
A_{CP}^+	$3.9 \cdot 10^{-4}$ (SM ³)	$-0.080 \pm 0.052 \pm 0.028$


Experimental results:

$$e^+ e^- \rightarrow Y_1 \bar{Y}_1 \rightarrow (Y_2 M_1)(\bar{Y}_2 \bar{M}_1) \rightarrow (B M_2 M_1)(\bar{B} \bar{M}_2 \bar{M}_1)$$

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+, \Xi^- \rightarrow \Lambda(\rightarrow p\pi^-)\pi^- + \text{c.c.} \quad (1)$$



- Data sample: $1.3 \cdot 10^9 J/\psi$
- $73.2 \cdot 10^3$ events with
 $N_{\text{bkg}} = 199 \pm 17$



Parameter	This work	Previous result
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$ [1]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010 [2]
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad [2]
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007$ rad	–
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$ [3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$ [3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–
$\Delta\phi_{CP}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–
A_{CP}^{Λ}	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$ [3]
$\langle \phi_{\Xi} \rangle$	$0.016 \pm 0.014 \pm 0.007$ rad	

- First measurement of Ξ^- polarisation at e^+e^- collider

¹ [PRD93(2016)072003] ² [PTEP2020(2020)083C01] ³ [PRL129(2022)131801] ⁴ [PRL93(2004)011802]

[Nature 606(2022)64]

- First measurement of Ξ^- polarisation at e^+e^- collider
- First direct determination of all $\Xi^-\bar{\Xi}^+$ decay parameters

Parameter	This work	Previous result
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$ [1]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010 [2]
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad [2]
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007$ rad	–
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$ [3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$ [3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–
$\Delta\phi_{CP}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$ [3]
$\langle\phi_{\Xi}\rangle$	$0.016 \pm 0.014 \pm 0.007$ rad	

¹[PRD93(2016)072003] ²[PTEP2020(2020)083C01] ³[PRL129(2022)131801] ⁴[PRL93(2004)011802]

- First measurement of Ξ^- polarisation at e^+e^- collider
- First direct determination of all $\Xi^-\bar{\Xi}^+$ decay parameters
- Independent measurement of Λ decay parameters
 - Excellent agreement with previous BESIII results

Parameter	This work	Previous result
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$ [1]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010 [2]
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad [2]
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007$ rad	–
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$ [3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$ [3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–
$\Delta\phi_{CP}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$ [3]
$\langle \phi_{\Xi} \rangle$	$0.016 \pm 0.014 \pm 0.007$ rad	

¹ [PRD93(2016)072003] ² [PTEP2020(2020)083C01] ³ [PRL129(2022)131801] ⁴ [PRL93(2004)011802]

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+, \Xi^- \rightarrow \Lambda(\rightarrow p\pi^-)\pi^- + c.c. \quad (2)$$

[Nature 606(2022)64]

- First measurement of Ξ^- polarisation at e^+e^- collider
- First direct determination of all $\Xi^-\bar{\Xi}^+$ decay parameters
- Independent measurement of Λ decay parameters
 - Excellent agreement with previous BESIII results
- Two independent CP tests

Parameter	This work	Previous result
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$ [1]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010 [2]
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad [2]
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007$ rad	–
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$ [3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$ [3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–
$\Delta\phi_{CP}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$ [3]
$\langle \phi_{\Xi} \rangle$	$0.016 \pm 0.014 \pm 0.007$ rad	

¹ [PRD93(2016)072003] ² [PTEP2020(2020)083C01] ³ [PRL129(2022)131801] ⁴ [PRL93(2004)011802]

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+, \Xi^- \rightarrow \Lambda(\rightarrow p\pi^-)\pi^- + c.c. \quad (2)$$

- First measurement of Ξ^- polarisation at e^+e^- collider
- First direct determination of all $\Xi^-\bar{\Xi}^+$ decay parameters
- Independent measurement of Λ decay parameters
 - Excellent agreement with previous BESIII results
- Two independent CP tests
- First measurement of weak phase difference
 $(\xi_P - \xi_S)_{\text{SM}} = (-2.1 \pm 1.7) \cdot 10^{-4} \text{ rad}$
[\[PRD105\(2022\)116022\]](#)

[Nature 606(2022)64]

Parameter	This work	Previous result	
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$	[1]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016 \text{ rad}$	–	
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010	[2]
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009 \text{ rad}$	$-0.037 \pm 0.014 \text{ rad}$	[2]
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–	
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007 \text{ rad}$	–	
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$	[3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$	[3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2} \text{ rad}$	–	
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2} \text{ rad}$	$(10.2 \pm 3.9) \times 10^{-2} \text{ rad}$	[4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–	
$\Delta\phi_{\text{CP}}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3} \text{ rad}$	–	
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$	[3]
$\langle \phi_{\Xi} \rangle$	$0.016 \pm 0.014 \pm 0.007 \text{ rad}$		

¹ [PRD93(2016)072003] ² [PTEP2020(2020)083C01] ³ [PRL129(2022)131801] ⁴ [PRL93(2004)011802]

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+, \Xi^- \rightarrow \Lambda(\rightarrow p\pi^-)\pi^- + c.c. \quad (2)$$

- First measurement of Ξ^- polarisation at e^+e^- collider
- First direct determination of all $\Xi^-\bar{\Xi}^+$ decay parameters
- Independent measurement of Λ decay parameters
 - Excellent agreement with previous BESIII results
- Two independent CP tests
- First measurement of weak phase difference
 $(\xi_P - \xi_S)_{SM} = (-2.1 \pm 1.7) \cdot 10^{-4}$ rad
[\[PRD105\(2022\)116022\]](#)

Parameter	This work	Previous result
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–
α_{Ξ}	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010
ϕ_{Ξ}	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad
$\bar{\alpha}_{\Xi}$	$0.371 \pm 0.007 \pm 0.002$	–
$\bar{\phi}_{\Xi}$	$-0.021 \pm 0.019 \pm 0.007$ rad	–
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.7519 \pm 0.0036 \pm 0.0019$ [3]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.7559 \pm 0.0036 \pm 0.0029$ [3]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [4]
A_{CP}^{Ξ}	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–
$\Delta\phi_{CP}^{\Xi}$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-2.5 \pm 4.6 \pm 1.1) \times 10^{-3}$ [3]
$\langle \phi_{\Xi} \rangle$	$0.016 \pm 0.014 \pm 0.007$ rad	


¹[PRD93(2016)072003] ²[PTEP2020(2020)083C01] ³[PRL129(2022)131801] ⁴[PRL93(2004)011802]

- Study of $e^+e^- \rightarrow J/\psi \rightarrow \Xi^0\bar{\Xi}^0, \Xi^0 \rightarrow \Lambda(\rightarrow p\pi^-)\pi^0 + c.c.$ [PRD108(2023)L031106]
 - Data sample: $10^{10} J/\psi$
 - First measurement of Ξ^0 polarization: $\Delta\Phi = 1.168 \pm 0.019 \pm 0.018$
 - Improved result of weak phase difference: $(0.0 \pm 1.7 \pm 0.2) \times 10^{-2}$

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+ \rightarrow (\Lambda(p\pi^-)\pi^-)(\bar{\Lambda}(\bar{n}\pi^0)\pi^+) + \text{c.c.} \quad (1)$$

- Data sample: $10^{10} J/\psi$
- $144 \cdot 10^3$ and $123 \cdot 10^3$ events for $(p2\pi^-)(\bar{n}\pi^0\pi^+)$ and $(n\pi^0\pi^-)(\bar{p}2\pi^+)$, respectively
- Result is consistent with $\Xi^-\bar{\Xi}^+ \rightarrow (p2\pi^-)(\bar{p}2\pi^+)$

[Nature 606(2022)64]



Parameters	This work	Previous result
α_0/α_-	$0.877 \pm 0.015^{+0.014}_{-0.010}$	1.01 ± 0.07 [1]
$\bar{\alpha}_0/\alpha_+$	$0.863 \pm 0.014^{+0.012}_{-0.008}$	$0.913 \pm 0.028 \pm 0.012$ [2]

¹[PTEP2022(2022)083C01] ²[Nature Phys.15(2019)631] ³[PRL129(2022)131801]


$$A_{\text{CP}}^- = \frac{\alpha_- + \alpha_+}{\alpha_- - \alpha_+} = -0.007 \pm 0.008^{+0.002}_{-0.003} \quad * [3]$$

$$A_{\text{CP}}^0 = \frac{\alpha_0 + \bar{\alpha}_0}{\alpha_0 - \bar{\alpha}_0} = 0.001 \pm 0.009^{+0.005}_{-0.007}$$

$$A_{\text{CP}}^\Lambda = (2A_{\text{CP}}^- + A_{\text{CP}}^0)/3 = -0.004 \pm 0.007^{+0.003}_{-0.004}$$

- Data sample:

- $10^{10} J/\psi$ with $N_{\text{sig}} \sim 1.1 \cdot 10^6$
- $2.7 \cdot 10^9 \psi(2S)$ with $N_{\text{sig}} \sim 52 \cdot 10^3$



Parameters	This work	Previous result ¹
$\alpha_{J/\psi}$	$-0.4133 \pm 0.0035 \pm 0.0077$	-0.449 ± 0.022
$\Delta\Phi_{J/\psi}$ [rad]	$-0.0828 \pm 0.0068 \pm 0.0033$	—
$\alpha_{\psi(2S)}$	$0.814 \pm 0.028 \pm 0.028$	0.71 ± 0.12
$\Delta\Phi_{\psi(2S)}$ [rad]	$0.512 \pm 0.085 \pm 0.034$	—
α_{Σ^0}	$-0.0017 \pm 0.0021 \pm 0.0018$	—
$\bar{\alpha}_{\Sigma^0}$	$0.0021 \pm 0.0020 \pm 0.0022$	—


First result

¹[PRD95(2017)052003]

$e^+e^- \rightarrow$

- Data samples
- $10^{10} J/\psi$
- $2.7 \cdot 10^{10} \psi(2S)$


First result
First result



Summary and Outlook

- BESIII has performed

- Measurements of **polarisation** and **spin correlations**
 - * $\Lambda\bar{\Lambda}, \Sigma^0\bar{\Sigma}^0, \Sigma^+\bar{\Sigma}^-, \Xi^0\bar{\Xi}^0, \Xi^+\bar{\Xi}^-$
using partial and full J/ψ and $\psi(2S)$ statistics
- Determination of **hyperon and antihyperon decay parameters**
- **CP tests** comparing hyperon and antihyperon
 - * Separation of strong and weak decay phases
 \implies more **sensitive** CP tests




Summary and Outlook

- BESIII has performed

- Measurements of **polarisation and spin correlations**
 - * $\Lambda\bar{\Lambda}, \Sigma^0\bar{\Sigma}^0, \Sigma^+\bar{\Sigma}^-, \Xi^0\bar{\Xi}^0, \Xi^+\bar{\Xi}^-$
using partial and full J/ψ and $\psi(2S)$ statistics
- Determination of **hyperon and antihyperon decay parameters**
- **CP tests** comparing hyperon and antihyperon
 - * Separation of strong and weak decay phases
 \Rightarrow more **sensitive** CP tests

- Future prospects with BESIII

- **More interesting results** are coming using full collected statistics:
 10^{10} J/ψ and $3 \cdot 10^9 \psi(2S)$
- BEPCII upgrade in 2024-25 with increasing $E_{ee} \in (4...5)$ GeV



Summary and Outlook

- BESIII has performed


- Measurements of **polarisation and spin correlations**
 - * $\Lambda\bar{\Lambda}, \Sigma^0\bar{\Sigma}^0, \Sigma^+\bar{\Sigma}^-, \Xi^0\bar{\Xi}^0, \Xi^+\bar{\Xi}^-$
using partial and full J/ψ and $\psi(2S)$ statistics
- Determination of **hyperon and antihyperon decay parameters**
- **CP tests** comparing hyperon and antihyperon
 - * Separation of strong and weak decay phases
 \Rightarrow more **sensitive** CP tests

- Future prospects with BESIII

- **More interesting results** are coming using full collected statistics:
 10^{10} J/ψ and $3 \cdot 10^9 \psi(2S)$
- BEPCII upgrade in 2024-25 with increasing $E_{ee} \in (4...5)$ GeV

- Longer time scale prospects with STCF [FrontPhys(Beijing)19(2024)14701] [PRD105(2022)116022]

- Planning produce more than 10^{12} J/ψ events
- Polarized electron beam
- Statistical precision will be comparable to the SM predictions



Summary and Outlook

- BESIII has performed


- Measurements of **polarisation** and **spin correlations**
 - * $\Lambda\bar{\Lambda}, \Sigma^0\bar{\Sigma}^0, \Sigma^+\bar{\Sigma}^-, \Xi^0\bar{\Xi}^0, \Xi^+\bar{\Xi}^-$
using partial and full J/ψ and $\psi(2S)$ statistics
- Determination of **hyperon and antihyperon decay parameters**
- **CP tests** comparing hyperon and antihyperon
 - * Separation of strong and weak decay phases
 \Rightarrow more **sensitive** CP tests

- Future prospects with BESIII

- More interesting results are coming using full collected statistics:
 10^{10} J/ψ and $3 \cdot 10^9 \psi(2S)$
- BEPCII upgrade in 2024-25 with increasing $E_{ee} \in (4...5)$ GeV

- Longer time scale prospects with STCF [FrontPhys(Beijing)19(2024)14701] [PRD105(2022)116022]

- * Planning produce more than 10^{12} J/ψ events
- * Polarized electron beam
- * Statistical precision will be comparable to the SM predictions




Stay tuned!



"I ALWAYS BACK UP EVERYTHING."

Complementarity of hyperon and kaon decays



$$(\xi_P - \xi_S)_{\text{BSM}} = \frac{C'_B}{B_G} \left(\frac{\epsilon'}{\epsilon} \right)_{\text{BSM}} + \frac{C_B}{\kappa} \epsilon_{\text{BSM}}$$

[PRD69(2004)076008]

- BSM predictions [PRD105(2022)116022]



$$|A_\Lambda + A_\Xi| \leq 11 \cdot 10^{-4}$$

Decay mode	$ \xi_P - \xi_S $ [rad]	C_B	C'_B
$\Lambda \rightarrow p\pi^-$	$\leq 5.3 \cdot 10^{-3}$	0.9 ± 1.8	0.4 ± 0.9
$\Xi^- \rightarrow \Lambda\pi^-$	$\leq 3.7 \cdot 10^{-3}$	-0.5 ± 1.0	0.4 ± 0.7

with $0.5 < B_G < 2$ and $0.2 < |\kappa| < 1$ [PRD61(2000)071701]

$|\epsilon'/\epsilon|_{\text{BSM}} \leq 1 \cdot 10^{-3}$ and $|\epsilon|_{\text{BSM}} \leq 2 \cdot 10^{-4}$ [JHEP12(2020)097]

$$e^+ e^- \rightarrow J/\psi \rightarrow \Sigma^+ \bar{\Sigma}^- \rightarrow (p\gamma)(\bar{p}\pi^0) + c.c.$$



- Data sample: 10^{10} J/ψ
- 1189 ± 38 and 1306 ± 39 events for $(p\gamma)(\bar{p}\pi^0)$ and $(p\pi^0)(\bar{p}\gamma)$, respectively

$$\mathcal{B} = (0.996 \pm 0.021 \pm 0.018) \cdot 10^{-3}$$

$$\langle \alpha_\gamma \rangle = -0.651 \pm 0.056 \pm 0.020$$

$$\Delta_{CP} = \frac{\mathcal{B} - \bar{\mathcal{B}}}{\mathcal{B} + \bar{\mathcal{B}}} = 0.006 \pm 0.011 \pm 0.004$$

$$A_{CP} = \frac{\bar{\alpha}_\gamma + \alpha_\gamma}{\bar{\alpha}_\gamma - \alpha_\gamma} = 0.095 \pm 0.087 \pm 0.018$$

- Data sample: $10^{10} J/\psi$
- $3.3 \cdot 10^5$ events with 2% bkg
- First measurement of Ξ^0 polarisation
- Improved measurement:
 - All $\Xi^0\bar{\Xi}^0$ decay parameters
 - Weak phase difference
 - Two independent CP tests

Parameter	This work	Previous result	
$\alpha_{J/\psi}$	$0.514 \pm 0.006 \pm 0.015$	0.66 ± 0.06	[1]
$\Delta\Phi(\text{rad})$	$1.168 \pm 0.019 \pm 0.018$	-	
α_{Ξ}	$-0.3750 \pm 0.0034 \pm 0.0016$	-0.358 ± 0.044	[2]
$\bar{\alpha}_{\Xi}$	$0.3790 \pm 0.0034 \pm 0.0021$	0.363 ± 0.043	[2]
$\phi_{\Xi}(\text{rad})$	$0.0051 \pm 0.0096 \pm 0.0018$	0.03 ± 0.12	[2]
$\bar{\phi}_{\Xi}(\text{rad})$	$-0.0053 \pm 0.0097 \pm 0.0019$	-0.19 ± 0.13	[2]
α_{Λ}	$0.7551 \pm 0.0052 \pm 0.0023$	0.7519 ± 0.0043	[3]
$\bar{\alpha}_{\Lambda}$	$-0.7448 \pm 0.0052 \pm 0.0017$	-0.7559 ± 0.0047	[3]
$\xi_P - \xi_S(\text{rad})$	$(0.0 \pm 1.7 \pm 0.2) \times 10^{-2}$	-	
$\delta_P - \delta_S(\text{rad})$	$(-1.3 \pm 1.7 \pm 0.4) \times 10^{-2}$	-	
A_{CP}^{Ξ}	$(-5.4 \pm 6.5 \pm 3.1) \times 10^{-3}$	$(-0.7 \pm 8.5) \times 10^{-2}$	[2]
$\Delta\phi_{CP}^{\Xi}(\text{rad})$	$(-0.1 \pm 6.9 \pm 0.9) \times 10^{-3}$	$(-7.9 \pm 8.3) \times 10^{-2}$	[2]
A_{CP}^{Λ}	$(6.9 \pm 5.8 \pm 1.8) \times 10^{-3}$	$(-2.5 \pm 4.8) \times 10^{-3}$	[3]
$\langle \alpha_{\Xi} \rangle$	$-0.3770 \pm 0.0024 \pm 0.0014$	-	
$\langle \phi_{\Xi} \rangle(\text{rad})$	$0.0052 \pm 0.0069 \pm 0.0016$	-	
$\langle \alpha_{\Lambda} \rangle$	$0.7499 \pm 0.0029 \pm 0.0013$	0.7542 ± 0.0026	[3]

¹[PLB770(2017)217] ³[PRL129(2022)131801]

²[PRD108(2023)L011101] $\psi(2S) \rightarrow \Xi^0\bar{\Xi}^0$, $0.45 \cdot 10^9 \psi(2S)$, $N_{\text{sig}} \sim 2 \cdot 10^3$ with 1.2% bkg

$$e^+ e^- \rightarrow J/\psi \rightarrow \Xi^- \bar{\Xi}^+ \rightarrow (\Lambda(p\pi^-)\pi^-)(\bar{\Lambda}(\bar{n}\pi^0)\pi^+) + \text{c.c.} \quad (2)$$

[PRL132(2024)101801]

- If CP is conserved,

$$\alpha_- \alpha_{\Xi} = \alpha_+ \bar{\alpha}_{\Xi} \text{ and}$$

$$R(\cos \theta_i, \cos \theta_{\bar{i}}) = \frac{1 + \alpha_- \alpha_{\Xi} \cos \theta_i}{1 + \alpha_+ \alpha_{\Xi} \cos \theta_{\bar{i}}} \text{ with}$$

$i = \{p, n\}$ are flat and equal to unity

- If no $\Delta I = 3/2$ transition in Λ decay,

$$\alpha_- = \alpha_0 \text{ and}$$

$$R(\cos \theta_{\vec{n}}, \cos \theta_{\vec{p}}) = \frac{1 + \overset{(n)}{\alpha}_0 \overset{(n)}{\alpha}_{\Xi} \cos \theta_{\vec{n}}}{1 + \overset{(n)}{\alpha}_- \overset{(n)}{\alpha}_{\Xi} \cos \theta_{\vec{p}}}$$

are flat and equal to unity

- If CP is conserved,

$$\alpha_- \alpha_{\Xi} = \alpha_+ \bar{\alpha}_{\Xi} \text{ and}$$

$$R(\cos \theta_i, \cos \theta_{\bar{i}}) = \frac{1 + \alpha_- \alpha_{\Xi} \cos \theta_i}{1 + \alpha_+ \bar{\alpha}_{\Xi} \cos \theta_{\bar{i}}} \text{ with}$$

$i = \{p, n\}$ are flat and equal to unity

- Consistent with CP symmetry test

$$R_1 = \frac{1 + \alpha_{\Lambda} \alpha_{\Xi} \cos \theta}{1 + \bar{\alpha}_{\Lambda} \bar{\alpha}_{\Xi} \cos \theta} \text{ with } \alpha_{\Lambda} = (2\alpha_- + \alpha_0)/3$$

- If no $\Delta I = 3/2$ transition in Λ decay,

$$\alpha_- = \alpha_0 \text{ and}$$

$$R(\cos \theta_{\bar{n}}, \cos \theta_{\bar{p}}) = \frac{1 + \bar{\alpha}_0 \bar{\alpha}_{\Xi} \cos \theta_{\bar{n}}}{1 + \bar{\alpha}_- \bar{\alpha}_{\Xi} \cos \theta_{\bar{p}}}$$

are flat and equal to unity

- Indication of $\Delta I = 3/2$ contribution
in Λ decay

$$R_2 = \frac{1 + \langle \alpha_0 \rangle \langle \alpha_{\Xi} \rangle \cos \theta}{1 + \langle \alpha_- \rangle \langle \alpha_{\Xi} \rangle \cos \theta}$$

$$e^+e^- \rightarrow J/\psi \rightarrow \Xi^-\bar{\Xi}^+ \rightarrow (\Lambda(p\pi^-)\pi^-)(\bar{\Lambda}(\bar{n}\pi^0)\pi^+) + \text{c.c.} \quad (2)$$

- If CP is conserved,

$$\alpha_- - \alpha_{\Xi} = \alpha_+ - \bar{\alpha}_{\Xi} \text{ and}$$

$$R(\cos \theta_i, \cos \theta_{\bar{i}}) = \frac{1 + \alpha_- - \alpha_{\Xi} \cos \theta_i}{1 + \alpha_+ - \bar{\alpha}_{\Xi} \cos \theta_{\bar{i}}} \text{ with}$$

$i = \{p, n\}$ are flat and equal to unity

- Consistent with CP symmetry test

$$R_1 = \frac{1 + \alpha_{\Lambda} \alpha_{\Xi} \cos \theta}{1 + \bar{\alpha}_{\Lambda} \bar{\alpha}_{\Xi} \cos \theta} \text{ with } \alpha_{\Lambda} = (2\alpha_- + \alpha_0)/3$$



- If no $\Delta I = 3/2$ transition in Λ decay,
 $\alpha_- = \alpha_0$ and

$$R(\cos \theta_{\bar{n}}, \cos \theta_{\bar{p}}) = \frac{1 + \bar{\alpha}_0 \bar{\alpha}_{\Xi} \cos \theta_{\bar{n}}}{1 + \bar{\alpha}_- \bar{\alpha}_{\Xi} \cos \theta_{\bar{p}}}$$

are flat and equal to unity

- Indication of $\Delta I = 3/2$ contribution
in Λ decay

$$R_2 = \frac{1 + \langle \alpha_0 \rangle \langle \alpha_{\Xi} \rangle \cos \theta}{1 + \langle \alpha_- \rangle \langle \alpha_{\Xi} \rangle \cos \theta}$$



$$(\Delta I = 3/2) / (\Delta I = 1/2)$$

$$\text{in } S \text{ wave: } 0.0349 \pm 0.0017^{+0.0012}_{-0.0013} \pm 0.0047$$

$$\text{in } P \text{ wave: } -0.0752 \pm 0.0078^{+0.0067}_{-0.0062} \pm 0.0044$$

constraint for IQCD [PRD102(2020)054509] and dual QCD [EPJC74(2014)2871] approach