



How well do we know the vector quarkonia?

Nils Hüsken
Johannes Gutenberg-Universität Mainz
DPG Spring Meeting Cologne, March 12th 2025




work done with/by the BESIII collaboration,
R. Mitchell, E. Swanson, R. Lebed, Y.Q. Wang, C.Z. Yuan

50 years of charmonium (just last year)




the ψ as seen at SLAC




S. Ting with the J particle


Understanding hadrons





Understanding hadrons



Understanding hadrons




Understanding hadrons




there are many different hadrons

Understanding hadrons



there are many different hadrons

Understanding hadrons




there are many, many (...) different hadrons



variety & patterns also teach us about the strong interaction

→ hadron spectroscopy

Understanding hadrons





Why quarkonium?



Quarkonium

quarkonium: $Q\bar{Q}$ -meson (where $Q = c, b$)

strong interaction version of positronium (or the hydrogen atom)




we can calculate the spectrum using similar methods

Godfrey & Isgur, PRD 32 (1985) 189-231





Barnes, Godfrey, Swanson, PRD 72 (2005) 054026

Godfrey & Moates, PRD 92 (2015) 054034






from Povh, Rith, Scholz, Zetsche, Rodejohann, Teilchen und Kerne

Charmonium





from Povh, Rith, Scholz, Zetsche, Rodejohann, Teilchen und Kerne

Charmonium




Belle, PRL 91 (2003) 262001


Charmonium?




Charmonium?



Vector charmonium




$D\bar{D}$





this decay
should dominate

Vector charmonium




this decay
should dominate


$D\bar{D}$



$$e^+e^- \rightarrow J/\psi \pi^+ \pi^-$$







Vector charmonium




this decay
should dominate





$D\bar{D}$



Vector charmonium





this decay
should dominate




Vector charmonium

$$e^+e^- \rightarrow \text{hadrons}$$





us with the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$

S. Ting with the J particle








Vector charmonium





us with the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$

$e^+e^- \rightarrow \text{hadrons}$








Vector charmonium




us with the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$

can we reliably extract information on vector charmonia in this way?



$e^+e^- \rightarrow \text{hadrons}$




How about vector bottomonium?




Bottomonium




Vector bottomonium





Vector bottomonium




$B\bar{B}$



$$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$$




Vector bottomonium






$B\bar{B}$



this decay
should dominate




$$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$$



Vector bottomonium



Vector bottomonium



$B\bar{B}$

PRL54 (1985) 381

⁹The choice of Gaussian peak shape is arbitrary and is made only for convenience in fitting. Theory predicts not Breit-Wigner peaks but rather irregular shapes, sensitive to assumptions about decay channels and thresholds; see, for example, E. Eichten *et al.*, Phys. Rev. D 21, 203 (1980).


PRL102 (2009) 012001

The number of states is, *a priori*, unknown as are their energy dependencies. Therefore, a proper coupled channel approach [16,17] including the effects of the various thresholds outlined earlier would be likely to modify the results obtained from our simple fit. As an illustration of

CPC 44 (2020) no.8 083001



also complicated. The results from these fits may change dramatically by including more information on each exclusive mode.

The data obtained in this analysis can be used to extract resonant parameters of these states if a better parametrization of the cross sections is developed.




The database: $e^+e^- \rightarrow b\bar{b}$

all exclusive data is from Belle
inclusive data from Belle & BaBar




The database: $e^+e^- \rightarrow b\bar{b}$

all exclusive data is from Belle
inclusive data from Belle & BaBar



two-body open-flavour




$(b\bar{b})\pi\pi$ three-body
channels (small)

$e^+e^- \rightarrow b\bar{b}$:
inclusive (Σ of all)


$B^{(*)}\bar{B}^{(*)}\pi$ multi-
body not measured

How to improve: coupled channel analysis



Coupled channel analysis

from the experiment, we have an almost complete picture of bottom hadron production in e^+e^- annihilation




idea: use all the data to learn more about the intermediate vector bottomonium resonances

Coupled channel analysis


the most simple model we could think of:

$$M = b + BW(s)$$




Coupled channel analysis

unfortunately, that is insufficient:




we could have an intermediate pair of hadrons $j\bar{j}$


Coupled channel analysis




Coupled channel analysis



Coupled channel analysis





Coupled channel analysis



Coupled channel analysis

$$\sigma(e^+e^- \rightarrow b\bar{b}) - \sum_f \sigma(e^+e^- \rightarrow f)$$

unmeasured processes




unitarity:

$$\sigma_{b\bar{b}} = \sum_f \sigma(e^+e^- \rightarrow f)$$

here, this is the sum of all exclusive processes

The resonances

resonance:
pole of matrix element
 $M(s)$ at $\sqrt{s} = m - i\frac{\Gamma}{2}$




What do we learn?

- for the first time, decay branching fractions are determined

→ sensitive input for interpretation



versus




- we can improve upon old measurements


→ accounting for non-resonant processes is important!

- we can provide some guidance for future measurements


→ 80% of $\Upsilon(11020)$ peak intensity has not been measured yet!




example: decays of the $\Upsilon(11020)$



Charmonium: A piece of the puzzle





The case of $G(3900)$




The case of $G(3900)$






Eichten et al., Phys. Rev. D 21 (1980) 203



In our calculation there is some weak structure in the 3.9–4.0 GeV region. It does not arise from a $c\bar{c}$ resonance, but from the opening of the $D\bar{D}^* + D^*\bar{D}$ channel and a decrease in the $D\bar{D}$ channel due to a nearby zero in the $3S$ decay amplitude.



The case of $G(3900)$








— fit result
 68% CL
 90% CL

Data from:

BESIII (unoff.) Andy Julin, University of Minnesota
 BESIII: PRL 133 (2024) 8, 081901
 BESIII: JHEP 05 (2022) 155
 Belle: Phys.Rev.D 97 (2018) 1, 012002
 CLEO: Phys.Rev.D 80 (2009) 072001
 BES: PRL 88, 101802 (2002)
 BESII: PRL 97, 262001 (2006)
 SPEAR: PRL 39, 526 (1977);
 A. Osterheld et al. 86; Schindler 79

The case of $G(3900)$




— fit result
 68% CL
 90% CL


Data from:

BESIII (unoff.) Andy Julin, University of Minnesota
 BESIII: PRL 133 (2024) 8, 081901
 BESIII: JHEP 05 (2022) 155
 Belle: Phys.Rev.D 97 (2018) 1, 012002
 CLEO: Phys.Rev.D 80 (2009) 072001
 BES: PRL 88, 101802 (2002)
 BESII: PRL 97, 262001 (2006)
 SPEAR: PRL 39, 526 (1977);
 A. Osterheld et al. 86; Schindler 79

The case of $G(3900)$



- we find no $G(3900)$ pole!
- effects from the opening of other channels matter when interpreting the data!





(others disagree: Lin et al., PRL 133, 241903)

Summary

How well do we know vector quarkonia?

- a lot of effort has gone into finding new exotic hadrons - but surprisingly little is known about regular vector quarkonia
- we have the necessary data, but there is no free lunch:
 - this is no bump-hunt, simple interpretations tend to fail
 - coupled channel effects matter & global analyses are key - but hard!
- the future is bright: BESIII and Belle-II keep producing high quality data



- very active field with many different approaches - some examples:
- many things still to be learned!
do $\psi(4160)$, $\psi(4230)$, $\psi(4360)$ all exist?

Lin et al., PRL 133, 241903
Nakamura et al., arXiv:2312.17658 [hep-ph]
Cleven et al., PRD 90 (2014) 7, 074039
L. von Detten, PRD 109 (2024) 11, 116002

**Thank you for
your attention!**