


3 QFT analysis of $\pi\pi \rightarrow \pi\pi$ and $\pi\pi \rightarrow \pi\pi\gamma$

We consider the reaction, both on-shell and off shell,

$$\pi^-(\tilde{p}_a) + \pi^0(\tilde{p}_b) \longrightarrow \pi^-(\tilde{p}_1) + \pi^0(\tilde{p}_2) \quad (3.1)$$



We have always energy - momentum conservation

$$\tilde{p}_a + \tilde{p}_b = \tilde{p}_1 + \tilde{p}_2 \quad (3.2)$$

In relations which hold both on- and off shell we denote the momenta by $\tilde{p}_a, \dots, \tilde{p}_2$.

As kinematic variables we have the masses of the, in general off shell, pions, an energy and a momentum transfer variable.

$$\begin{aligned}\tilde{v} &= \tilde{p}_a \cdot \tilde{p}_b + \tilde{p}_1 \cdot \tilde{p}_2, & \tilde{t} &= (\tilde{p}_a - \tilde{p}_1)^2 = (\tilde{p}_b - \tilde{p}_2)^2, \\ m_a^2 &= \tilde{p}_a^2, & m_b^2 &= \tilde{p}_b^2, & m_1^2 &= \tilde{p}_1^2, & m_2^2 &= \tilde{p}_2^2.\end{aligned}\quad (3.3)$$

Following Low we use here $\tilde{v}$ instead of Mandelstam's variable

$$\tilde{s} = \tilde{v} + \frac{1}{2} (m_a^2 + m_b^2 + m_1^2 + m_2^2). \quad (3.4)$$

The scattering amplitude for $\pi^- \pi^0 \rightarrow \pi^- \pi^0$ can only depend on the above variables

$$T(\tilde{p}_a, \tilde{p}_b, \tilde{p}_1, \tilde{p}_2) = M(\tilde{v}, \tilde{t}, m_a^2, m_b^2, m_1^2, m_2^2). \quad (3.5)$$

For the on-shell amplitude we have $\tilde{p}_a \rightarrow p_a, \dots, \tilde{p}_2 \rightarrow p_2,$

$$m_a^2 = m_b^2 = m_1^2 = m_2^2 = m_\pi^2, \quad \tilde{v} \rightarrow v, \quad \tilde{t} \rightarrow t,$$

$$T(p_a, p_b, p_1, p_2) \big|_{\text{on shell}} = \mathcal{M}(\nu, t, m_\pi^2, m_\pi^2, m_\pi^2, m_\pi^2) \\ \equiv \mathcal{M}^{(\text{on})}(\nu, t).$$

(3.6)

Now we come to the photon-emission reaction (on shell)

$$\pi^-(p_a) + \pi^0(p_b) \longrightarrow \pi^-(p_1') + \pi^0(p_2') + \gamma(k, \varepsilon) \quad (3.7)$$

where we have from energy-momentum conservation

$$p_a + p_b = p_1' + p_2' + k. \quad (3.8)$$

Note that for $k \neq 0$ we must have a change of p_1', p_2' from the values p_1, p_2 for $k = 0$.




The amplitude for the above reaction is

$$\langle \gamma(k, \varepsilon), \pi^-(p_1'), \pi^0(p_2') | T | \pi^-(p_a), \pi^0(p_b) \rangle = \varepsilon^{\lambda*} \mathcal{M}_\lambda. \quad (3.9)$$


In the following we consider $\mathcal{M}_\lambda$ for real and timelike virtual photons

$$\mathcal{M}_\lambda = \mathcal{M}_\lambda(p_a, p_b, p_1', p_2', k), \quad k^2 \geq 0, k^0 \geq 0. \quad (3.10)$$


Diagrams for m_λ



pion propagator:



vertex $\gamma \pi \pi$:



$ie \Gamma_\lambda(p', p)$

(3.11)

(a), (b) 1 particle reducible

(c) 1 particle irreducible

With the $\pi\pi \rightarrow \pi\pi$ off-shell amplitude, the pion propagator, and the $\gamma\pi\pi$ vertex we get $m_\lambda = m_\lambda^{(a)} + m_\lambda^{(b)} + m_\lambda^{(c)}$, where

$$m_\lambda^{(a)} = -e m^{(a)} \Delta[(p_a - k)^2] \Gamma_\lambda(p_a - k, p_a),$$

$$\begin{aligned} m^{(a)} &= T(p_a - k, p_b, p_1', p_2') \Big|_{\text{off shell}} \\ &= \mathcal{M}[(p_a - k, p_b) + p_1' \cdot p_2', (p_b - p_2')^2, (p_a - k)^2, m_\pi^2, m_\pi^2, m_\pi^2], \end{aligned}$$

$$m_\lambda^{(b)} = -e \Gamma_\lambda(p_1', p_1' + k) \Delta[(p_1' + k)^2] m^{(b)}, \quad (3.12)$$

$$\begin{aligned} m^{(b)} &= T(p_a, p_b, p_1' + k, p_2') \Big|_{\text{off shell}} \\ &= \mathcal{M}[p_a \cdot p_b + (p_1' + k, p_2'), (p_b - p_2')^2, m_\pi^2, m_\pi^2, (p_1' + k)^2, m_\pi^2]. \end{aligned}$$

(3.13)

We shall now use the best tool from QFT: gauge invariance.

With this we get the Ward-Takahashi identity

$$(p' - p)^\lambda \Gamma_\lambda(p', p) = \Delta^{-1}(p'^2) - \Delta^{-1}(p^2) \quad (3.14)$$

and the condition

$$k^\lambda \left[m_\lambda^{(a)} + m_\lambda^{(b)} + m_\lambda^{(c)} \right] = 0. \quad (3.15)$$

As a consequence of these we find


$$k^\lambda m_\lambda^{(c)} = -e m^{(a)} + e m^{(b)}. \quad (3.16)$$

4 Soft photon theorem I

In this section we shall give the expansion of the amplitude $\mathcal{M}_\lambda$ around the phase-space point $(k=0, \hat{p}_1' = \hat{p}_1)$. In a small neighbourhood of this phase-space point we set:

$$\hat{p}_1' = \hat{p}_1 - \vec{\ell}_{1\perp} / |\vec{p}_1|, \quad \vec{\ell}_{1\perp} \cdot \hat{p}_1 = 0, \quad k, \vec{\ell}_{1\perp} = \mathcal{O}(\omega). \quad (4.1)$$

This neighbourhood has 6 dimensions. Schematically we represent it as follows:



Below we shall study $\mathcal{M}_\lambda$ on such a ray starting from the origin.


For $(k=0, \vec{l}_{1\perp}=0)$ we have the kinematics of the reaction without radiation, $\pi^-(p_a) + \pi^0(p_b) \rightarrow \pi^-(p_1) + \pi^0(p_2)$. For $(k, \vec{l}_{1\perp})$ we have the kinematics of photon radiation with

$$\begin{aligned} p_1' &= p_1 - l_1, & p_2' &= p_2 - l_2, \\ l_1 + l_2 &= k, \end{aligned} \tag{4.2}$$

as required by energy-momentum conservation

$$p_a + p_b = p_1' + p_2' + k = p_1 + p_2.$$

Working in the c.m. system we find



$$p_1 = \begin{pmatrix} p_1^0 \\ |\vec{p}_1| \hat{p}_1 \end{pmatrix}, \quad p_2 = \begin{pmatrix} p_1^0 \\ -|\vec{p}_1| \hat{p}_1 \end{pmatrix}$$

$$s = (p_1 + p_2)^2, \quad p_1^0 = \frac{1}{2} \sqrt{s}, \quad |\vec{p}_1| = \sqrt{\frac{s}{4} - m_\pi^2}, \quad |\hat{p}_1| = 1, \quad (4.3)$$

$$l_1 = \begin{pmatrix} \frac{p_2 \cdot k}{\sqrt{s}} \\ \frac{p_1^0}{|\vec{p}_1| \sqrt{s}} (p_2 \cdot k) \hat{p}_1 + \vec{l}_{1\perp} \end{pmatrix} + \mathcal{O}(\omega^2), \quad l_2 = \begin{pmatrix} \frac{p_1 \cdot k}{\sqrt{s}} \\ \vec{k} - \frac{p_1^0 \hat{p}_1}{|\vec{p}_1| \sqrt{s}} (p_2 \cdot k) - \vec{l}_{1\perp} \end{pmatrix} + \mathcal{O}(\omega^2)$$

$$p_1 \cdot l_1 = 0 + \mathcal{O}(\omega^2), \quad p_2 \cdot l_2 = 0 + \mathcal{O}(\omega^2). \quad (4.4)$$


Now we come to the expansion of the amplitude M_λ for $\omega \rightarrow 0$.
To be precise we set, in the c.m. system with $\omega \geq 0$:

$$k = \omega \begin{pmatrix} 1 \\ \vec{\tilde{k}} \end{pmatrix}, \quad |\vec{\tilde{k}}| \leq 1, \quad \vec{l}_{1\perp} = \omega \vec{\tilde{l}}_{1\perp}, \quad |\vec{\tilde{l}}_{1\perp}| = O(1). \quad (4.5)$$

We keep $\vec{\tilde{k}}$ and $\vec{\tilde{l}}_{1\perp}$ fixed and consider the expansion of the radiative amplitude for $\omega \rightarrow 0$. That is, we consider M_λ on a ray starting at the origin in the phase space $\{(k, \vec{l}_{1\perp})\}$.

Of course, we shall get a Laurent expansion for M_λ .

We illustrate this for the term $M_\lambda^{(a)}$.



$$M_{\lambda}^{(a)} = -e T(p_a - k, p_b, p_1', p_2') \big|_{\text{off shell}} \Delta[(p_a - k)^2] \Gamma_{\lambda}(p_a - k, p_a), \quad (4.6)$$

$$T(p_a - k, p_b, p_1', p_2') \big|_{\text{off shell}} = \mathcal{M}[(p_a - k, p_b) + (p_1 - l_1, p_2 - l_2), (p_b - p_2 + l_2)^2, (p_a - k)^2, m_{\pi}^2, m_{\pi}^2, m_{\pi}^2]. \quad (4.7)$$

From the generalised Ward identity we find for $\omega \rightarrow 0$

$$\Delta[(p_a - k)^2] \Gamma_{\lambda}(p_a - k, p_a) = \frac{(2p_a - k)_{\lambda}}{-2p_a \cdot k + k^2} + \mathcal{O}(\omega). \quad (4.8)$$

To get $M_{\lambda}^{(a)}$ to the orders ω^{-1} and ω^0 we need $\mathcal{M}$ to the orders ω^0 and ω .

This expansion of $M[]$ is straightforward, remembering that k, l_1, l_2 are all of order ω .

Treating in this way $M_\lambda^{(a)}$ and $M_\lambda^{(b)}$ and determining $M_\lambda^{(c)}$ to the order ω^0 from the gauge invariance condition (3.16) we get for the case of real photon emission, $k^2 = 0$, the following

$$\begin{aligned}
 M_\lambda(p_a, p_b, p_1', p_2', k) = & e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1'\lambda}}{p_1' \cdot k} \right] M^{(on)}(\nu, t) \\
 & - 2e \left[p_{a\lambda} \frac{p_{b\lambda} \cdot k}{p_a \cdot k} - p_{b\lambda} \right] \frac{\partial}{\partial \nu} M^{(on)}(\nu, t) \\
 & - 2e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}}{p_1 \cdot k} \right] \left[(p_a - p_1, k) - (p_a \cdot l_1) \right] \frac{\partial}{\partial t} M^{(on)}(\nu, t) + O(\omega).
 \end{aligned} \tag{4.9}$$

We can, for consistency, still expand

$$\frac{p_{1'\lambda}}{p_1' \cdot k} = \frac{(p_1 - l_1)_\lambda}{(p_1 - l_1, k)} = \frac{p_{1\lambda}}{p_1 \cdot k} + \frac{1}{(p_1 \cdot k)^2} \left[p_{1\lambda} (l_1 \cdot k) - l_{1\lambda} (p_1 \cdot k) \right] + O(\omega). \tag{4.10}$$

This gives us our final result for real photon emission

$$\begin{aligned}
 \mathcal{M}_\lambda(p_a, p_b, p'_1, p'_2, k) = & e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}}{p_1 \cdot k} \right] \mathcal{M}^{(on)}(\nu, t) \Big|_{\omega^{-1}} \\
 & - e \frac{1}{(p_1 \cdot k)^2} [p_{1\lambda}(\ell_1 \cdot k) - \ell_{1\lambda}(p_1 \cdot k)] \mathcal{M}^{(on)}(\nu, t) \\
 & - 2e \left[p_{a\lambda} \frac{p_b \cdot k}{p_a \cdot k} - p_{b\lambda} \right] \frac{\partial}{\partial \nu} \mathcal{M}^{(on)}(\nu, t) \\
 & - 2e \left[(p_a - p_1) \cdot k - (p_a \cdot \ell_1) \right] \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}}{p_1 \cdot k} \right] \frac{\partial}{\partial t} \mathcal{M}^{(on)}(\nu, t) \Big|_{\omega^0} \\
 & + \mathcal{O}(\omega).
 \end{aligned} \tag{4.11}$$

$$\nu = s - 2m_\pi^2, \quad t = (p_a - p_1)^2 = (p_b - p_2)^2.$$

With (4.11) we have given the Laurent expansion of $\mathcal{M}_\lambda$ to the orders ω^{-1} and ω^0 around the phase space point $(k=0, \vec{\ell}_{1\perp}=0)$ corresponding to $k=0, p'_1=p_1, p'_2=p_2$

The amplitude $M^{(0n)}(\nu, t)$ corresponds to the basic process

$$\pi^-(p_a) + \pi^0(p_b) \rightarrow \pi^-(p_1) + \pi^0(p_2).$$

The pole term $\propto \omega^{-1}$ in (4.11) is exactly Weinberg's soft-photon term. He writes

"Hence the effect of attaching one soft-photon line to an arbitrary diagram is simply to supply an extra factor,

$$\sum_n e_n \eta_n p_n^\mu / [p_n \cdot q - i\eta_n \varepsilon], \quad (4.12)$$

the sum running over all external lines in the original diagram."

Here $\eta_n = +1$ (-1) for an outgoing (incoming) charged particle.

In our work we have given the next to leading term, $\mathcal{O}(\omega^0)$, to Weinberg's pole term.

5 Soft photon theorem II

Now we want to discuss Low's version of soft-photon theorem.

Of course, as starting point he considers the diagrams (a), (b), (c) for M_λ . He also uses the generalised Ward identity which gave us (4.8) for $\Delta[(p_a - k)^2] \Gamma_\lambda(p_a - k, p_a)$. Considering only real photon emission we have then for $M_\lambda^{(a)}$ (see eq. (2.11) of Low)

$$M_\lambda^{(a)} = e M \left[(p_a - k, p_b) + p_1' \cdot p_2', (p_b - p_2')^2, m_a^2 = (p_a - k)^2, m_\pi^2, m_\pi^2, m_\pi^2 \right] \frac{p_a \cdot k}{p_a \cdot k} . \quad (5.1)$$

Now Low expands M with respect to k keeping p_1' and p_2' fixed.

That is, he expands with respect to k which is explicit in the chosen parametrisation. In this way we get :

$$\begin{aligned}
m_{\lambda}^{(a)}(p_a, p_b, p_1', p_2', k) = & e \frac{p_{a\lambda}}{(p_a \cdot k)} \left\{ m^{(on)}[p_a \cdot p_b + p_1' \cdot p_2', (p_b - p_2')^2] \right. \\
& - (p_b \cdot k) \frac{\partial}{\partial \nu} m^{(on)}[p_a \cdot p_b + p_1' \cdot p_2', (p_b - p_2')^2] \\
& \left. - 2(p_a \cdot k) \frac{\partial}{\partial m_a^2} \mathcal{M}[p_a \cdot p_b + p_1' \cdot p_2', m_a^2, m_{\pi}^2, m_{\pi}^2, m_{\pi}^2] \right|_{m_a^2 = m_{\pi}^2} \Big\} + \mathcal{O}(k).
\end{aligned} \tag{5.2}$$

Note an important point: whereas the expansion of the scalar function $\mathcal{M}[\]$ with respect to k keeping p_1' and p_2' fixed is completely legitimate and a usual expansion, this is not the case for $m_{\lambda}^{(a)}$. We must remember the energy - momentum conservation:

$$p_a + p_b = p_1' + p_2' + k. \tag{5.3}$$

Keeping p_1' and p_2' fixed, also k is fixed and (5.2) makes sense only for this one value of k .

Now we can treat $M_\lambda^{(b)}$ in a similar way and then determine $M_\lambda^{(c)}$ approximately from the gauge-invariance condition (3.16). The result is Low's formula (eq. (2.16) of Low):

$$M_\lambda(p_a, p_b, p'_1, p'_2, k) = e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p'_{1\lambda}}{p'_1 \cdot k} \right] M^{(on)}(\nu_L, t_2) \\ - e \left[p_{a\lambda} \frac{p_b \cdot k}{p_a \cdot k} + p'_{1\lambda} \frac{p'_2 \cdot k}{p'_1 \cdot k} - p_{b\lambda} - p'_{2\lambda} \right] \frac{\partial}{\partial \nu_L} M^{(on)}(\nu_L, t_2) + O(k),$$

(5.4)

where

$$\nu_L = p_a \cdot p_b + p'_1 \cdot p'_2 = s - 2m_\pi^2 - (p_a + p_b, k), \\ t_2 = (p_b - p'_2)^2 = (p_a - p'_1 - k)^2. \quad (5.5)$$

We emphasise again that (5.4) is not an expansion of M_λ around some phase-space point. The r.h.s of (5.4) gives an approximate expression for M_λ at a given phase-space point p'_1, p'_2, k .

We can use (5.4) only for one value of $k = p_a + p_b - p_1' - p_2'$.

If we use (5.4) at a different value of k we violate energy-momentum conservation.

Furthermore, the leading approximation in (5.4) does not give what is frequently called Low's theorem. We see this best by considering the reactions

$$\pi^-(p_a) + \pi^+(p_b) \longrightarrow \pi^-(p_1') + \pi^+(p_2') + \gamma(k, \varepsilon),$$

$$\pi^-(p_a) + \pi^+(p_b) \longrightarrow \pi^-(p_1) + \pi^+(p_2).$$

(5.6)

According to Low we get

$$\begin{aligned} \mathcal{M}_\lambda(p_a, p_b, p_1', p_2', k) = & e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}'}{p_1' \cdot k} \right] \mathcal{M}^{(on)}(\nu_L, t_2) \\ & + e \left[-\frac{p_{b\lambda}}{p_b \cdot k} + \frac{p_{2\lambda}'}{p_2' \cdot k} \right] \mathcal{M}^{(on)}(\nu_L, t_1) + \mathcal{O}(\omega^0), \end{aligned}$$

$$\nu_L = s - 2m_\pi^2 - (p_a + p_b, k), \quad t_1 = (p_a - p_1')^2, \quad t_2 = (p_b - p_2')^2, \quad (5.7)$$

According to Weinberg we have, see (4.12),

$$\begin{aligned} \mathcal{M}_\lambda(p_a, p_b, p_1', p_2', k) = & e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}}{p_1 \cdot k} - \frac{p_{b\lambda}}{p_b \cdot k} + \frac{p_{2\lambda}}{p_2 \cdot k} \right] \mathcal{M}^{(on)}(\nu, t) + \mathcal{O}(\omega^0) \\ \nu = s - 2m_\pi^2, \quad t = & (p_a - p_1)^2 = (p_b - p_2)^2. \end{aligned} \quad (5.8)$$

Low's formula (5.7) gives an approximate expression for $\mathcal{M}_\lambda$ at one given phase-space point. Weinberg's formula (which is frequently but incorrectly called Low's theorem) gives the pole term of the Laurent expansion of $\mathcal{M}_\lambda$ around the point $p_i' = p_i, k = 0$.

Let us go back to $\pi^- + \pi^0 \rightarrow \pi^- + \pi^0 + \gamma$. In (5.4) we have Low's formula which gives us an approximate expression for M_λ at a given phase-space point. We can construct, as we did in Sec. 4, the corresponding expansion of this approximate expression around the phase-space point $(k=0, \hat{p}_1)$. Inserting in (5.4) $p'_1 = p_1 - l_1$, $p'_2 = p_2 - l_2$ from (4.2) we get

$$\begin{aligned}
 M^{(0n)}(\nu_L, t_2) &= M^{(0n)}[\nu - (p_a + p_b, k), t - 2[(p_a - p_1, k) - p_a \cdot l_1]] + O(\omega^2) \\
 &= M^{(0n)}(\nu, t) - (p_a + p_b, k) \frac{\partial}{\partial \nu} M^{(0n)}(\nu, t) - 2[(p_a - p_1, k) - p_a \cdot l_1] \frac{\partial}{\partial t} M^{(0n)}(\nu, t) \\
 &\quad + O(\omega^2),
 \end{aligned}
 \tag{5.9}$$

$$\begin{aligned}
M_\lambda(p_a, p_b, p'_1, p'_2, k) = & e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p'_{1\lambda}}{p'_1 \cdot k} \right] m^{(on)}(\nu, t) \\
& - 2e \left[p_{a\lambda} \frac{p_b \cdot k}{p_a \cdot k} - p_{b\lambda} \right] \frac{\partial}{\partial \nu} m^{(on)}(\nu, t) \\
& - 2e \left[\frac{p_{a\lambda}}{p_a \cdot k} - \frac{p_{1\lambda}}{p_1 \cdot k} \right] \left[(p_a - p_1, k) - (p_a \cdot l_1) \right] \frac{\partial}{\partial t} m^{(on)}(\nu, t) + O(\omega).
\end{aligned}
\tag{5.10}$$

This is identical to our result (4.9) and from there we can go on to (4.11) using (4.10).

In this way we have given the relation between Low's formula and the Laurent series (4.11) where the pole term $\propto \omega^{-1}$ is given by Weinberg's soft-photon theorem and the next to leading term $\propto \omega^0$ by our calculation.