



AI and finding signatures of phase transition

Olga Soloveva
GSI & Goethe University
Poster #6



Interaction rate $> 10^7$ Hz


100 GB/s throughput ceiling

Only a **small fraction** of collisions can be stored -
Selection must happen online



The goal: Establish whether an ANN trigger can select QGP-sensitive events from reconstructed final states – and whether its decision survives changing the event generator and passing through a real detector chain.

Multi-output hybrid NN


28 x 20 x 20 x 20 input tensor



28	particle species	channels; seen $\geq$ once per 1000 events
20	$ p $ momentum	log-binned, dense at low momentum
20	θ polar angle	equal bins
20	ϕ azimuthal angle	equal bins



Cross-Model Transfer: The network works on physics it has never seen.



Result: The network learned generic QGP features, not the details of a single simulation.

Interpretable ANN: Rediscovering Strangeness Enhancement




The Insight

SHAP (Shapley Additive Explanations) analysis reveals that the network's decision is heavily driven by the presence of strange particles (K , Λ , Ξ , Ω) and antibaryons.

The Validation



No strangeness observables or physics-informed loss functions were ever imposed during training. The CNN learned the correct features associated with QGP production entirely on its own.

From Ideal Simulations to the Beam




Takeaway: Accuracy drops gracefully. The model is highly robust against incomplete detector information, tracking uncertainties, and partial topology loss.

Signatures of phase transition



! QGP fraction is small at low $\sqrt{s_{NN}}$

- Selective Antibaryon Suppression
- Softened Transverse Momentum (p_T)



A faster, accurate QGP trigger ready for the CBM Experiment

'Speed'

0.14 ms
per event.

Runs seamlessly
(~7,300 events/sec) on a
single laptop GPU

'The Core Proofs'



Model Independent
(Transfers from PHSD to UrQMD)



Physically Interpretable
(Driven by strangeness/antibaryons)



Detector Robust
(Survives full CBM reconstruction chain)

'Call to Action'



Scan for the full poster
and physics references.