



Contribution ID: 45

Type: Oral

Dipole toroidal resonance: vortical properties, deformation impact, relation to pygmy mode

Thursday, 3 September 2015 14:30 (15 minutes)

Properties of the isoscalar dipole toroidal resonance (TR) in spherical and deformed nuclei are discussed using our recent results obtained within the self-consistent random phase approximation with Skyrme forces [1-4]. The TR strength functions, transition densities and velocity fields are inspected. The resonance is compared to other exotic isoscalar dipole modes (compression and pygmy) [1,2]. The main attention is paid to: i) possibility to use TR as a measure of the nuclear vorticity [3], ii) anomalous TR deformation splitting [4], iii) relation of the TR and pygmy dipole resonance [2], iv) perspectives of experimental observation of the TR.

[1] J. Kvasil, V.O. Nesterenko, W. Kleinig, P.-G. Reinhard, and P. Vesely, Phys. Rev. C 84, 034303 (2011).

[2] A. Repko, P.-G. Reinhard, V.O. Nesterenko, and J. Kvasil, Phys. Rev. C 87, 024305 (2013).

[3] P.-G. Reinhard, V.O. Nesterenko, A. Repko, and J. Kvasil, Phys. Rev. C 89, 024321 (2014).

[4] J. Kvasil, V.O. Nesterenko, W. Kleinig, and P.-G. Reinhard, Phys. Scripta, v.89, n.5, 054023 (2014).

Primary author: NESTERENKO, Valentin (Joint Institute for Nuclear Research, Dubna)Presenter: NESTERENKO, Valentin (Joint Institute for Nuclear Research, Dubna)Session Classification: Nuclear Structure, Spectroscopy, and Dynamics IV