

# Hyperfine spectroscopy setup for antihydrogen and first results with a hydrogen beam

Thursday, September 18, 2014 5:00 PM (30 minutes)

The ASACUSA collaboration aims to measure the ground state hyperfine splitting of the antihydrogen atom, since this is a system where the CPT symmetry can be investigated with extremely high sensitivity. The principal idea is described in [1,2].

During the CERN LS1 shut down, antiprotons were not available. Therefore, a source of cold, polarized, and modulated atomic hydrogen has been constructed to enable comprehensive testing of the Rabi-like experimental setup consisting of a microwave spin flip cavity and superconducting sextupole magnet [3].

After shortly discussing the main components of the atomic hydrogen source and detector as well as the spectroscopy beamline I will present the latest experimental data, which allowed for a characterization of the focusing effect of the superconducting sextupole magnet and the resonance line shape of the spin flip cavity. Furthermore, a confirmation of the proposed measurement principle for the hyperfine splitting of antihydrogen was achieved by the determination of the ground state hyperfine splitting of atomic hydrogen with a precision on the 10 ppb level.

## References

[1] E. Widmann et al., *Hyperfine Interactions*, 215, 1-8 (2013)

[2] N. Kuroda et al., *Nature Communications*, 2089, 5 (2014)

[3] C. Malbrunot et al., *Hyperfine Interactions*, (2014) DOI: 10.1007/s10751-014-1013-z

**Author:** DIERMAIER, Martin (Austrian Academy of Sciences (AT))

**Co-authors:** Dr MALBRUNOT, Chloé (SMI (Austrian Academy of Science)); KLAUSHOFER, Christoph (Stefan-Meyer-Institute); Mr SAUERZOPF, Clemens (SMI Wien); Prof. WIDMANN, Eberhard (Stefan Meyer Institute); ZMESKAL, Johann (SMI); SIMON, Martin (Stefan-Meyer-Institute); OSWALD, Massiczek (Stefan-Meyer-Institute); WOLF, Michael (Stockholms Universitet); CARADONNA, Peter (unknown)

**Presenter:** DIERMAIER, Martin (Austrian Academy of Sciences (AT))