

Search for Long-Lived States of $\pi^+\pi^-$ atom

Friday, 19 September 2014 12:00 (30 minutes)

The observation of long-lived (metastable) states of $\pi^+\pi^-$ atoms ($A_2\pi$) opens the possibility to measure the energy difference between ns and np states and to determine the value of the combination $2a_0 + a_2$ of S-wave $\pi^+\pi^-$

scattering lengths with isotope spins 0,2 in a model-independent way. This result, together with the $A_2\pi$ lifetime

measurement that provides the value $|a_0 - a_2|$, allows to get a_0 and a_2 separately using only $\pi^+\pi^-$ atoms data.

In this experiment the proton beam with momentum 24 GeV/c interacts with a Be target with thickness 100mm and

generates $A_2\pi$ in short-lived ns states. Passing through the target a fraction of $A_2\pi$ interacts with Be-atoms and

get exited into long-lived $2p, 3p, 4p\dots$ states. From the Be target more than 6% of $A_2\pi$ come out to the vacuum in

the long-lived states. For the short lived $A_2\pi$, with Lorenz factor 20, the decay lengths of $2S, 3S$ and $4S$ are in the interval between 0.017 and 1.1mm, while the metastable atoms in the states $2P, 3P$ and $4P$ have the decay lengths

between 5.7cm and 44cm.

After the Be target at a distance of 100mm, it was installed a Pt foil in which only long-lived atoms break up, generating $\pi^+\pi^-$ pairs with small relative momentum Q in their c.m.s. In order to suppress the background from the

$\pi^+\pi^-$ generated in the Be target, a magnet with $BL=0.023$ Tm was installed between Pt foil and the target. At the

exit of this magnet the pairs produced on the Be target have their Q_y component increased of 12.7 MeV/c, while the

pairs generated on Pt foil, have their Q_y component increased only of 2.3MeV/c by the fringing magnetic field. In

this report we present the results of the analysis which select $\pi^+\pi^-$ pairs with small transverse component $Q_T < 1.5$

MeV/c. The distribution in the longitudinal component Q_L of these pairs shows a peak around $Q_L=0$ MeV/c. The

statistical significance of this peak is 5s and it could be explained by the long-lived $\pi^+\pi^-$ atoms breaking in the Pt foil.

Primary author: Prof. NEMENOV, Leonid (JINR Dubna)

Presenter: Prof. NEMENOV, Leonid (JINR Dubna)