



Contribution ID: 58

Type: **not specified**

## Reconstruction of $\pi^0$ and $\eta$ mesons via conversion method in Au+Au at 1.23A GeV with HADES

*Saturday, 27 September 2014 11:30 (30 minutes)*

Lepton pairs emerging from decays of virtual photons represent promising probes of matter under extreme conditions. In the energy domain of 1 - 2 GeV per nucleon, the HADES experiment at GSI Helmholtzzentrum fuer Schwerionenforschung in Darmstadt studies di-electrons and strangeness production in various reactions, i.e. collisions of pions, protons, deuterons and heavy-ions with nuclei. An accurate determination of the medium radiation depends on a precise knowledge of the underlying hadronic cocktail composed of various sources contributing to the net spectra. Therefore, a measurement of the neutral meson yields together with the dileptons is crucial. In this contribution, the capability of HADES to detect  $e^-e^+$  pairs from conversions of real photons will be demonstrated. We will present results from a two-photon analysis of Au+Au collisions

at 1.23 GeV/u providing information on neutral  $\pi^0$  and  $\eta$  mesons. Supported by BMBF (06FY9100I and 06FY7114), HIC for FAIR, EMMI, GSI, HGS-Hire and H-QM

**Primary author:** Mrs BEHNKE, Claudia (Institut fuer Kernphysik, Goethe-Universitaet, Frankfurt, Germany)

**Presenter:** Mrs BEHNKE, Claudia (Institut fuer Kernphysik, Goethe-Universitaet, Frankfurt, Germany)

**Session Classification:** Talks