



Contribution ID: 51

Type: **not specified**

Charged Kaon- and ϕ -reconstruction in Au+Au-collisions at 1.23 AGeV

Tuesday, 23 September 2014 17:30 (30 minutes)

In Au+Au-collisions at 1.23 AGeV, the complete strangeness production is below its nucleon-nucleon threshold. In baryon dominated matter K^+ and K^- exhibit different properties, because K^- can be resonantly absorbed by nucleons. Although strangeness exchange reactions have been proposed to be the dominant channel for K^- production below threshold, the production yield could also be explained in Ar+KCl-reactions at 1.76 AGeV based on a statistical hadronization model fit to the measured particle yields. To take care for strangeness conservation, strangeness is calculated canonically within R_c , and therefore the ratio of ϕ/K^- is predicted to rise with decreasing beam energies and hence the feed-down of ϕ -mesons to Kaons becomes more important.

In April and May 2012, 7.3 billion Au(1.23 GeV per nucleon)+Au collisions have been recorded by the HADES detector. In this contribution, we present results of the charged kaon and ϕ -meson production and test the above discussed prediction.

Primary author: Mrs SCHULDES, Heidi (IKF, Goethe-Universität)

Presenter: Mrs SCHULDES, Heidi (IKF, Goethe-Universität)

Session Classification: Talks