

Recent electron-spectroscopy experiments at the ESR

Monday, September 19, 2016 11:50 AM (20 minutes)

The energy distribution of electrons emitted in heavy-ion atom collision is a characteristic observable for understanding the underlying charge-transfer processes. Previously, we studied the collision system $U88+ + N_2$ at a projectile energy of 90 MeV/u at the ESR using the magnet forward-angle electron spectrometer [1-3]. For electrons released from the projectile ions in collisions with neutral target atoms, the shape of the electron-loss-to-continuum cusp is not a function of the target atomic number according to first-order perturbation theory, which was applied in ref. [2]. However, for the cusp shape of $U28+$ ions colliding with different atomic targets, which is relevant for understanding the lifetimes of beams in the heavy-ion accelerators of FAIR, we observed a strong asymmetry of the electron cusp shape that varies significantly with the atomic number of the target [4]. These spectra cannot be explained by current theories. In a recent experiment we therefore compared the cusp-electron spectra of $U89+$ projectiles colliding with neutral N_2 and Xe targets at 76 MeV/u. Amongst others, the experimental results provide tests to theory describing two-center effects in the double-differential cross-sections of projectile ionization of heavy ions [5].

[1] P.-M. Hillenbrand et al., Phys. Rev. A 90, 022707 (2014)

[2] P.-M. Hillenbrand et al., Phys. Rev. A 90, 042713 (2014)

[3] P.-M. Hillenbrand et al., Phys. Rev. A 91, 022705 (2015)

[4] P.-M. Hillenbrand et al., Phys. Rev. A 93, 042709 (2016)

[5] A. B. Voitkiv et al., Phys. Rev. A 76, 022709 (2007)

Primary author: HILLENBRAND, Pierre-Michel (GSI, Darmstadt)

Co-authors: HAGMANN, Siegbert (Goethe-Universität Frankfurt(UFfm-IKP)/GSI-Darmstadt); STÖHLKER, Thomas (GSI, Darmstadt); Dr LITVINOV, Yury (GSI, Darmstadt)

Presenter: HILLENBRAND, Pierre-Michel (GSI, Darmstadt)

Session Classification: SPARC Experiments