

# The new isotopes $^{240}\text{Es}$ and $^{236}\text{Bk}$

Joonas Konki

Department of Physics  
University of Jyväskylä

TASCA 16

The 15th Workshop on Recoil Separator  
for Superheavy Element Chemistry

26.8.2016

GSI, Darmstadt, Germany



# Outline

- 1 Short introduction
  - Motivation
  - Electron-capture delayed fission (ECDF)
- 2 Experimental details
  - R48 experiment (December 2014)
  - RITU and GREAT
- 3 Data analysis and results
  - ER- $\alpha$  correlations
  - ER- $\alpha$ - $\alpha$  correlations
  - ER- $\alpha$ , ER-Fission, ER- $\alpha$ -Fission time distributions
- 4 Conclusions
  - Decay properties of the new isotopes
  - ECDF probability systematics in Es and Bk isotopes



# Motivation

## Why study neutron-deficient heavy nuclei?

- Ideal cases to study the competition of the attractive nuclear and the repulsive Coulomb forces that determine the stability of nuclei
- Provide information on the nuclear mass surface close to the proton dripline
- Help understand the processes and stability of superheavy elements (SHE)
- Study low-energy fission properties of excited nuclei via ECDF



# Electron-capture delayed fission (ECDF)




Fig. from A. Andreyev et al. Rev. Mod. Phys. 85 (2013)

- Parent nucleus EC/ $\beta^+$  decays
- An excited state in the daughter nucleus is populated
- If  $Q_{\text{EC}} \sim B_f$  or greater, then fission may compete with other decay modes of the excited states

# R48 experiment (December 2014)

## Experiment details:

Spokespersons:

J. Khuyagbaatar (GSI),  
J. Uusitalo (JYFL)

Reaction:



${}^{209}\text{BiO}_2$  target  $500 \mu\text{g}/\text{cm}^2$

6 days of beam on target

$E_{\text{beam}} = 178, 174 \text{ MeV}$

C degrader foils:  $200, 400 \mu\text{g}/\text{cm}^2$

$\rightarrow E^* = 39, 36, 35, 34 \text{ MeV}$

Typical  $I_{\text{beam}} \approx 100\text{-}200 \text{ pA}$

Total beam dose  $\approx 3.7 \cdot 10^{17}$

RITU


DSSD-Y (80 strips)

DSSD-X (120 strips)


3 Clovers + Planar Ge + Pins at FP

2 BGO PMTs at target position

- ${}^{209}\text{Bi}({}^{34}\text{S}, 3n){}^{240}\text{Es}$  reaction to study the decay properties of the new isotopes  ${}^{240}\text{Es}$  and its  $\alpha$ -decay daughter  ${}^{236}\text{Bk}$
- A relatively high ECDF probability ( $P_{\text{ECDF}}$ ) expected for  ${}^{240}\text{Es}$  from systematics




# RITU and GREAT



RITU gas-filled recoil separator  
GREAT focal plane spectrometer


# Raw $\alpha$ -like events



- (a) Raw  $\alpha$ -like events, MWPC and Planar vetoed




# ER- $\alpha$ correlations



- Fusion-evaporation residue (ER) correlated  $\alpha$ -like events, 200 s searching time
- Short-lived  $\alpha$ -decays from slow transfer reaction products still visible (Region 1)




# ER- $\alpha$ correlations



- (a) Raw  $\alpha$ -like events, MWPC and Planar vetoed
- (b) ER- $\alpha$ ,  $\Delta t \leq 200$  s




# ER- $\alpha$ correlations



- (a) Raw  $\alpha$ -like events, MWPC and Planar vetoed
- (b) ER- $\alpha$ ,  $\Delta t \leq 200$  s
- (c) ER- $\alpha$ ,  $\Delta t \leq 30$  s

# ER- $\alpha$ - $\alpha$ correlations




- ER- $\alpha_1$ - $\alpha_2$ ,  
 $\Delta t_{\text{ER}-\alpha_1} \leq 200$  s,  
 $\Delta t_{\alpha_1-\alpha_2} \leq 1200$  s
- Can see ER correlated  $\alpha$  chains of interest:  
 ER- $\alpha_1$ (7.57 MeV)- $\alpha_2$ ( $^{236}\text{Cm}$ )  
 ER- $\alpha_1$ (8.19 MeV)- $\alpha_2$ ( $^{236}\text{Cm}$ )  
 ER- $\alpha_1$ (8.09 MeV)- $\alpha_2$ ( $^{236}\text{Cm}$ )  
 ER- $\alpha_1$ (8.02 MeV)- $\alpha_2$ ( $^{236}\text{Cm}$ )  
 ER- $\alpha_1$ (7.97 MeV)- $\alpha_2$ ( $^{236}\text{Cm}$ )

→ attributed to  $^{240}\text{Cf}$  and the new isotope  $^{240}\text{Es}$

—  
 $^{236}\text{Cm}$ :  $E_\alpha=6.954(20)$  MeV,  
 $T_{1/2}=410(50)$  s


# ER- $\alpha$ - $\alpha$ correlations



- (a) Raw  $\alpha$ -like events, MWPC and Planar vetoed
- (b) ER- $\alpha$ ,  $\Delta t \leq 200$  s
- (c) ER- $\alpha$ ,  $\Delta t \leq 30$  s
- (d) ER- $\alpha_1$ - $\alpha_2$ ,  $\Delta t_{\text{ER-}\alpha_1} \leq 200$  s,  $\Delta t_{\alpha_1-\alpha_2} \leq 1200$  s and  $E_{\alpha_2}$  in range 6.85-7.05 MeV


—  
 $^{236}\text{Cm}$ :  $E_{\alpha} = 6.954(20)$  MeV,  
 $T_{1/2} = 410(50)$  s

# ER- $\alpha$ , ER-Fission, ER- $\alpha$ -Fission time distributions



- 27 ER- $\alpha$ (8.19 MeV) events with  $T_{1/2} \sim 6$  s  
→ attributed to  $\alpha$  decay of  $^{240}\text{Es}$
- 60 ER- $\alpha$ (8.09 MeV) events with  $T_{1/2} \sim 6$  s  
→ attributed to  $\alpha$  decay of  $^{240}\text{Es}$

# ER- $\alpha$ , ER-Fission, ER- $\alpha$ -Fission time distributions



- 27 ER- $\alpha$ (8.19 MeV) events with  $T_{1/2} \sim 6$  s  
→ attributed to  $\alpha$  decay of  $^{240}\text{Es}$

- 60 ER- $\alpha$ (8.09 MeV) events with  $T_{1/2} \sim 6$  s  
→ attributed to  $\alpha$  decay of  $^{240}\text{Es}$

- 15 ER-Fission events with  $T_{1/2} \sim 5$  s  
→ attributed to ECDF of  $^{240}\text{Es}$

- 8 ER- $\alpha$ (8.09 MeV)-Fission events (4 of them are escape  $\alpha$ s) with  $T_{1/2} = 22^{+13}_{-6}$  s  
→ attributed to ECDF of  $^{236}\text{Bk}$


# Decay properties of the new isotopes ${}_{99}^{240}\text{Es}_{141}$ and ${}_{97}^{236}\text{Bk}_{139}$

## Decay properties of ${}^{240}\text{Es}$

- ER- $\alpha$ , ER- $\alpha$ - $\alpha$ , ER-fission correlations analysed
- $E_{\alpha} = 8.19(3)$  MeV,  $8.09(3)$  MeV,  $(8.02(3)$  MeV,  $7.97(3)$  MeV)  
 $T_{1/2} = 6(2)$  s
- $b_{\alpha} = 0.7(1)$   
 $b_{\text{EC}} = 0.3(1)$
- $P_{\text{ECDF}} = 0.16(6)$

## Decay properties of ${}^{236}\text{Bk}$


- ER- $\alpha$ ( ${}^{240}\text{Es}$ )-fission, 8 events
- $T_{1/2} = 22_{-6}^{+13}$  s
- No  $\alpha$ -decay branch seen
- $P_{\text{ECDF}} = 0.04(2)$



The proposed decay scheme of  ${}^{240}\text{Es}$  and  ${}^{236}\text{Bk}$

J. Konki et al. to be published.  
(Data for  ${}^{240}\text{Cf}$  and  ${}^{236}\text{Cm}$  are from literature.)

# ECDF probability systematics in Es and Bk isotopes



- Probability increases exponentially as a function of the energy difference  $Q_{\text{EC}} - B_{\text{sf}}$
- Similar behaviour observed for other neutron-deficient nuclei (Am, Np, Tl)
- Probability expected to approach saturation ( $P_{\text{ECDF}} \rightarrow 1$ ) in the next lighter odd-odd Es and Bk isotopes

$Q_{\beta}$  and  $B_{\text{sf}}$  from P. Möller et al. 1997, 2009

Other  $P_{\text{ECDF}}$  from A. Andreyev et al. Rev. Mod. Phys. 85 (2013)



# Thank you!

J. Khuyagbaatar, J. Uusitalo, P.T. Greenlees, K. Auranen, H. Badran, M. Block, R. Briselet, D.M. Cox, M. Dasgupta, A. Di Nitto, Ch. E. Düllman, T. Grahn, K. Hauschild, A. Herzán, R.-D. Herzberg, F. P. Heßberger, M. Hinde, R. Julin, S. Juutinen, E. Jäger, B. Kindler, J. Krier, M. Leino, B. Lommel, A. Lopez-Martens, D. Luong, M. Mallaburn, K. Nishio, J. Pakarinen, P. Papadakis, J. Partanen, P. Peura, P. Rahkila, K. Rezynkina, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, S. Stolze, B. Sulignano, Ch. Theisen, A. Ward, A. Yakushev and V. Yakusheva



JOHANNES GUTENBERG  
UNIVERSITÄT MAINZ



Australian  
National  
University



Science & Technology  
Facilities Council

