

COMPASS

Status and Recent Results

Jan Friedrich
Technische Universität München
on behalf of the COMPASS collaboration




Bad Honnef
2.12.2016



bmb+f - Förderschwerpunkt
COMPASS
Großgeräte der physikalischen
Grundlagenforschung

Collaboration: 249 members

22 institutes: Prague, Saclay, Bonn, Freiburg, Mainz, München, Calcutta, Tel Aviv, Turin, Trieste, Yamagata, Warshaw, Lisbon, Dubna, Moscow, Tomsk, CERN, Taiwan, Illinois



COMPASS II program

- 2012 Primakoff $\{\pi/K\}\gamma$
- 2014/15 Drell-Yan
- 2016/17 DVCS / SIDIS





The Physics Case: Exploring QCD

with a multi-purpose setup




- Deep-inelastic **muon scattering**
with longitudinal and transverse target polarisation
Mainz, Bonn, Freiburg
- High-energy **hadron scattering**
at low and intermediate excitation energies **München, Bonn**
- Primakoff reactions: π^\pm polarisability, chiral dynamics **München**
- (First) Polarised Drell-Yan **Bonn**
- Exclusive muon-induced processes **Freiburg, Bonn, Mainz**
 - ▶ deeply-virtual Compton scattering
 - ▶ hard exclusive meson production

Results on g_1^p and g_1^d


longitudinal spin-dependent structure functions

A_1



proton



deuteron




g_1




- first moment Γ_1 : verification of Bjorken sum rule (94% in cov. range)
- NLO QCD fit \rightarrow polarised parton distributions, $0.26 < \Delta\Sigma|_{Q^2=3} < 0.36$

Meson spectroscopy: 3π final state




arXiv:1509.00992



⇒




- 46 million exclusive 3π events
- partial-wave fit with 88 waves in narrow 3π -mass slices
- mass dependence fitted for 14 waves $\sim 75\,000$ data points (including interference terms)

Example: $0^{-+}0^+ f_0(980)\pi S$ wave

$$m_{\pi(1800)} = 1802.6^{+8}_{-3.5} \text{ MeV}/c^2 ; \Gamma_{\pi(1800)} = 218^{+11}_{-6} \text{ MeV}/c^2$$

$$m_{\pi(1800)}^{\text{PDG}} = 1812 \pm 12 \text{ MeV}/c^2 ; \Gamma_{\pi(1800)}^{\text{PDG}} = 208 \pm 12 \text{ MeV}/c^2$$

- $\pi(1800)$ previously observed to decay in $f_0(980)\pi$ and $f_0(1500)\pi$
→ “fixed f_0 isobars” assumed in the fit
- new analysis method: *this assumption can be tested!*

De'isobar'ing of $\pi(1800)$ 


- Coupling of $\pi(1800)$ to $f_0(980)\pi$ and $f_0(1500)\pi$ decay modes

→ 2 π -isobar parameters can be **extracted** from the 3 π final states (*ongoing*)

Results on 3π resonances


new: parameters of 11 resonances

- main known resonances reproduced
- all resonance parameters determined in one single fit
- new signal: $a_1(1420)$
- three π_2 states needed
- (broad) exotic 1^{-+} signal




Polarised Drell-Yan 2015




Sivers asymmetry - news about sign question expected soon



SIDIS




single-polarised DY



$$\vec{\mu}^\pm p \rightarrow \mu^\pm p \gamma$$



$$S_{CS,U} \equiv d\sigma(\mu^{+\leftarrow}) + d\sigma(\mu^{-\rightarrow}) \propto d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + K s_1^{\text{Int}} \sin\phi$$

$$\rightarrow d\sigma^{DVCS}/d|t| \sim \exp(-B|t|)$$





$$\langle r_\perp^2(x_B) \rangle \approx 2B(x_B)$$

$r_\perp \rightarrow$ distance between struck and spectator partons




Transverse Shape of the Proton

Data taken in 2012



November 2016:
long run part 1 just finished
new DAQ München





Conclusions

diverse, exciting, long-range physics program



- Inclusive and Semi-inclusive DIS

- Meson Spectroscopy

- ▶ 2008 and 2009: 3π resonances studies with unprecedented precision
- ▶ > 2018 dedicated future Kaon programme with RF-separated beam (★)

- Chiral Dynamics

- ▶ 2012 Primakoff run → high-precision α_π, β_π

- Polarised Drell-Yan

- ▶ 2015: successful beam time
- ▶ 2018 proposed followup

- Generalized Parton Distributions

- ▶ 2016 and 2017 → GPD H *ongoing*
- ▶ > 2018 polarised target (★) → GPD E

(★) March 2016: COMPASS “BEYOND 2020” Workshop

<https://indico.cern.ch/event/502879/>

March 2017: IWHSS workshop in Cortona (Tuscany)

Citation: C. Pestrignani et al. (Particle Data Group), Chin. Phys. C, 40, 100001 (2016)



$$I^G(J^P) = 1^-(0^-)$$

π ELECTRIC POLARIZABILITY α_π

SEE HOLSTEIN 14 for a general review on hadron polarizability.

VALUE (10^{-4} fm 3)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.0 \pm 0.6 \pm 0.7$	63k	1 ADOLPH	15A SPEC	$\pi^-\gamma \rightarrow \pi^-\gamma$ Compton scatt.

¹Value is derived assuming $\alpha_\pi = -\beta_\pi$.



$$I^G(J^P C) = 1^-(1^{++})$$

OMITTED FROM SUMMARY TABLE

a₁(1420) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1414^{+15}_{-13}	1 ADOLPH	15C COMP	190 $\pi^-\rho \rightarrow \pi^-\pi^+\pi^-p$

¹Using the isobar model and partial-wave analysis with 88 waves.

some of the new COMPASS entries
in the RPP2016 edition



Thank you for your attention!





Backup



Is Peak in $1^{++} 0^+ f_0(980)\pi P$ Wave a Model Artifact?

Novel analysis method

(inspired by E791 analysis, PRD **73** (2006) 032204)

- Replace $J^{PC} = 0^{++}$ isobar parametrizations by piece-wise constant amplitudes in $m_{\pi^+\pi^-}$ bins
- Extract $m_{3\pi}$ dependence of 0^{++} isobar amplitude from data
 - Drastic reduction of model bias
 - *Caveat:* significant increase in number of fit parameters
- Result: the $a_1(1420)$ signal is indep. on the $f_0(980)$ description

