

# Thermal dileptons from coarse-grained approach as fireball probes




TECHNISCHE  
UNIVERSITÄT  
DARMSTADT

CBM-STAR joint Workshop, Darmstadt

Florian Seck – TU Darmstadt




in collaboration with T. Galatyuk, R. Rapp & J. Stroth



# Electromagnetic probes in heavy-ion collisions

## Experiments across the QCD phase diagram



▲ NA60 ( $\mu+\mu^-$ ) : H.J.Specht: AIP Conf. Proc. 1322 (2010)

### ► Search for

#### ► phase boundary(ies)

→ fluctuations of conserved quantum numbers

→ flavor production (multi-strange, charm)

#### ► change in microscopic degrees of freedom

#### ► restoration of chiral symmetry

#### ► emitting source temperature


→ electromagnetic probes leave collision zone  
undistorted

→ real  $\gamma$  characterized by transverse momentum


→ dileptons carry extra information: invariant mass

# Electromagnetic probes in heavy-ion collisions

## CBM cocktail – invariant mass of dielectrons



dilepton spectra reflect the  
whole history of a collision



➔ necessary ingredients:



- ▶ realistic emission rates
- ▶ accurate description of fireball evolution

# Electromagnetic probes in heavy-ion collisions

## Insights from theory

- ▶ integrated yield of thermal radiation in the mass range  $0.3\text{-}0.7\text{ GeV}/c^2$  is sensitive to the lifetime of the fireball  
R. Rapp, H. van Hees: Phys. Lett. B **753** (2016) 586
- ▶ dilepton yield determined by interplay between temperature and fireball volume
- ▶ slope of dileptons in the intermediate-mass range constitutes a blue-shift free fireball thermometer

## ▶ What happens at low energies?




# Realistic dilepton emission rates

## 8-differential thermal production rate

$$\frac{dN_{ll}}{d^4x d^4q} = \frac{\alpha_{\text{EM}}^2}{\pi^3 M^2} f^B(q \cdot u; T) \text{Im}\Pi_{\text{EM}}(M, q; \mu_B, T)$$

$$R = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} \propto \frac{\text{Im}\Pi_{\text{EM}}^{\text{vac}}}{M^2}$$




$$\text{Im}\Pi_{\text{EM}}^{\text{vac}}(M) = \begin{cases} \sum_{v=\rho,\omega,\phi} \left(\frac{m_v^2}{g_v}\right)^2 \text{Im}D_v^{\text{vac}}(M), & M < M_{\text{dual}}^{\text{vac}} \simeq 1.5 \text{ GeV}/c^2 \\ -\frac{M^2}{12\pi} \left(1 + \frac{\alpha_s(M)}{\pi} + \dots\right) N_c \sum_{q=u,d,s} (e_q)^2, & M > M_{\text{dual}}^{\text{vac}} \end{cases}$$


# Realistic dilepton emission rates

## The $\rho$ meson in nuclear matter

R. Rapp, J. Wambach: Eur. Phys. J. A **6** (1999) 415




vacuum



medium

$$D_\rho(M, q; \mu_B, T) = [M^2 - m_\rho^2 - \Sigma_{\rho\pi\pi} - \Sigma_{\rho B} - \Sigma_{\rho M}]^{-1}$$



The  $\rho$  spectral function strongly broadens in the medium as the  $\rho$  meson couples to baryons !

additional contributions to the  $\rho$  meson self-energy in the medium

# Realistic dilepton emission rates

## Hadronic matter

- ▶ parameterization of Rapp-Wambach in-medium  $\rho$  spectral function

R. Rapp, J. Wambach: Eur. Phys. J. A **6** (1999) 415

depends on

- ▶ temperature  $T$
- ▶ effective baryon density  $\rho_{\text{eff}}$
- ▶ pion chemical potential  $\mu_{\pi}$

$$\rho_{\text{eff}} = \rho_N + \rho_{\bar{N}} + \frac{1}{2} (\rho_R + \rho_{\bar{R}})$$

- ▶ reproduces excess in experimental data


- ▶ CERES
- ▶ NA60
- ▶ STAR (including BES)
- ▶ PHENIX with HBD

- ▶ at higher masses: include hadronic continuum radiation

E. V. Shuryak: Rev. Mod. Phys. **69** (1993) 1

# Space-time evolution of a heavy-ion collision


Au+Au at 1.23 AGeV ( $\sqrt{s_{NN}} = 2.4$  GeV)  $\rightarrow$  HADES energy regime






# Description of the fireball evolution

## Coarse-graining of hadronic transport





- ▶ “combine” the advantages of both descriptions: hydrodynamics & transport
- ▶ simulate events with a transport model
  - ensemble average to obtain smooth space-time distributions
- ▶ divide space-time evolution into 4-dimensional cells
  - 21 x 21 x 21 space cells (1fm<sup>3</sup>), 30 time steps → ~ 280 k cells
- ▶ determine for each cell the bulk properties like  $T$ ,  $\rho_B$  &  $v_{\text{coll}}$
- ▶ calculate dilepton rates based on these inputs
  - parameterization of RW in-medium spectral function
- ▶ sum up the contributions of all cells
- ▶ similar approaches by
  - ▶ Huovinen *et al.*: PRC **66** (2002) 014903
  - ▶ Endres *et al.*: PRC **91** (2015) 054911, PRC **92** (2015) 014911, PRC **93** (2016) 054901, PRC **94** (2016) 024912



# Local thermalization

## Momentum distributions of nucleons ( $n_{\text{coll}} \geq 3$ ) & evolution of $n_{\text{coll}}$




- ▶ Gaussian shaped  $p_z$  distribution builds up for nucleons with  $n_{\text{coll}} \geq 3$
- ▶  $m_t$  spectra have exponential shape
- ▶ check for every cell  $\rightarrow$  deviations are kept in space-time evolution

# Determination of bulk properties

(Baryon) density, collective flow velocity & temperature

- ▶ baryon density via 4-current
- ▶ Lorentz-boost to local rest frame (LRF) where the baryon current vanishes




- ▶ in Boltzmann approximation

$$\frac{d^3N}{d\vec{p}} = \frac{d^3N}{dp_z p_t dp_t d\theta} \propto \exp(-E/T)$$

↓

$$\frac{1}{m_t^{3/2}} \frac{dN}{dm_t} \propto \exp(-m_t/T)$$




- ▶ fill  $m_t$  spectra with particle momenta in LRF (mean flow  $v_{\text{coll}}$  vanishes)
- ▶ fit exponential function to extract  $T$  (species of choice: pions)



# Out of chemical equilibrium?

## Build-up of effective chemical potentials



- ▶ thermal emission rates assume chemical equilibrium
- ▶ chemical non-equilibrium possible, e.g. after chemical freeze-out
  - ▶ no more inelastic interactions → pion number conserved
  - ▶ system in thermal equilibrium cools down further → over-population of pions
  - ▶ build-up of an effective chemical potential  $\mu_\pi$
- ▶ induces a factor  $(z_\pi)^\kappa$  in the dilepton rates with the fugacity  $z = \exp\left(\frac{\mu_\pi}{T}\right)$ 
  - ▶ exponent  $\kappa$  reflects the main production mechanism of  $\rho$  mesons
  - ▶ at HADES energies UrQMD suggests  $\kappa = 1.12$



# Time-evolution


Au+Au at 1.23 AGeV

- ▶ evolution of  $T$ ,  $\rho_{\text{eff}}$  and  $\mu_{\pi}$  in the central cube of  $7 \times 7 \times 7$  cells
- ▶ trajectories of the cells in the temperature-density plane



# Interplay temperature – fireball volume



Au+Au at 1.23 AGeV



# Dileptons as fireball probes

## Au+Au at 1.23 AGeV

- ▶ time evolution of cumulative dilepton yield in mass window  $M = 0.3-0.7 \text{ GeV}/c^2$
- ▶ active radiation window  $\sim 13 \text{ fm}/c$  follows build-up of collective medium flow  $\Rightarrow$  fireball lifetime
- ▶ strong medium effects on  $\rho$ -meson  $\Rightarrow$  remarkably structure-less low-mass spectrum
- ▶  $dR_{ll}/dM \propto (MT)^{3/2} \exp(-M/T)$
- ▶ inverse slope parameter:  $T_S = 88 \pm 5 \text{ MeV}$  in IMR,  $T_S = 64 \pm 5 \text{ MeV}$  in LMR






T. Galatyuk *et al.*: Eur. Phys. J. A **52** (2016) 131

# Dileptons as fireball probes

Ar+KCl at 1.76 AGeV ( $\sqrt{s_{NN}} = 2.6$  GeV)



- ▶ evolution of  $T$ ,  $\rho_{\text{eff}}$  and  $\mu_{\pi}$  in the inner cube of  $5 \times 5 \times 5$  cells
- ▶ invariant mass spectrum for the thermal radiation
- ▶ window for dilepton radiation & build-up of collectivity  $\sim 8 \text{ fm}/c$





# Excitation function of dilepton production




## Yield in low-mass window tracks fireball lifetime



- ▶ fireball dominated by incoming nucleons at lower energies
- ▶ number of charged particles  $N_{ch}$  not a good proxy for thermal excitation energy
- ▶ normalization to number of charged pions  $N_{\pi}$
- ▶ lifetime from dilepton yield in mass window  $0.3\text{-}0.7 \text{ GeV}/c^2$ :  $\frac{N_{l+l-}}{N_{\pi^\pm}} \cdot 10^6 \simeq 1.45 \cdot \tau_{fb}$


# Comparison to experimental excess spectra

Ar+KCl at 1.76 AGeV & Au+Au at 1.23 AGeV (min. bias)



# Exploring the QCD phase diagram –


## – with dileptons



$\frac{\langle \bar{q}q \rangle_{T, \mu_B}}{\langle \bar{q}q \rangle_{T=0, \mu_B=0}}$  : B.J. Schaefer and J. Wambach

- ▶ chemical freeze-out from measured particle yields analyzed with SHM
- ▶ trajectories extracted from inner cube of cells with coarse-grained UrQMD
- ▶ time-window of dilepton emission
  - ▶ radiation stops shortly after chemical freeze-out
  - ▶ access to hot and dense stage of the heavy-ion collision

# Exploring the QCD phase diagram – – with dileptons



- ▶ NA60 intermediate mass  $\mu^+\mu^-$
- ▶ trajectories at SIS18
- ▶ trajectories at SIS100

$\frac{\langle q\bar{q} \rangle_{T, \mu_B}}{\langle q\bar{q} \rangle_{T=0, \mu_B=0}}$  : B.J. Schaefer and J. Wambach  
 ▲ NA60 ( $\mu^+\mu^-$ ) : H.J. Specht: AIP Conf. Proc. 1322 (2010)

# Summary

THANK YOU FOR YOUR ATTENTION !


- ▶ dileptons are excellent fireball probes
  - ▶ thermometer & chronometer
  - ▶ new insights into the matter created under extreme conditions
- ▶ thermal dilepton spectra from highest to lowest energies
  - ▶ realistic thermal dilepton emission rates
  - ▶ accurate description of fireball evolution in terms of  $T$ ,  $\rho_{\text{eff}}$ ,  $v_{\text{coll}}$  and  $\mu_{\text{T}}$
  - ▶ coarse-graining of hadronic transport at SIS energies
- ▶ baseline for future experimental explorations
  - ▶ any significant deviation can indicate new physics!

---

# Backup slides

---



# Excitation function of hadron yields



A. Andronic, arXiv:1407.5003

# Final-state pion cocktail

PID of mother particle





- ▶ 90% (95%) of pions are emitted before 30 (40) fm/c






# Final-state pion spectra

- ▶ Dominant contribution:  $\Delta(1232)$  decays (cyan)
- ▶ Many more resonances contribute especially at higher  $p_T$






# Final-state pion spectra: density dependent

$\rho/\rho_0 < 0.1$  at emission or  $t_{\text{emission}} > 30 \text{ fm}/c$



# Final-state pion spectra: density dependent

$\rho/\rho_0 < 0.5$  at emission or  $t_{\text{emission}} > 30 \text{ fm}/c$



# Final-state pion spectra: density dependent

$\rho/\rho_0 > 1$  at emission

