



Contribution ID: 21

Type: not specified

Observation of a long regular band structure at high spin in ^{89}Zr

The high spin states of an atomic nucleus provides understanding of it's shape, dynamics and interaction between the constituent nucleons. The high spin states of a nucleus near the shell closure provide unique test bench for studying the excitations of valance shell nucleons.

Among some of the very interesting phenomena that have been observed in different doubly magic nuclei is the evolution of collectivity at high

spin [1]. Nuclei around ^{90}Zr having $Z=40$ shell gap and $N = 50$ major shell closure are perfect testing ground to probe such emergent phenomena. Recently, the high spin states of ^{89}Zr was populated with $^{13}\text{C}(^{80}\text{Se},5n)$ reaction using ^{13}C beam at 50 and 60 MeV from TIFR-BARC pelletron facility [2,3]. The gamma-rays decaying from the residual nuclei were detected using the Indian National Gamma Array (INGA) spectrometer with 18 Compton suppressed clover HPGe detectors [4].

The ^{89}Zr levelscheme has been extended up to $49/2 \hbar\omega$ and a regular dipole band has been observed. The spin and parity of the levels have been assigned using DCO and polarisation method. The lifetimes of some of the states have also been measured. The cranked Nilsson Strutinsky model has been used to describe the observed rotational bands. The calculation suggests that the the band structure is originated from the longest axis rotation of the triaxially deformed nucleus.

Primary author: Dr SAHA, Sudipta

Co-authors: SINGH, Ajay K. (Indian Institute of Technology Kharagpur(IITKGP)); Dr SETHI, Jasmine (TIFR); Dr SINGH, Purnima (TIFR); Prof. PALIT, Rudrajyoti (TIFR); Ms BISWAS, Sayani (TIFR); Dr NAG, Somnath (Department of Physics, National Institute of Technology Raipur)

Presenter: Dr SAHA, Sudipta