




The CBM Experiment: Program and Status

Norbert Herrmann
Heidelberg Univ.



Dense Baryonic Matter



Neutron stars

Temperature
 $T < 20 \text{ MeV}$

Density
 $\rho < 10 \rho_0$

Lifetime
 $T \sim \text{infinity}$




Neutron star merger

Temperature
 $T < 70 \text{ MeV}$

Density
 $\rho < 3\rho_0$

Reaction time
(GW170817)
 $T \sim 10 \text{ s}$

Heavy ion collisions at SIS100




Compressed Baryonic Matter

Temperature
 $T < 120 \text{ MeV}$

Density
 $\rho < 8\rho_0$

Reaction time
 $t \sim 10^{-23} \text{ s}$

CBM – Goals



Mission:


Systematically explore QCD matter at large baryon densities with high accuracy and rare probes.

Outline:

- Current experimental knowledge
- Experimental and theoretical expectations / speculations
- Experiment setup
- Status of developments
- Planned Fair Phase-0 and Phase-1 measurements


Baryon densities at SIS100

Baryon densities in central Au+Au collisions






I.-C. Arsene et al., Phys. Rev. C 75, 24902 (2007)



5 A GeV



10 A GeV



Observables in Heavy – Ion Collisions




Hard probes
(initial state)


Penetrating probes
(integrate over collision history)
Relicts
(produced in dense phase)

Freeze-out
(final state particles)
Thermalized (?) hadrons

Chemical Freeze-out data



Analyses in framework of
Statistical Hadronisation Model



High energies:
grandcanonical ensemble

$$n_i(\mu, T) = \frac{N_i}{V} = -\frac{T}{V} \frac{\partial \ln Z_i}{\partial \mu} = \frac{g_i}{2\pi^2} \int \frac{p^2 dp}{e^{\frac{E_i - \mu_i}{T}} \pm 1}$$


$$\mu_i = \mu_B B_i + \mu_S S_i + \mu_{I_3} I_{3,i}$$

Lower energies / small systems:
canonical ensemble,
strangeness suppression factor γ_s

Equilibrium achieved in small systems?

Equilibrium as signature for phase transition?

Freeze-out line at large baryon densities
as phase boundary to quarkyonic matter ?




A. Andronic et al.,
Nucl. Phys. A837 (2010) 65

HADES: Sub-threshold Ξ^- - production




Ar+KCl reactions at 1.76A GeV


- Ξ^- yield by appr. factor 25 higher than thermal yield




G. Agakishiev et al. (HADES), PRL103, 132301, (2009)




Reminder: Subthreshold Kaon – measurements (KAOS at SIS18)



C.Sturm et al. (KaoS), PRL 86 (2001) 39



C. Fuchs et al., PRL 86 (2001) 1974




Strong sensitivity to Equation Of State
due to multistep production
(formation of nucleon resonances)
=> soft EOS (K=200 MeV)


Equation of State & Neutron stars



C. Fuchs,
Prog. Part. Nucl. Phys. 56 (2006) 1



P.B. Demorest (2010)
doi:10.1038/nature09466




Soft EOS (Skyrme, $K = 200$ MeV) is not repulsive enough to allow for a neutron star with 2 solar masses.


DBHF BONN A corresponds to AP4, however, does not contain strange baryons.

Stiffening of EOS must occur in the range of densities up to $4 \rho_0$ (SIS100 energy range).

Final state particle abundance



Particle yields from central Au + Au collisions



C.Blaum, SQM2017)


Strange and charmed particle production thresholds in pp - collisions

reaction	\sqrt{s} (GeV)	T _{lab} (GeV)
$pp \rightarrow K^+ \Lambda p$	2.548	1.6
$pp \rightarrow K^+ K^- pp$	2.864	2.5
$pp \rightarrow K^+ K^+ \Xi^- p$	3.247	3.7
$pp \rightarrow K^+ K^+ K^+ \Omega^- n$	4.092	7.0
$pp \rightarrow \Lambda \bar{\Lambda} pp$	4.108	7.1
$pp \rightarrow \Xi^- \bar{\Xi}^+ pp$	4.520	9.0
$pp \rightarrow \Omega^- \bar{\Omega}^+ pp$	5.222	12.7
$pp \rightarrow J/\Psi pp$	4.973	12.2

Hyperons as probes of dense matter

PHSD interpretation of Ξ^- - production

A. Palmese et al. Phys.Rev. C94 (2016) no.4, 044912




Predicted sensitivities
of production yields:

strong dependence on
Chiral Symmetry Restoration (CSR)

Measurable dependence on
Equation of State (NL1, NL3)


Antihyperon – production



Prediction of PHSD transport model

(E. Bratkovskaya, W. Cassing)


I. Vassiliev, CBM, private communication



Large sensitivity to
partonic degrees of freedom
in SIS100 energy range
(deconfinement phase transition)


Mapping out the phase structure
requires systematic measurements.

CBM experimental setup (MSV)



- Tracking acceptance:
 $2^\circ < \theta_{\text{lab}} < 25^\circ$
- Free streaming DAQ
 $R_{\text{int}} = 10 \text{ MHz} (\text{Au+Au})$
with
 $R_{\text{int}} (\text{MVD}) = 0.1 \text{ MHz}$
- Software based event selection

CBM – Strategy





Exploration of QCD phase diagram
as international effort:

- | | |
|-------------|------------------|
| NA61 | @ SPS / CERN |
| BM@N | @ Nuclotron/JINR |
| STAR (F.t.) | @ RHIC/BNL |
| MPD | @ NICA / JINR |

CBM's unique feature
High statistics measurement of rare probes

CBM data processing system




Reaction rate: Au + Au, 10^7 collisions per second
Data rate: ~ 1 TB/s

Main features:

- radiation tolerant detectors and front-end electronics
- no hardware trigger of events,
- free streaming (triggerless) data,
- all detector hits with time stamps,
- software based event selection

CBM physics and observables



Eur.Phys.J. A53 (2017) 60

QCD equation-of-state

- collective flow of identified particles
- particle production at threshold energies

Phase transition

- excitation function of hyperons
- excitation function of LM lepton pairs

Critical point

- event-by-event fluctuations of conserved quantities

Chiral symmetry restoration at large ρ_B


- in-medium modifications of hadrons
- dileptons at intermediate invariant masses

Strange matter


- (double-) lambda hypernuclei
- Search for meta-stable objects (e.g. strange dibaryons)

Heavy flavour in cold and dense matter

- excitation function of charm production




CBM experimental setup (day-1)



- Tracking acceptance:
 $2^\circ < \theta_{\text{lab}} < 25^\circ$
 - Free streaming DAQ
- $R_{\text{int}} \approx 0.5 \text{ MHz (Au+Au)}$
- full bandwidth:
Det. – Entry nodes
reduced bandwidth
Entry nodes – Comp. farm
- with
 $R_{\text{int}} (\text{MVD}) = 0.1 \text{ MHz}$
- Software based event selection

Day-1 setup = MSV setup – Compute Performance - ECAL
Phase-1 = Day1 with full Compute Performance + ECAL


CBM day-1 – program




Observables: Strangeness and Dileptons

Excitation function of yields and phase-space distributions of multi-strange hyperons and lepton pairs in AA (C+C, Au+Au) collisions from 2-11 A GeV.
Search for hypernuclei (no data available in this energy range).


multi-strange hyperons




hypernuclei




dilepton invariant mass



Dileptons as probes for dense matter




[R. Rapp, H. v.Hees, PLB 753 (2016) 586]




LMR: ρ – chiral symmetry restoration
fireball space – time extension

IMR: access to fireball temperature
 ρ - a_1 chiral mixing


Measurement program:
e.g. excitation function of IMR - slope



- 1M Au+Au ($b=0$ fm), 8 AGeV
- IMR: S/B > 1/100
- Statistical accuracy of 10% requires ~1 week of beamtime




CBM Day 1 – unique measurements




Di-electron measurement

Full performance,
uses MVD, limited to 100 kHz




Au+Au, 8A GeV,


Hyperon measurements, e.g. Au+Au at 10A GeV :




Hypernuclei measurement,
e.g. Au + Au at 10A GeV




Di-Muon
LM measurement at 8A GeV




CBM – FAIR Phase 0 projects (2018 – 2022)




1. Install, commission and use 430 out of 1100 CBM RICH multi-anode photo-multipliers (MAPMT) including FEE in HADES RICH photon detector




2. Install, commission and use 10% of the CBM TOF modules including read-out chain at STAR/RHIC (BES II 2019/2020)



3. Upgrade BM@N experiment with 4 Silicon stations of CBM/STS design in the BM@N experiment at the Nuclotron JINR/Dubna (Au-beams in late 2020)



4. Install, commission and use the Project Spectator Detector at the BM@N experiment





5. mini CBM (mCBM@SIS18) demonstrator for full CBM data taking and analysis chain

CBM FAIR Phase 0 project at SIS18: mCBM



Demonstrator for full CBM data taking and analysis chain




the mCBM test-setup (“mini-CBM”) will focus on

- test of final detector prototypes
- free streaming data transport to a computer farm
- online reconstruction and event selection
- offline data analysis

under full load conditions (Au + Au, 10^7 interactions/s)



<https://cbm-wiki.gsi.de/foswiki/pub/Public/Documents/mcbm-proposal2GPAC-fullVersion.pdf>

mCBM performance benchmark








(Sub)threshold Λ – baryon reconstruction.



Event based MC simulation of 10^8 events
(measurement time: 10 s)



Acceptance
&
Efficiency




mCBM Cave (HTD)





- Modified switching magnet (HTD – MU1)
- New beam dump
- Additional shielding


FLUKA calculations:
 10^8 Au ions s^{-1} , 1.24 AGeV,
 2.5 mm Au target ($P_{int} = 10\%$)
 vertical section: **beam level**



Status of the cave reconstruction




Schedule of mCBM construction








Schedule

10/2017	cave & beam line: reconstruction started, procurement started
11/2017	mDAQ test stand @ Heidelberg operational
12/2017	beam dump mounted
03/2018	cave reconstruction completed
04/2018	mFLES cluster @ Green IT Cube installed
05/2018	beam line installed and commissioned
05/2018	installation of detector stations
06/2018	start commissioning w/o beam
08/2018	start commissioning with beam



eTOF & HPC software in STAR at RHIC (BNL)



Participating CBM groups:

- Tsinghua Univ. Beijing
- GSI Darmstadt
- TU Darmstadt
- Univ. Frankfurt
- Univ. Heidelberg
- USTC Hefei
- CCNU Wuhan

Test module installed	(Oct. 2016),
Module is operational	(Oct. 2016),
STAR DAQ interface	(Jan. 2017),
Full sector test	(Spring 2018),
Wheel installation	(Summer 2018),
BES II data taking	(2019/2020),
Transfer of modules to FAIR	(2021/22).

27

STAR – BES II physics program




Physics Program for the STAR/CBM eTOF Upgrade

The STAR Collaboration
The CBM Collaboration eTOF Group
(Dated: September 19, 2016)

arXiv:1609.05102v1 [nucl-ex]

Example: kaon acceptance




Topics to be studied with extended acceptance in energy range $\sqrt{s}_{NN} = 3 - 62$ GeV:

- Excitation function and phase-space distributions of hyperons, hypernuclei, anti-protons, ...
→ Equilibration, phase transitions
- Collective Flow (v_1 , v_2)
→ Equation-of-State, phase transitions
- Fluctuations of conserved quantum numbers (baryon, charge, strangeness)
→ Critical point
- Dilepton yields
→ Chiral symmetry restoration


STS & PSD in BM@N (JINR)



*Silicon Tracking
Stations*



Au beams up to 4.5 GeV/u




PSD calorimeter

Improvement in efficiency
& signal / background

BM@N timeline: NICA white paper
(Eur. Phys. J. A (2016) 213)

- 2018 Installation of PSD detector (MoU signed)
- 2020 Installation of 4 Si tracking stations (MoU signed)
- 2020 Au beams from Nuclotron



CBM – Collaboration: 55 institutions, 470 members



China:

CCNU Wuhan
Tsinghua Univ.
USTC Hefei
CTGU Yichang
Chongqing Univ.

Czech Republic:

CAS, Rez
Techn. Univ. Prague

France:

IPHC Strasbourg

Germany:

Darmstadt TU
FAIR
Frankfurt Univ. IKF
Frankfurt Univ. FIAS
Frankfurt Univ. ICS
GSI Darmstadt
Giessen Univ.
Heidelberg Univ. P.I.
Heidelberg Univ. ZITI
HZ Dresden-Rossendorf
KIT Karlsruhe
Münster Univ.
Tübingen Univ.
Wuppertal Univ.
ZIB Berlin

India:

Aligarh Muslim Univ.
Bose Inst. Kolkata
Panjab Univ.
Univ. of Jammu
Univ. of Kashmir
Univ. of Calcutta
B.H. Univ. Varanasi
VECC Kolkata
IOP Bhubaneswar
IIT Kharagpur
IIT Indore
Gauhati Univ.

Korea:

Pusan Nat. Univ.

Poland:

AGH Krakow
Jag. Univ. Krakow
Warsaw Univ.
Warsaw TU

Romania:

NIPNE Bucharest
Univ. Bucharest

Hungary:

KFKI Budapest
Eötvös Univ.

Russia:


IHEP Protvino
INR Troitzk
ITEP Moscow
Kurchatov Inst., Moscow
VBLHEP, JINR Dubna
LIT, JINR Dubna
MEPHI Moscow
PNPI Gatchina
SINP MSU, Moscow

Ukraine:


T. Shevchenko Univ. Kiev
Kiev Inst. Nucl. Research



CBM Scientists



Summary / Conclusion



CBM scientific program at SIS100 is unique

explore QCD matter at neutron star core densities

employ high statistics capability

to achieve high-precision of multi-differential observables

to enable rare processes as sensitive probes

CBM day-1 setup allows start of program with significant discovery potential

excitation function of hyperons production

excitation function of di-lepton production

study of hypernuclei

CBM Phase 0 activities targeted towards usage and understanding
of major components & production of visible physics results with CBM devices

CBM – RICH sensors & readout

in HADES at SIS18

CBM – TOF and HPC software

in STAR at RHIC/BNL

CBM – PSD and CBM - STS

in BM@N at Nuclotron/JINR

Integration of all subsystems & FLES

in mCBM at SIS18