

# Structure of Exotic Compounds

M. Bayar

Physics Department  
Kocaeli University

2nd EMMI Workshop on anti-matter, hyper-matter and  
exotica production at the LHC


09 November, 2017

Collaborators: E. Oset, C. W. Xiao, W.H. Liang, T. Uchino, P. Fernandez-Soler,  
Zhi-Feng Sun, Xiu-Lei Ren, B. Durkaya

- Introduction
- Formalism
  - The Faddeev equations under the Fixed Center Approximation
- The Light Unflavored and Strange Sectors
- The Charm Sector
- The Beauty Sector

# Introduction

Quantum Chromodynamics (QCD)  $\Rightarrow$  the theory behind the strong interactions.




At large distances ( $\geq 0.1$  fm):

- Confinement
- Quarks build coherent bound states - hadrons
- QCD perturbation theory is inapplicable

At small distances:

- Asymptotic freedom
- Quasi-free quark propagation
- QCD perturbation theory

Quantum Chromodynamics (QCD)  $\Rightarrow$  the theory behind the strong interactions.




Non-perturbative methods:


- Phenomenological quark models
- Lattice QCD
- Effective lagrangian methods
- QCD sum rules
- Chiral Perturbation theory
- .....

# Hadron Spectroscopy

- Ordinary hadrons




Meson ( $q\bar{q}$ )




Baryon ( $qqq$ )

- Gluonic excitations




Hybrid meson:  
 $q\bar{q}$  with gluonic excitations,




Glueball:  
only gluons, no valence quarks

- Multiquark states



Tetraquark:


Two quarks and two antiquarks,



Hadronic molecule:

composed of two or more color-neutral hadrons

- Pentaquark




four quarks and one antiquark bound together

# Formalism



# The Faddeev equations under the Fixed Center Approximation (FCA)

The FCA to the Faddeev equations is an effective tool to deal with multi-hadron interaction



- $T_1$ : all diagrams beginning with interaction in particle 1.
- $T_2$ : all diagrams beginning with interaction in particle 2.

$$T_1 = t_1 + t_1 G_0 T_2, \quad T_2 = t_2 + t_2 G_0 T_1, \quad T = T_1 + T_2$$

$$T = T_1 + T_2 = \frac{\tilde{t}_1 + \tilde{t}_2 + 2 \tilde{t}_1 \tilde{t}_2 G_0}{1 - \tilde{t}_1 \tilde{t}_2 G_0^2}.$$

The function  $G_0$  :

$$G_0(s) = \int \frac{d^3 \vec{q}}{(2\pi)^3} F_R(q) \frac{1}{q^{02} - \vec{q}^2 - m_3^2 + i\epsilon}, \quad q^0(s) = \frac{s + m_3^2 - M_R^2}{2\sqrt{s}}.$$

$F_R(q)$  is the cluster form factor

$$F_R(q) = \frac{1}{\mathcal{N}} \int_{|\vec{p}| < \Lambda', |\vec{p} - \vec{q}| < \Lambda'} d^3 \vec{p} \frac{1}{2E_1(\vec{p})} \frac{1}{2E_2(\vec{p})} \frac{1}{M_R - E_1(\vec{p}) - E_2(\vec{p})} \frac{1}{2E_1(\vec{p} - \vec{q})} \frac{1}{2E_2(\vec{p} - \vec{q})} \frac{1}{M_R - E_1(\vec{p} - \vec{q}) - E_2(\vec{p} - \vec{q})}, \quad (1)$$

$$\mathcal{N} = \int_{|\vec{p}| < \Lambda'} d^3 \vec{p} \left( \frac{1}{2E_1(\vec{p})} \frac{1}{2E_2(\vec{p})} \frac{1}{M_R - E_1(\vec{p}) - E_2(\vec{p})} \right)^2,$$

(J.Yamagata-Sekihara, J. Nieves, E. Oset Phys. Rev. D 83,014003 (2011))

# Two Body Scattering

The Bethe-Salpeter equation in coupled channels

$$t = V + VGt$$




A loop function of pseudoscalar and vector mesons  $G_I$ :

$$G_I(\sqrt{s}) = i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(P-q)^2 - M_I^2 + i\epsilon} \frac{1}{q^2 - m_I^2 + i\epsilon}$$

## The Light Unflavored and Stange Sectors

# Description of $\rho(1700)$ as a $\rho K \bar{K}$ system

(M. Bayar, W. H Liang, T. Uchino and C. W Xiao, Eur.Phys.J. A50 (2014))



- A cluster of two bound particle ( $K\bar{K}$  ( $l = 0$ ),  $f_0(980)$ )
- $K\bar{K}$ +coupled channel  $\rightarrow f_0(980)$  (J. A. Oller, E. Oset, Nuclear Physics A 620,438 (1997))  
 $\Rightarrow \rho - (K\bar{K}) \rightarrow$  one needs  $t$  for  $\rho K(\rho\bar{K})$
- $\rho K(\rho\bar{K})$  unitarized scattering amplitude, (L. S. Geng, E. Oset, L. Roca and J. A. Oller, Phys.

Rev. D 75, 014017 (2007), L. Roca, E. Oset and J. Singh, Phys. Rev. D 72, 014002 (2005). )

# $\rho K(\rho \bar{K})$ unitarized scattering amplitude

- The Bethe-Salpeter equation in coupled channels:  $t = [1 + V\hat{G}]^{-1}(-V)\vec{\epsilon} \cdot \vec{\epsilon}'$
- The two meson loop function:

$$G_I(\sqrt{s}) = i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(P-q)^2 - M_I^2 + i\epsilon} \frac{1}{q^2 - m_I^2 + i\epsilon}$$

- In the dimensional regularization scheme the loop function:

$$G_I(\sqrt{s}) = \frac{1}{16\pi^2} \left\{ a(\mu) + \ln \frac{M_I^2}{\mu^2} + \frac{m_I^2 - M_I^2 + s}{2s} \ln \frac{m_I^2}{M_I^2} \right. \\ \left. + \frac{q_I}{\sqrt{s}} \left[ \ln(s - (M_I^2 - m_I^2) + 2q_I\sqrt{s}) \right. \right. \\ \left. \left. + \ln(s + (M_I^2 - m_I^2) + 2q_I\sqrt{s}) \right. \right. \\ \left. \left. - \ln(-s + (M_I^2 - m_I^2) + 2q_I\sqrt{s}) \right. \right. \\ \left. \left. - \ln(-s - (M_I^2 - m_I^2) + 2q_I\sqrt{s}) \right] \right\}$$

- $q_I$  determined at the center of mass frame,  $q_I = \frac{\sqrt{[s - (M_I - m_I)^2][s - (M_I + m_I)^2]}}{2\sqrt{s}}$
- $\mu \Rightarrow$  a scale parameter in this scheme,  $a(\mu) \Rightarrow$  the subtraction constant

# For the normalization

The S-matrix for the single scattering

$$S^{(1)} = -i(2\pi)^4 \delta^4(k_\rho + k_{f_0} - k'_\rho - k'_{f_0}) \times \frac{1}{\mathcal{V}^2} \frac{1}{\sqrt{2\omega_\rho}} \frac{1}{\sqrt{2\omega'_\rho}} \frac{1}{\sqrt{2\omega_K}} \frac{1}{\sqrt{2\omega'_K}} t,$$

S-matrix for the double scattering,

$$S^{(2)} = -i(2\pi)^4 \delta^4(k_\rho + k_{f_0} - k'_\rho - k'_{f_0}) \times \frac{1}{\mathcal{V}^2} \frac{1}{\sqrt{2\omega_\rho}} \frac{1}{\sqrt{2\omega'_\rho}} \frac{1}{\sqrt{2\omega_K}} \frac{1}{\sqrt{2\omega'_K}} \\ \times \frac{1}{\sqrt{2\omega_K}} \frac{1}{\sqrt{2\omega'_K}} \int \frac{d^3q}{(2\pi)^3} F_{f_0}(q) \frac{1}{q^{02} - \vec{q}^2 - m_\rho^2 + i\epsilon} tt,$$


The full three-body scattering is given by

$$S = -i(2\pi)^4 \delta^4(k_\rho + k_{f_0} - k'_\rho - k'_{f_0}) \frac{1}{\mathcal{V}^2} \frac{1}{\sqrt{2\omega_\rho}} \frac{1}{\sqrt{2\omega'_\rho}} \frac{1}{\sqrt{2\omega_{f_0}}} \frac{1}{\sqrt{2\omega'_{f_0}}} T.$$

Using the low energy reduction,  $\sqrt{2\omega} \sim \sqrt{2m} \Rightarrow \tilde{t} = \frac{2m_{f_0}}{2m_K} t.$

● Finally  $T = T_1 + T_2 = \frac{\tilde{t}_1 + \tilde{t}_2 + 2 \tilde{t}_1 \tilde{t}_2 G_0}{1 - \tilde{t}_1 \tilde{t}_2 G_0^2}.$

# Description of $\rho(1700)$ as a $\rho K \bar{K}$ system








	single	full	full + $\Gamma_\rho$	PDG
Mass (MeV)	1777.9	1734.8	1748.0	$1720 \pm 20$
Width (MeV)	144.4	63.7	160.8	$250 \pm 100$




# A description of the $f_2(1270)$ , $\rho_3(1690)$ , $f_4(2050)$ , $\rho_5(2350)$ and $f_6(2510)$ resonances as multi- $\rho(770)$ states (L. Roca and E. Oset PRD 82 054013 (2010).)

$\rho$ - $\rho$  interaction in the hidden gauge approach (R. Molina, D. Nicmorus, E. Oset PRD78 (2008))

- scattering of  $f_2$  with  $f_2 \Rightarrow$  the  $f_4$
- $\rho$  interaction with  $f_4 \Rightarrow \rho_5$
- $f_2$  with  $f_4 \Rightarrow f_6$



Modulus squared of the unitarized multi- $\rho$  amplitudes. Solid line: full model  $\Lambda' |_{f_4} = 1500$  MeV; dashed line: full model  $\Lambda' |_{f_4} = 875$  MeV; dotted line: only single-scattering contribution.




Masses of the dynamically generated states as a function of the number of constituent  $\rho(770)$  mesons,  $n_\rho$ .


# On the nature of the $K_2^*(1430)$ , $K_3^*(1780)$ , $K_4^*(2045)$ , $K_5^*(2380)$ and $K_6^*$ as

$K^*$ -multi- $\rho$  states, (J. Yamagata, L.Roca and E. Oset PRD 82 094017 (2010).)





$\rho K^*$  interaction  $\Rightarrow K_2^*(1430)$  (L. S. Geng and E. Oset, Phys. Rev. D 79 (2009))



etc



	A	B ( $b_1 b_2$ )
two-body	$\rho$	$K^*$
three-body	$K^*$	$f_2(\rho\rho)$
four-body	$\rho$	$K_2^*(\rho K^*)$
five-body	$f_2$	$K_2^*(\rho K^*)$
five-body	$K^*$	$f_4(f_2 f_2)$
five-body	$\rho$	$K_4^*(f_2 K_2^*)$
six-body	$K_2^*$	$f_4(f_2 f_2)$
six-body	$f_2$	$K_4^*(f_2 K_2^*)$



## Results for the masses of the dynamically generated states:



generated resonance	amplitude	mass (MeV), PDG	mass (MeV) only single scatt.	mass (MeV) full model
$K_2^*(1430)$	$\rho K^*$	$1429 \pm 1.4$	—	1430
$K_3^*(1780)$	$K^* f_2$	$1776 \pm 7$	1930	1790
$K_4^*(2045)$	$f_2 K_2^*$	$2045 \pm 9$	2466	2114
$K_5^*(2380)$	$K^* f_4$	$2382 \pm 14 \pm 19$	2736	2310
$K_6^*$	$K_2^* f_4 - f_2 K_4^*$	—	3073-3310	2661-2698

## The Charm Sector

# The $\rho D\bar{D}$ system

M. Bayar, B. Durkaya Phys.Rev. D92 2015

- Clusters :  $D\bar{D} \Rightarrow X(3700)(I=0)$  and  $\rho D \Rightarrow D_1(2420)(I=1/2)$   
The  $\rho(D\bar{D})_{X(3700)}$  and  $\bar{D}(\rho D)_{D_1(2420)}$
- $\rho D(D_1(2420)) \Rightarrow \pi D^*, D\rho, KD_s^*, D_s K^*, \eta D^*, D\omega, \eta_c D^*, DJ/\psi, (I=1/2)$  and  $\pi D^*, D\rho, (I=3/2)$  (D. Gamermann, E. Oset, D. Strottman, and M. J. Vicente Vacas, PRD76(2007), D. Gamermann and E. Oset, EPJA33(2007))
- $D\bar{D}(X(3700)) \Rightarrow D\bar{D}, K\bar{K}, \pi\bar{\pi}, \eta\eta, \eta_c\eta, D_s\bar{D}_s, (I=0), D\bar{D}, K\bar{K}, \pi\bar{\pi}, \pi\eta, \eta_c\pi, (I=1)$  (J. A. Oller and E. Oset, Nucl. Phys. A620 438 (1997), 4652 (1999))



$\rho(D\bar{D})_{X(3700)}, m \sim 4320 \text{ MeV}, \Gamma \sim 25 \text{ MeV}, I=1$      $\bar{D}(\rho D)_{D_1(2420)}, I=1, m \sim 4256 \text{ MeV}, \Gamma \sim 30 \text{ MeV}$

PDG: the  $X(4360), I^G(J^{PC}) = ?^?(1^{--}), m \simeq 4341 \text{ MeV}, \Gamma \sim 102 \pm 9 \text{ MeV}$

the  $X(4260), I^G(J^{PC}) = ?^?(1^{--}), m \simeq 4230 \text{ MeV}, \Gamma \sim 55 \pm 19 \text{ MeV}$




# The $\rho D^* \bar{D}^*$ System with $J = 3$

M. Bayar, X. L. Ren, E. O, Eur. Phys. J. A 2015

$D^* \bar{D}^*$  cluster with  $J^P = 2^+ \Rightarrow$  peak around 3920 MeV  $\Rightarrow$  the X(3915) or the Z(3940)  
(with  $l = 0$ )

the  $\rho$  interact with  $D^* \bar{D}^*$





A state with the  $l = 1$ ,  $J^P = 3^-$  and hidden charm is predicted around 4330 MeV.



# The $DK\bar{K}$ system

V. R. Debastiani, J. M. Dias, E. Oset, Phys.Rev. D96 (2017)



Narrow bound state around  $Df_0(980)$  threshold (2855 MeV) in both prescriptions.

$M_{DK\bar{K}} = 2833 - 2858$  MeV,


QCD Sum Rules:  $M_{Df_0} = (2926 \pm 237)$  MeV, Full Faddeev equations:  $M_{Df_0} = 2890$

MeV, (A. Martinez Torres, K. P. Khemchandani, M. Nielsen and F. S. Navarra, Phys. Rev. D 87, no. 3, 034025 (2013))

# A prediction of $D^*$ -multi- $\rho$ Molecular States



C. W. Xiao, M. Bayar, E. Oset, Phys.Rev. D86 (2012) 094019

- $\rho\rho$  interaction in  $l = 0$  and  $S = 2$  is very strong  
 $\Rightarrow f_2(1270)$  is a molecule of two  $\rho(770)$
  - $\rho D^*$  interaction in  $l = 1/2$  and  $S = 2$  is also very strong  
 $\Rightarrow D_2^*(2460)$  is a molecule of  $\rho(770)$  and  $D^*$
- $\Rightarrow$  Hence,  $\rho\rho$  and  $\rho D^*$  are the clusters



particles:	3	R (1,2)	amplitudes
Two-body	$\rho$	$D^*$	$t_{\rho D^*}$
	$\rho$	$\rho$	$t_{\rho\rho}$
Three-body	$D^*$	$f_2(\rho\rho)$	$T_{D^*} - f_2$
	$\rho$	$D_2^*(\rho D^*)$	$T_{\rho - D_2^*}$
Four-body	$D_2^*$	$f_2(\rho\rho)$	$T_{D_2^*} - f_2$
	$f_2$	$D_2^*(\rho D^*)$	$T_{f_2 - D_2^*}$
Five-body	$D^*$	$f_4(f_2 f_2)$	$T_{D^*} - f_4$
	$\rho$	$D_4^*(f_2 D_2^*)$	$T_{\rho - D_4^*}$
Six-body	$D_2^*$	$f_4(f_2 f_2)$	$T_{D_2^*} - f_4$
	$f_2$	$D_4^*(f_2 D_2^*)$	$T_{f_2 - D_4^*}$

### Three-body interaction:



2800-2850 MeV ( $I_{total} = \frac{1}{2}$ ) ———


$\sim 400$  MeV below  $D^* - f_2$  thr. ———

3120 MeV ( $I_{total} = \frac{3}{2}$ )

$|T_{\rho - D_2^*}^{l=3/2}|^2 \ll (30 \text{ times}) |T_{\rho - D_2^*}^{l=1/2}|^2$


**New  $D_3^*$  state** ; mixture of  $D^* - f_2$  and  $\rho - D_2^*$  m  $\sim 2800$ -2850 MeV  $\Gamma \sim 60$ -100 MeV

## Four and Five-body interactions:



3200 MeV,  $\Gamma \sim 200$  MeV — 3075 MeV,  $\Gamma \sim 400$  MeV

New  $D_4^*$  resonance ;  $m \sim 3075$ -3200 MeV,  $\Gamma \sim 200$ -400 MeV



3375 MeV,  $\Gamma \sim 200$  MeV — 3360 MeV,  $\Gamma \sim 400$  MeV


New  $D_5^*$  resonance ;  $m \sim 3360$ -3375 MeV,  $\Gamma \sim 200$ -400 MeV

# The Beauty Sector

# States of $\rho B^* \bar{B}^*$ with $J = 3$

M. Bayar, P. Fernandez-Soler, Zhi-Feng Sun, E. Oset, Eur.Phys.J. A52 (2016) no.4, 106

- A  $\rho$  meson and a  $B^* \bar{B}^*$  cluster
- The  $B^* \bar{B}^*$  cluster  $\Rightarrow$  the  $J = 2, I = 0$  (A. Ozpineci, C.W. Xiao, E. Oset, Phys.Rev. D88 (2013) 034018)




- Taking the  $\rho B^*$  interaction in  $J = 2$  (P. F. Soler, Z. F. Sun, J. Nieves and E. Oset, Eur.Phys.J. C76 (2016))
- We find a  $I(J^{PC}) = 1(3^{--})$  state of mass  $10987 \pm 40$  MeV and width  $40 \pm 15$  MeV

# On the binding of the $B\bar{D}\bar{D}$ system

J. M. Dias, V. R. Debastiani, L. Roca, S. Sakai, E. Oset, arXiv:1709.01372

- $BD$  cluster and  $\bar{D}$  scattering from that cluster
- $I(J^P) = 1/2(0^-)$  bound state for the  $B\bar{D}\bar{D}$  system at an energy around 8925 – 8985 MeV



$|T_{B\bar{D}\bar{D}}|^2$  with prescriptions I, II for  $\sqrt{s_{DB}}$ ,  $\sqrt{s_{DD}}$  and  $q_{\max} = 600$  MeV with and without considering width (from  $\eta\eta$  channel) for the  $X(3700)$  through the  $D\bar{D}$  interaction. The two curves with  $\eta\eta$  channel were multiplied by a factor  $10^4$  for comparison.

- The FCA to the Faddeev equations is an effective tool to deal with multi-hadron interaction
- $\rho(1700)$  appears as resonance of  $\rho K \bar{K}$
- Multirho states could be identified with meson states of increasing spin
- $K^*$  -multirho states can also be identified with  $K^*$  states of increasing spin
- In the charm sector the method is repeated and new charmed resonances,  $D_3^*$ ,  $D_4^*$ ,  $D_5^*$  and  $D_6^*$  are predicted
- The method is expanded to the beauty sector ....



**Thank you for your attention!**