

Chiral Symmetry Restoration by Parity Doubling and the Structure of Neutron Stars

M. Marczenko, D. Blaschke, K. Redlich, C. Sasaki

Institute of Theoretical Physics,
University of Wrocław, Poland

Phys. Rev. D97 (2018) no.3, 036011,
Phys. Rev. D98 (2018) no.10, 103021

EMMI Workshop, GSI
28.03.2019



Uniwersytet
Wrocławski



NATIONAL SCIENCE CENTRE
POLAND

Common approach to EoS


Hadronic EoS: p^+ , n^+
(incomplete chiral physics)

+

Quark EoS
(chiral physics)

↓


Maxwell construction
(deconfinement)




courtesy: N.-U. F. Bastian

- Striking problem: No chiral physics in the resulting EoS

Common approach to EoS



Parity doubling in lattice QCD Aarts et al, JHEP 1706, 034 (2017)



- Imprint of chiral symmetry restoration in the baryonic sector
- Expected to occur at low temperature

Particle Identification

p

n

$N(1535) \ 1/2^-$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$


$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } ****$$

C. Patrignani et al. (Particle Data Group), Chin. Phys. C, 40, 100001 (2016) and 2017

Parity doubling in SU(2) chiral models DeTar, Kunihiro PRD 39 (1989)

$$m^{\pm} = \frac{1}{2} \left[\sqrt{(g_1 + g_2)^2 \sigma^2 + 4m_0^2} \mp (g_1 - g_2) \sigma \right] \xrightarrow{\sigma \rightarrow 0} m_0$$




Parity doublet model + quark-meson coupling



Statistical confinement:

- UV cutoff for nucleons: $f_N \rightarrow \theta (\alpha^2 b^2 - \mathbf{p}^2) f_N$
- IR cutoff for quarks: $f_q \rightarrow \theta (\mathbf{p}^2 - b^2) f_q$
- α - model parameter




- b - scalar field

$$V_b = -\frac{1}{2}\kappa_b^2 b^2 + \frac{1}{4}\lambda_b b^4$$

$$\begin{aligned} b(\mu_B = 0) &> 0 && \text{favors nucleons} \\ b(\mu_B \rightarrow \infty) &= 0 && \text{favors quarks} \end{aligned}$$




Phase Diagram for Isospin-Symmetric Matter

- 1st order deconfinement transition
- Order of chiral transition (from low to high α)
 - 1st order $\rightarrow$ critical point $\rightarrow$ crossover
- Sequential phase transitions (may coincide for smaller m_0)






MM, C. Sasaki, Phys. Rev. D **97** 036011 (2018)

Higher-Order Cumulants




Higher-Order Cumulants in pQM model

Skokov, Friman, Redlich, PRC 83 (2011)




Equation of State Under NS Conditions

- $\alpha \rightarrow$ stiffening of EoS
- $\alpha \rightarrow$ strength of the chiral phase transition





Mass-Radius Relation


- chiral transition in high-mass part of the sequence
- $2M_{\odot}$ with chirally restored but confined core
- deconfinement above $2M_{\odot}$



Different Realizations of $2.01(4) M_{\odot}$ Stars



Composition of a Compact Star Matter



Threshold for Direct URCA: Conventional Scenario


- d.o.f.: p^+ , n^+ , e , μ

- Charge neutrality

$$\rho_{p^+} = \rho_e + \rho_\mu$$

- Momentum conservation

$$f_{n^+} \leq f_{p^+} + f_e$$



$$Y_{p^+} = \frac{1}{1 + (1 + \sqrt[3]{Y_e})^3} \Rightarrow 11\% - 15\%, \quad Y_e = \frac{\rho_e}{\rho_e + \rho_\mu}$$

Threshold for Direct URCA: Parity Doubling

- χ -symmetry broken

- d.o.f.: p^+ , n^+ , e , μ

- Charge neutrality

$$\rho_{p^+} = \rho_e + \rho_\mu$$

- Momentum conservation

$$f_{n^+} \leq f_{p^+} + f_e$$

- Proton Fraction Threshold

$$\frac{1}{1 + (1 + \sqrt[3]{Y_e})^3} \Rightarrow 11\% - 15\%$$

- χ -symmetry restored

- d.o.f.: p^+ , n^+ , p^- , n^- , e , μ

- Charge neutrality

$$\rho_{p^+} + \rho_{p^-} = 2\rho_{p^+} = \rho_e + \rho_\mu$$


- Momentum conservation

$$f_{n^+} \leq f_{p^+} + f_e$$

- Proton Fraction Threshold



$$\frac{1}{1 + (1 + \sqrt[3]{Y_e})^3} \Rightarrow 8\% - 11\%$$

Threshold for Direct URCA




- DU triggered already in chirally broken phase
- DU excluded below BRP (due to cooling phenomenology)

Constraints from GW170817




- $\Lambda = \frac{2}{3} k_2 C^{-5}$, where $C = M/R$ is compactness
- stiff EoS are rather excluded
- constraint on radius $R < 13.6$ km at $1.4 M_\odot$

Compilation of All Constraints



- $2M_\odot \rightarrow$ stiff EoS
- Direct Urca $\rightarrow$ soft EoS
- Tidal Deformability $\rightarrow$ soft EoS

Back to Symmetric-Matter Phase Diagram



- Critical Point at low temperature or even absent!


Conclusions

Parity doubling yields non-trivial implications for the physics of neutron stars:


- $2M_{\odot}$ with **chirally restored** but **confined** core
- High-mass stars $\rightarrow$ **not necessarily** signal of **deconfinement**
- Parity doubling $\rightarrow$ **modification** of direct URCA threshold
 - new estimate for the proton fraction threshold
 - impact on neutron star cooling
- Phase diagram with CEP at rather low temperature

Thank you for your attention

Parity Doubling for Light Baryons Aarts et al, arXiv:1710.08294 (2017)



Parity Doubling for Light Baryons Aarts et al, arXiv:1710.08294 (2017)



- Naive and **mirror** assignments under $SU(2)_L \times SU(2)_R$

$$\mathcal{L}_N = i\bar{\psi}_1 \not{\partial} \psi_1 + i\bar{\psi}_2 \not{\partial} \psi_2 + \mathbf{m}_0 \left(\bar{\psi}_1 \gamma_5 \psi_2 - \bar{\psi}_2 \gamma_5 \psi_1 \right)$$

For finite **m**₀, chiral symmetry is

- explicitly broken under naive assignment
 - remains unbroken under **mirror** assignment
- Parity doublet model for cold and dense nuclear matter

Hatsuda, Prakash, Phys.Lett. B **224** (1989)

Zschesche et al, Phys. Rev. C **75**, 055202 (2007)

$$\mathcal{L} = \mathcal{L}_N + \mathcal{L}_M + \sum_{k=1,2} g_k \bar{\psi}_k (\sigma \pm i\gamma_5 \boldsymbol{\tau} \cdot \boldsymbol{\pi}) \psi_k - g_\omega \bar{\psi}_k \psi_k \psi_k$$

- Fermions coupled to bosons: σ, π, ω
- $\mathcal{L}_M \rightarrow$ Linear σ -model

Full HQMN model Lagrangian

$$\blacksquare \mathcal{L} = \mathcal{L}_N + \mathcal{L}_M + \mathcal{L}_q$$

$$\begin{aligned} \mathcal{L}_N = & \sum_{k=1,2} \bar{\psi}_k i \not{\partial} \psi_k + m_0 (\bar{\psi}_2 \gamma_5 \psi_1 - \bar{\psi}_1 \gamma_5 \psi_2) \\ & + \sum_{k=1,2} g_k \bar{\psi}_k (\sigma \pm i \gamma_5 \boldsymbol{\tau} \cdot \boldsymbol{\pi}) \psi_k - g_\omega \bar{\psi}_k \psi_k \psi_k \end{aligned}$$

$$\mathcal{L}_q = \bar{q} i \not{\partial} q + g_q \bar{q} (\sigma + i \gamma_5 \boldsymbol{\tau} \cdot \boldsymbol{\pi}) q$$


$$\mathcal{L}_M = \frac{1}{2} (\partial_\mu \sigma)^2 + \frac{1}{2} (\partial_\mu \boldsymbol{\pi})^2 - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - V_\sigma - V_\omega - V_b$$

$$V_\sigma = -\frac{1}{2} \bar{\mu}^2 (\sigma^2 + \boldsymbol{\pi}^2) + \frac{\lambda}{4} (\sigma^2 + \boldsymbol{\pi}^2)^2 - \epsilon \sigma$$


$$V_\omega = -\frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu$$

$$V_b = -\frac{1}{2} \kappa_b^2 b^2 + \frac{1}{4} \lambda_b b^4$$


mass-density




mass-radius




mass-radius




mass-radius




Matter composition ($\alpha b_0 = 350$ MeV)




Matter composition ($\alpha b_0 = 370$ MeV)




Matter composition ($\alpha b_0 = 400$ MeV)




Matter composition ($\alpha b_0 = 450$ MeV)




Phase diagram in $(T - \mu_B)$ -plane ($\alpha b_0 = 310$ MeV)




Threshold for direct URCA







Mean fields at $T = 10$ MeV ($\alpha b_0 = 300$ MeV)



Equation of state at $T = 10$ MeV ($\alpha b_0 = 300$ MeV)



Masses at $T = 10$ MeV ($\alpha b_0 = 300$ MeV)



Masses at $T = 10$ MeV ($\alpha b_0 = 390$ MeV)

