

# Laser-driven X-ray sources for investigating extreme states of matter generated by intense heavy ion beams

*Monday, 27 January 2020 17:00 (25 minutes)*

One of the unique features of the infrastructure and facilities at GSI is the possibility to carry out experiments combining the heavy-ion beam of the accelerator with the high-power laser PHELIX. With the new beamline guiding the long-pulse (nanosecond) beam of PHELIX to the HHT experimental cave, which is currently being installed at GSI, new diagnostic methods for the investigation of heavy-ion heated states of matter will become available in the near future. These include using laser-driven X-rays for diffraction, imaging or spectroscopy for investigating the behavior of matter under such conditions. After a brief report on the status of the beamline we will discuss our plans for first experiments using the new capabilities at HHT. We have also performed a preliminary study into characterizing the laser-driven X-ray source in order to experimentally confirm the feasibility of our plans, which we will also present here.

**Primary author:** SLATTERY-MAJOR, Zsuzsanna (GSI, Darmstadt)

**Co-authors:** Dr ZIELBAUER, Bernhard (GSI Helmholtzzentrum für Schwerionenforschung GmbH); Dr NEU-MAYER, Paul (GSI, Darmstadt); EISENBARTH, Udo (GSI, Darmstadt); Dr BAGNOUD, Vincent (GSI, Darmstadt)

**Presenter:** SLATTERY-MAJOR, Zsuzsanna (GSI, Darmstadt)

**Session Classification:** Activities of HED@FAIR