

Holographic Flavor Transport

Andy O'Bannon
Max Planck Institute for Physics
Munich, Germany

String Theory and Extreme Matter
EMMI Workshop
Heidelberg, Germany

Credits

- 0705.3870 A. Karch and A. O'B.
- 0708.1994 A. O'B.
- 0808.1115 A. O'B.
- 0812.3629 A. Karch, A. O'B., E. Thompson
- 0908.2625 M. Ammon, H.T. Ngo, A. O'B.

Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

Motivation

REAL Strongly-coupled Systems

Quantum Chromodynamics (QCD)

Relativistic Heavy-Ion Collider (RHIC)

QCD at $T \leq 2 \times T_c$

$T_c \approx 170$ MeV

Strongly-coupled, Nearly-ideal FLUID

Motivation

Well-described by hydrodynamics

Inputs for hydrodynamics:

I. Thermodynamics (equation of state)

II. Transport Coefficients

η ζ D σ

Question:

Can we compute

TRANSPORT COEFFICIENTS

for QCD

at RHIC temperatures?

Question:

Can we compute

TRANSPORT COEFFICIENTS

for QCD

at RHIC temperatures?

Answer:

NO.

Philosophy:
CHANGE the Question

Question:

Can we find **ANY**

STRONGLY-COUPLED

system for which we **CAN** compute

TRANSPORT COEFFICIENTS?

Answer:
YES!

$\mathcal{N} = 4$ supersymmetric $SU(N_c)$ Yang-Mills (SYM)

$$N_c \rightarrow \infty$$

$$\lambda \rightarrow \infty$$

$$\lambda = g_{YM}^2 N_c$$

Use Gauge-gravity Duality

Shear Viscosity

$$\frac{\eta}{s} = \frac{\hbar}{4\pi k_B}$$

GOAL

Compute a
CONDUCTIVITY
associated with
“Quarks” or “Electrons”
using
Gauge-gravity Duality

RESULT

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

Current $\langle J^x \rangle = \sigma E$ nonlinear in E

Pair Production

Drude Conductivity

Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

The Theory

$\mathcal{N} = 4$ supersymmetric $SU(N_c)$ Yang-Mills (SYM)

$$N_c \rightarrow \infty$$

$$\lambda \rightarrow \infty$$

$$\beta = 0$$

The Theory

$\mathcal{N} = 4$ supersymmetric $SU(N_c)$ Yang-Mills (SYM)

$$N_c \rightarrow \infty$$

$$\lambda \rightarrow \infty$$

No Quarks!

ADD

$N_f \mathcal{N} = 2$ hypermultiplets

$$\beta = +\mathcal{O}\left(\frac{N_f}{N_c}\right)$$

“Probe Limit”

$$N_f \text{ fixed} \ll N_c \rightarrow \infty$$

Scales

Temperature

T

Mass

m

$U(1)_B$

symmetry current

J^μ

Baryon Number Density

$\langle J^t \rangle$


Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

Electric Field E

“Two-fluid” picture

Lorentz force = Drag Force



$$\langle J^x \rangle = \sigma E$$


Steady-state???

Translation Invariance



Momentum
Conservation

Net Charge
+ Constant Electric
Field



Net Work

ENTIRE SYSTEM
ACCELERATES FOREVER

NO

DISSIPATION!

Probe Limit MIMICS Dissipation

$$\langle T_{\mu\nu} \rangle = \mathcal{O}(N_c^2)_{\mu\nu} + \mathcal{O}(N_f N_c)_{\mu\nu}$$

$$\partial_\mu \langle T^{\mu\nu} \rangle = F^{\nu\sigma} \langle J_\sigma \rangle$$

$$\partial_t \langle T^{tt} \rangle = E \langle J_x \rangle \quad \partial_t \langle T^{tx} \rangle = -E \langle J_t \rangle$$

$$\langle J_\mu \rangle = \mathcal{O}(N_f N_c)$$

$$E = \mathcal{O}(1)$$

Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

The Gravity Dual

$\mathcal{N} = 4$ SYM
 $N_c, \lambda \rightarrow \infty$

=

Supergravity
 $AdS_5 \times S^5$

Finite temperature

=

AdS-Schwarzschild

N_f $\mathcal{N} = 2$ hypers.

=

N_f probe D7-branes
 $AdS_5 \times S^3$

m

=

Embedding

J^μ

=

A_μ

Anti-de Sitter Space

$$ds_{AdS_5}^2 = g_{rr}(r)dr^2 + g_{tt}(r)dt^2 + g_{xx}(r)d\vec{x}^2$$

Boundary

$$r \rightarrow \infty$$

$$g_{rr} \sim 1/r^2 \quad |g_{tt}| \sim g_{xx} \sim r^2$$

Horizon

$$g_{tt}(r_H) = 0 \quad r_H = \pi T R^2$$

$$ds_{S^5}^2 = d\theta^2 + \sin^2 \theta ds_{S^1}^2 + \cos^2 \theta ds_{S^3}^2$$

The Gravity Dual

$\mathcal{N} = 4$ SYM
 $N_c, \lambda \rightarrow \infty$

=

Supergravity
 $AdS_5 \times S^5$

Finite temperature

=

AdS-Schwarzschild

N_f $\mathcal{N} = 2$ hypers.

=

N_f probe D7-branes
 $AdS_5 \times S^3$

m

=

Embedding

J^μ

=

A_μ

D7-Brane Action

$$S_{D7} = -N_f T_{D7} \int d^8 x \sqrt{-\det (g_{ab}^{D7} + (2\pi\alpha') F_{ab})}$$



Extended along $AdS_5 \times S^3$

Can move along θ and S^1 directions

Ansatz: $\theta(r)$

$$ds_{D7}^2 = (g_{rr} + \theta'(r)^2) dr^2 + g_{tt} dt^2 + g_{xx} d\vec{x}^2 + \cos^2 \theta(r) ds_{S^3}^2$$

Boundary Conditions



$$\theta(r_H) \in \left[0, \frac{\pi}{2}\right] \quad \theta'(r_H) = 0$$

The Gravity Dual

$\mathcal{N} = 4$ SYM
 $N_c, \lambda \rightarrow \infty$

=

Supergravity
 $AdS_5 \times S^5$

Finite temperature

=

AdS-Schwarzschild

N_f $\mathcal{N} = 2$ hypers.

=

N_f probe D7-branes
 $AdS_5 \times S^3$

m

=

Embedding

J^μ

=

A_μ

Dual Operators

$$\mathcal{O}_m = \frac{\partial}{\partial m} \mathcal{L} = \bar{\psi}\psi + (\text{scalars})$$

$$\theta(r) = \frac{m}{r} + \frac{\langle \mathcal{O}_m \rangle}{r^3} + \dots$$

$$\theta(r_H) = 0 \quad \Rightarrow \quad m = 0$$

$$\theta(r_H) \rightarrow \pi/2 \quad \Rightarrow \quad m \rightarrow \infty$$

The Gravity Dual

$\mathcal{N} = 4$ SYM
 $N_c, \lambda \rightarrow \infty$

=

Supergravity
 $AdS_5 \times S^5$

Finite temperature

=

AdS-Schwarzschild

N_f $\mathcal{N} = 2$ hypers.

=

N_f probe D7-branes
 $AdS_5 \times S^3$

m

=

Embedding

J^μ

=

A_μ

Dual Operators

We want

$$\mu_B \quad \langle J^t \rangle$$

$$A_t(r) = \mu_B + \frac{\langle J^t \rangle}{r^2} + \dots$$

We also want

$$E \quad \langle J^x \rangle$$

$$A_x(t, r) = -Et + \frac{\langle J^x \rangle}{r^2} + \dots$$

Calculating the Conductivity

$$S_{D7} \sim -N_f T_{D7} \int_{r_H}^{\infty} dr g_{xx} \cos^3 \theta(r)$$

$$\sqrt{|g_{tt}|g_{xx} (g_{rr} + \theta'(r)^2) - (2\pi\alpha')^2 \left(g_{xx} A_t'^2 + g_{tt} A_x'^2 + (g_{rr} + \theta'(r)^2) \dot{A}_x'^2 \right)}$$

$$\partial_r \frac{\delta S_{D7}}{\delta A'_\mu} = 0 \quad \frac{\delta S_{D7}}{\delta A'_\mu} = \langle J^\mu \rangle$$

Solve for

$$A'_t \quad A'_x$$

in terms of

$$\begin{array}{cc} \langle J^t \rangle & \langle J^x \rangle \\ E & \theta(r) \end{array}$$

Calculating the Conductivity

Insert the solutions back into the action:

$$S_{D7} \sim - \int_{r_H}^{\infty} dr g_{xx}^{5/2} |g_{tt}|^{1/2} (g_{rr} + \theta'(r)^2)^{1/2} \cos^6 \theta(r)$$

$$\sqrt{\frac{|g_{tt}|g_{xx} - (2\pi\alpha')^2 E^2}{c^2 |g_{tt}|g_{xx}^3 \cos^6 \theta(r) + |g_{tt}|\langle J^t \rangle^2 - g_{xx}\langle J^x \rangle^2}}$$

$$c^2 = \frac{\lambda N_f^2 N_c^2}{(2\pi)^6}$$

Calculating the Conductivity

Insert the solutions back into the action:


$$S_{D7} \sim - \int_{r_H}^{\infty} dr g_{xx}^{5/2} |g_{tt}|^{1/2} (g_{rr} + \theta'(r)^2)^{1/2} \cos^6 \theta(r)$$

$$\sqrt{\frac{|g_{tt}|g_{xx} - (2\pi\alpha')^2 E^2}{c^2 |g_{tt}|g_{xx}^3 \cos^6 \theta(r) + |g_{tt}|\langle J^t \rangle^2 - g_{xx}\langle J^x \rangle^2}}$$

Numerator changes sign!

Calculating the Conductivity

Imaginary Action



INSTABILITY

Minkowski Space

$$\sqrt{1 - (2\pi\alpha')^2 E^2}$$

$$E > \frac{1}{2\pi\alpha'}$$

Strings ripped apart!

AdS-Schwazrschild

$$\sqrt{|g_{tt}|g_{xx} - (2\pi\alpha')^2 E^2}$$

Calculating the Conductivity

$$\sqrt{\frac{|g_{tt}|g_{xx} - (2\pi\alpha')^2 E^2}{c^2 |g_{tt}|g_{xx}^3 \cos^6 \theta(r) + |g_{tt}|\langle J^t \rangle^2 - g_{xx}\langle J^x \rangle^2}}$$

$$\left[|g_{tt}|g_{xx} - (2\pi\alpha')^2 E^2 \right]_{r_*} = 0$$

To AVOID the instability: DEMAND

$$\left[c^2 |g_{tt}|g_{xx}^3 \cos^6 \theta(r) + |g_{tt}|\langle J^t \rangle^2 - g_{xx}\langle J^x \rangle^2 \right]_{r_*} = 0$$

Solve for

$$\langle J^x \rangle = \sigma E$$

Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

$$e = \frac{E}{\frac{\pi}{2} \sqrt{\lambda} T^2}$$

$$d = \frac{\langle J^t \rangle}{\frac{\pi}{2} \sqrt{\lambda} T^2}$$

$$f(m) = \cos^6 \theta(r_*)$$

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

Depends on E !

$$\langle J^x \rangle = \sigma(E)E$$

Linearize in E

$$\langle J^x \rangle = \sigma(0)E$$

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

$$\langle J^t \rangle = 0 \quad \text{BUT} \quad \sigma \neq 0$$

Pair Production

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

$$m \rightarrow \infty \Rightarrow f(m) \rightarrow 0$$

$$m \rightarrow 0 \Rightarrow f(m) \rightarrow 1$$

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

$$\langle T^{tx} \rangle \propto \langle J^t \rangle$$

NO momentum flow at **zero density**

The Conductivity

$$\sigma = \sqrt{\frac{N_f^2 N_c^2}{16\pi^2} T^2 \sqrt{e^2 + 1} f(m) + \frac{d^2}{e^2 + 1}}$$

Drude Conductivity

Linearize in E

$\sigma(0)$

Take

$m \rightarrow \infty$

$$\sigma \rightarrow d = \frac{\langle J^t \rangle}{\frac{\pi}{2} \sqrt{\lambda} T^2}$$

Why $m \rightarrow \infty$?

Charges behave as semi-classical quasi-particles:

$$\frac{dp}{dt} = -\mu p + E$$

Separate calculation

$$\mu m = \frac{\pi}{2} \sqrt{\lambda T^2}$$

$$\mu m = \frac{\pi}{2} \sqrt{\lambda T^2}$$

$$\sigma \rightarrow d = \frac{\langle J^t \rangle}{\frac{\pi}{2} \sqrt{\lambda T^2}} = \frac{\langle J^t \rangle}{\mu m}$$

Drude Conductivity

Outline:

- Motivation
- The Theory
- The Conductivity I
- The Gravity Dual
- The Conductivity II
- Summary and Outlook

Summary + Outlook

Probe Branes are Great!

Computed CONDUCTIVITY

for a “DISSIPATIVE”

STRONGLY-COUPLED

Non-Abelian Gauge Theory

FUTURE DIRECTIONS

MORE TRANSPORT COEFFICIENTS:

Magnetic Fields

Anomalous currents

Condensed Matter Applications:

Thermo-electric Transport

Quantum Hall Effect

Non-relativistic Theories

Superfluidity and Fermi Surfaces

$$N_f = 2 \implies U(2) = U(1) \times SU(2)$$

$$J_a^\mu \quad a = 1, 2, 3$$


Chemical potential for

Ground state: nonzero





$$\langle J_1^x \rangle$$

p-wave superfluid!

$$U(1) \subset SU(2) \quad \langle J_3^t \rangle$$



What does the Fermi Surface look like?

T_c  $0.91T_c$  $0.69T_c$  $0.43T_c$ 

Thank You.